Advances in Physics Theories and Applications
ISSN 2224-719X (Paper) ISSN 2225-0638 (Online)
Vol 14, 2013

Ultra-Secure Internet, Bell's Inequalities, Entanglement Distillation, Bizarre Quantum Physics And Encryption, Antibunching Dip In The Resonance Fluorescence, "Of The Self-Overcoming", Charginos, Neutralinos, Sleptons And Supersymmetric Standard Model, Embodiment And Expression Of Human Values Is Will To Power, Wormhole Between 'Entangled' Quantum Particles, And Other Reports Of Similar Interest: Das Ganze Ist Mehr Als Die Summe Seiner Telle: Whole Is Greater Than Sum Of Its Parts: Lopado--Temacho--Selacho--Galeo--Kranio--Leipsano--Drim--Hypo--Trimmato--Silphio--Parao--Melito--Katakechy--Meno--Kichl--Epi--Kossypho--Phatto--Perister--Alektryon--Opte--Kephallio--Kigklo--Peleio--Lagoio--Siraio--Baphe--Tragano--Pterygon Models: Pot Pourri, Olio Podrida Of A Various Dishes Models

@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@

[1]Dr. K.N.P.Kumar

Post Doctoral Scholar, Department of Mathematics, Kuvempu University, Janna Sahyadri, Shankaraghatta, Karnataka, India

[2]Prof. C.J.Prabhakara

Chairman and Professor Department of Computer Science, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

[2]Prof. S.K. Narasimha Murthy

Chairman and Professor Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

[3]Dr. B.J.Gireesha

Assistant Professor Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@

Abstract: Nature's General ledger continues to be written. Apprehensional - argumentation, bounds-comprehension, dialectics-discernment, generalizational induction, inferential intellect, soundness-speculation,

expositional ground, cerebrate cogitation, expostulational justification, determinative- development, argumentational assertion, polemical proof, innervation inspirit, congruity- consistency, adaptational compliance, composition-compromise, adjustment-affiliation, consistency-correspondence, accord-coetaneousness, coevality-coincidence, concurrence-conformity, conjunction-contemporaneousness, connection-consistency of the following systems are studied. Homologous interchangeability, undifferentiated homologosness, coextensive comparability, correlative correspondence or for that matter incongruous- inconsistency, inexorable-inflexibility, inharmonious-intransigence, Distinctiveness oneness, of the systems are also identified for concatenation and a statement of a holistic model, notwithstanding the fact that the model is posted and letters are given in piecemeal of nine modules for felicity of understanding and typewriting problems what with hundreds of suffixes. Following are the systems taken in to consideration for consummation, consolidation, corporation, consubstantiation and concatenation: Shiva = Pure Consciousness (Here All Dualities Cease: See Earlier Models. Being= Becoming, Self=Nonself Order = Chaos Etc.,); Shakti =Impure Consciousness (Mother Divine: Responsible For The Phenomenal Manifestation), Vijnāna-Bhairava Tantra, Language As (=) The Sound (Nāda) Integral To Shiva's (Recall Assets=Liabilities Example Creative Power, Pure Consciousness, (U-A Theories) Interpret Their Central Myth Of Shiva's Emanation And Control Of The Universe Through Shakti As Itself An Act Of Self-Recognition (Ahampratyavamarsha, Pratyabhijnā), Prakāsha Is The "Bare Subjective Awareness" That Validates Each Cognition, So That One Knows That One Knows, Everything Is Consciousness And Consciousness Is Everything, "Semantic Exclusion" (Apoha) Nonepistemic "Coordination" (Sārūpya) Between Languages And Momentary Perceptual Data As The Basis For Successful Reference In Communication And Behaviors, Pratyabhijnā Ontology: The Syntax Of Empowered Identity, Shiva's Mythic Action Through Shakti As Self-Recognition That Constitutes (Eb) All Experience And Objects Of Experience, "Being Is The Agency Of The Act Of Becoming, Which Is, The Freedom Characteristic Of An Agent Regarding All Actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, And 1:258-59).", Parā, Parā-Aparā And Aparā Śakti, Pure Consciousness (Cessation Of Dualities: Initial Conditions For The Creation Of Universe) And Individual Consciousness (Individual General Ledger), Anuttara, Svatantrya, Self-Created Free Will, Function Of Svātantrya Is Self Reflection And Contemplation Of One's Nature, Spanda, Ultra-Secure Internet, Multipartite Einstein-Podolsky-Rosen Steering, Einstein-Podolsky-Rosen Steering And Multipartite Quantum Nonlocality, Entanglement Distillation, Bizarre Quantum Physics And Encryption, Antibunching Dip In The Resonance Fluorescence, Principle Of Locality, Local Realism, Mind-Independent Property Does Not Have To Be The Value Of Some Physical Variable Such As Position Or Momentum, "Detection Loophole" Or The "Communication Loophole.", No Physical Theory Which Is Realistic And Also Local In A Specified Sense Can Agree With All Of The Statistical Implications Of Quantum Mechanics, Guidance Equation Ensures Agreement With The Statistical Predictions Of Standard Quantum Mechanics, Existence Of Such Mappings For Which These Intuitive Interpretations Are Maintained And Conditions (1) – (4) (See Previous Module And Extant One) Are Satisfied, Truth Value Which M Assigns To A Subspace S Depends Upon The Context C Of Propositions Measured In Tandem With The One Associated With S, Outcome Of A Measurement Made On One Constituent Depends Upon The Action Of The "Guiding Wave" Upon Constituents That Are Far Off, Which In General Will Depend On The Measurement Arrangement On That Side, Violation Constitutes A Special Case Of Bell's Theorem For It Shows That No Local Realistic Theory Subsumed Under The Framework Of Bell's 1964 Paper Can Agree With All Of The Statistical Predictions Of Quantum Theory, Bell's Own Version Of His Theorem Assumed That S And T Are Both Bivalent, Remote Context Independence , "Bell-Clauser-Horne-Shimony-Holt (BCHHSH) Inequality.", Local Realistic Theory, Prob Φ (1,−1|a,b) = probΦ(−1,1|a,b) = (½)sin2σ. Maximum Amount Of Violation Is Achieved With A Quantum State Of Maximum Degree Of Entanglement, Bell's Theorem Shows That Such An Ascription (Entangled Quantum State) Will Have Statistical Implications In Disagreement With Those Of Quantum Mechanics, Advantages And Disadvantages Of The Experiment, "Fair Sampling Assumption", − (¼) ≤ [D(π/8)/D0 − D(3π/8)/D0] ≤ ¼ :Consequence Of The BCHSH Inequality, And In Disagreement With The Quantum Mechanical Prediction, Variant Of The Holt-Pipkin Experiment, Detection Loophole And Its Remedy, Local Realistic Theories Can Be Consistent With The Quantum Mechanical Predictions Provided That The Detectors Are Properly Selective, BCH Inequality, BCH Inequality In Principle This Inequality Could Be Used To Adjudicate Between The Family Of Local Realistic Theories And Quantum Mechanics, Dissociation Is A Two-Body Process,

"No Enhancement" Assumption, Communication Loophole And Its Remedy, Maintenance Of Entanglement Between Particles Of A Pair That Are Well Separated, Entanglement Is A Property That May Dwindle With Spatial Separation, "Time Tag" Attached To Each Detected Photon, Quantum Non-Separability (Aspect 1999), Some Variants Of Bell's Theorem, No Deterministic Local Realistic Theory Can Agree With All The Predictions Of Quantum Mechanics, Incompatibility Of A Deterministic Local Realistic Theory With Quantum Mechanics, Quantum Mechanics Is Non-Local, Ways In Which Habits Are Formed Attests To The Desires At The Heart Of Our Social Milieu, Locality Cannot Be Married To The Assignment Of Determinate Values In The Recommended Way, Violations Of Locality Are The Result Of Conspiracy Plotted In The Overlap Of The Backward Light Cones Of The Analysis-Detection Events In The 1 And 2. Stability Solutions and other phenomenological correlates follow.

Key words: Truth Value Which M Assigns To A Subspace S Depends Upon The Context C Of Propositions Measured In Tandem With The One Associated With S, Outcome Of A Measurement Made On One Constituent Depends Upon The Action Of The "Guiding Wave" Upon Constituents That Are Far Off, Which In General Will Depend On The Measurement Arrangement On That Side, Violation Constitutes A Special Case Of Bell's Theorem For It Shows That No Local Realistic Theory Subsumed Under The Framework Of Bell's 1964 Paper Can Agree With All Of The Statistical Predictions Of Quantum Theory, Bell's Own Version Of His Theorem Assumed That S And T Are Both Bivalent, Remote Context Independence , "Bell-Clauser-Horne-Shimony-Holt (BCHHSH) Inequality.", Local Realistic Theory, Prob Φ (1,−1|a,b) = probΦ(−1,1|a,b) = (½)sin2σ. Maximum Amount Of Violation Is Achieved With A Quantum State Of Maximum Degree Of Entanglement, Bell's Theorem Shows That Such An Ascription (Entangled Quantum State) Will Have Statistical Implications In Disagreement With Those Of Quantum Mechanics, Advantages And Disadvantages Of The Experiment, "Fair Sampling Assumption", − (¼) ≤ [D(π/8)/D0 − D(3π/8)/D0] ≤ ¼ :Consequence Of The BCHSH Inequality, And In Disagreement With The Quantum Mechanical Prediction, Variant Of The Holt-Pipkin Experiment, Detection Loophole And Its Remedy, Local Realistic Theories Can Be Consistent With The Quantum Mechanical Predictions Provided That The Detectors Are Properly Selective, BCH Inequality, BCH Inequality In Principle This Inequality Could Be Used To Adjudicate Between The Family Of Local Realistic Theories And Quantum Mechanics, Dissociation Is A Two-Body Process, "No Enhancement" Assumption, Communication Loophole And Its Remedy, Maintenance Of Entanglement Between Particles Of A Pair That Are Well Separated

@@@

Fundamental constituents of reality are strings of the Planck length (about 10−33 cm) which vibrate at resonant frequencies. Every string, in theory, has a unique resonance, or harmonic. Different harmonics determine different fundamental particles. The tension in a string is on the order of the Planck force (1044 newtons). The graviton (the proposed messenger particle of the gravitational force), for example, is predicted by the theory to be a string with wave amplitude zero (Wikipedia). With that statement string theorists' are making an insinuation that the fundamental reality is absolute subjectivity, one that is filled with raga and Dwesha, two main elements responsible for pathology of the mind. Verily we are circumscribed by a holographic duality transcending which is the only avowed objective of the human being

Those in full consciousness of Terrestial Brahman-Anti Brahman to be the governing principle of the material manifestation can understand and know Brahman-AntiBrahman at the time of death caused by Brahman-Anti Brahman for he has become one with Brahman Anti Brahman, the essence of Principe of system disintegration.

Empirical subjectivity is (=) constituted in the mind under the influence of (e&eb) the principles affecting it; the mind therefore does not (e) have the characteristics of a preexisting subject. (ES 29) These associations account not only for (e) experience in the basic sense, but up to the highest level of social and cultural life: this is the basis for Hume's rejection (e) of a social contract model of society (such as Hobbes'), in favour of convention alone. Morals, feelings, bodily comportment, all of these elements of subjectivity are explained, not by (e) transcendental structures, such as Kant will propose, but by (e) the immanent activity of association. Once this habitual structure of the self is (=) in place, Deleuze suggests, the Humean concept of belief comes into play (eb), which is resolutely a central part of human nature. It describes (eb) the particularly human way of going beyond (e) the given. When we expect the sun to come up tomorrow, we do not do so because (e) we know that it will, but because of a belief based (e) on a habit. This in turn reverses (e&eb) the hierarchy of knowledge and belief, and results, for Deleuze, in a, "great conversion (e&eb) of theory to practice." (PI 36) Every act of belief is (=) a practical application of habit, without (e) any reference to an a priori ability to judge. Not only is (=) the human being thus habitual, on Deleuze's reading, but also creative (eb) even in the most mundane moments of life. Finally, Deleuze insists that one of Hume's greatest contributions to modern philosophy is (=) his insistence that all relations are (=) external to their terms: this is the essence of Hume's anti-transcendental stance. Human nature cannot (e) unite itself, there is no 'I' which stands before experience, but only moments of experience themselves, unattached and meaningless without (e) any necessary relation to each other. A **flash of red, a movement, a gust of wind, dashing of a woman against you, obscene gestures, spitting at you, talking extempore about you (my addition) and these elements must be externally related (e&eb) to each other to create (eb) the sensation of a tree in autumn. In the social world, this externality attests (eb) to the always-already interested nature of life: no relation is necessary, or governed by neutral laws, so every relation has (e) a localised and passional motive. The ways in which habits are formed attests (eb) to the desires at the heart of our social milieu. Subjectivity, as Deleuze describes it through his reading of Hume, is (=) a practical, passional, empiricist concept, immediately located at the heart of the conventional, which is to say the social.**

Life is not unfair; Do not get used to unfairness; donot stay foolish or stay stupid expand your individual consciousness; Terrestial Brahman-AntiBrahman is responsible for all such acts of omissions and commissions; there is no reason why you have to be satisfied and satiated with a promotion while the other is surfeited with a trillion dollars

Poverty penury and perdition and despondency is also business; in their name Brahman Anti Brahman get fish and loaves only to give pittance to them without any compunction and contrition gormandizing the lion's share of the manna from heaven

Energy of Terrestial Brahman-AntiBrahman comprises of various forms like Kama, krodha, lobha, moha, mada, mastarya is difficult to overcome and all people fall prey to it. Those who surrender to Energy of Terrestial Brahman-AntiBrahman shall be freed of all their sins and court cases by the virtue of and connections of Energy of Terrestial Brahman-AntiBrahman

Deleuze describes three different levels of time within (eb) which repetition occurs. Deleuze takes as axiomatic the notion that there is (=) no time but the present, which contains (e) past and future. These layers describe different ways in which past and future can be inscribed in (eb) a present. As this inscription grows more complicated, the status of the present itself becomes (=) more abstract. Basic processes of the universe have (e) a momentum that they carry into each present moment. A 'contraction' (e) of reality refers to the collection of a diffuse ongoing force into the present. Prior thought and behavior, all substance performs (eb) contraction. "We are made of (e) contracted water, earth, light, and air...Every organism, in its receptive and perceptual elements, but also in its viscera, is (=) a sum of contractions, of retentions and expectations" (73). Passive synthesis is exemplified by (e) habit. First the association was established between colors and the status quo or status quo ante of propositions that I was interested in. Like Scientists or actors or for that matter anyone, repetition forms the core essence of Brahman-AntiBrahman. Later on I started associating with them as a habit. Albeit repetition is done many times I have "self overcome "such feelings, which many people would not be able to. Habit incarnates (eb) the past (and gestures to the future) in the present by transforming the weight of experience into (e&eb) urgency. Habit creates (eb) a multitude of "larval selves," each of which functions like a small ego with (e) desires and satisfactions. In Freudian discourse, this is (=) the domain of bound excitations associated with the pleasure

principle. Deleuze cites Hume and Bergson as relevant to his understanding of the passive synthesis. The second level of time is (=) organized by the active force of memory, which introduces (eb) discontinuity into the passage of time by sustaining (eb) relationships between more distant events. A discussion of destiny makes clear how memory transforms time and enacts a more profound form of repetition: **Destiny never consists (e&eb) in step-by-step deterministic relations between presents which succeed one another according to the order of a represented time. Rather, it implies (eb) between successive presents non-localisable connections, actions at a distance, systems of replay, resonance and echoes, objective chances, signs, signals, and roles which transcend spatial locations and temporal successions. (83). Like one might have an insinuation and innuendo that he shall make it as an actor , or a scientist, or a top administrator ,as it happened in my case.** Relative to the passive synthesis of habit, memory is (=) virtual and vertical. It deals with events in their depth and structure rather than in (eb) their contiguity in time. Structure is important point here. Criminal intimidation, death threats, physical assaults, throwing of rats with entrails drawn out in the vicinity, throwing of human excretea changed the structure of the portending things time to time of the events. Where passive syntheses created a field of 'me's,' active synthesis is (=) performed by 'I.' In the Freudian register, this synthesis describes the displaced energy of Eros, which becomes (=) a searching and problematizing force rather than a simple stimulus to gratification. As already mentioned, problematizing human events is one of the ways of Terrestial Brahman Anti Brahman, depending on which the next action is taken. Although I fooled, I paid a heavy price for the extreme pressure and Problematization. I had heart attack. Like all of us Terrestial Brahman Anti Brahman lacks extraordinary ideas and resorts to same tactics. Just any one can be made to do anything either with money or intimidation or some other favours which may run in to billions of dollars. Very rarely person with Rajas, tamas and Sattava can circumvent such a situation. The third layer of time still exists in the present, but it does so in a way that breaks free (e) from the simple repetition of time. This level refers to an ultimate event so powerful that it becomes (=) omnipresent. It is a great symbolic event, like the murder to be committed by Oedipus or Hamlet. Upon rising to this level, an actor effaces herself as such and joins the abstract realm of eternal return. **The me and the I give way to "the man without name, without family, without qualities, without self or I...the already-Overman whose scattered members gravitate around the sublime image" (90). The man without name is Terrestial Brahman –Anti Brahman. I donot know whether Celestial Brahman Anti Brahman has any hand in this (cosmic general ledger: cosmic consciousness May be it has due to belief systems and or lack of it which shall confront you with more pronounced situation.)** Empty time is associated (e&eb) with Thanatos, a desexualized energy that runs through all matter and supersedes the particularity of an individual psychic system. Deleuze is careful to point out that there is no reason for Thanatos to produce (eb) a specifically destructive impulse or 'death instinct' in the subject; he conceives of Thanatos as simply indifferent. In Problematization of human events such a possibility of Thanatos being associated with the death instinct or suicidal tendencies doth exist. In fact it will as I thought many a time. **Repetition for**

Itself (Wikipedia) some additions and interpretations are mine. It has got nothing to with Deleuzean interpretation

Terrestial Brahman Anti Brahman controls both the field of activities and the finite enjoyer; This is what perhaps Adman smith meant by "invisible hand" and Hobbes Bellum Omnium Contra omnes

Intelligence of Terrestial Brahman Anti Brahman arises from the stupid "moon" theorems we have in jurisprudence.

While tantrism is a complex and controversial subject, one of its most definitive characteristics for contemporary classifications—if not its most definitive one—is the pursuit of power. Tantric traditions are (=) thus those that aim at increasing (eb) the power of the practitioner. The theological designation for the essence of such power is (=) Shakti (the female counterpart to the male divine principle, whose **essence is power**). The manifestations of Shakti that the practitioner of tantra aspire after vary greatly, from relatively limited magical proficiencies (siddhis or vibhūtis), through royal power, to (eb) the deindividualized and liberated saint's omnipotence to the performance of God's cosmic acts. What is God or Para Brahman's act? What is Karma? Take my case for example. I was a professional banker who took VRS, an aspiring star, and an academician who wanted to just make it to the top irrespective of the fact whether I had the abilities and capacities. I got in to a Chakravyuha like Abhimanyu, and lead a life of trials and tribulations and travails for a decade. What is that I did to be punished for this or still being punished? I will assume that God (Parà Brahman will take in to account what I had done in my earlier incarnations (I donot think I did anything like that in this incarnation) wherein I could have tormented or produced a sense of deprivation balance either equivalent to or less than what I am going through now. Or God must have added interest and given me the punishment. I will assume both and give models One with the deprivation balance equivalent to what I had done, and the other deprivation balance which includes the interest for the period between my act of commission and this incarnation. I will also take in to account the third possibility of Brahman-AntiBrahman who like spider weave the web, see the prey undergoing tormentations in the web and increase it just for "play" .In his seminal essay, "Purity and Power among the Brahmans of Kashmir," the Oxford historian Alexis Sanderson elucidates that the tantric pursuit of such power transgresses (e) orthodox, mainstream Hindu norms that delimit human agency for the sake of symbolic and ritual purity (shuddhi) (Sanderson 1985).**Violating**

prescriptions regarding caste, sexuality, diet and death, many of the tantric rites were originally performed in (eb) cremation grounds. Whereas in Shākta tantrism, Shakti as a Goddess is herself the ultimate deity, in monistic Kashmiri Shaivism she is incorporated into (eb) the metaphysical essence of the God Shiva. Shiva = pure consciousness (Here all dualities cease: See earlier models. Being= becoming, self=Nonself order = Chaos etc.,) Shakti =Impure consciousness (Mother divine: responsible for the manifestation). The basic pattern of spiritual practice, which also reflects the appropriation of Goddess worship (Shaktism) by Shaivism, is the approach to Shiva through (e&eb) Shakti. As the Shaiva scripture Vijnāna-Bhairava proclaims, Shakti is the **door**. The adept pursues the realization of identity with the omnipotent Shiva by assuming (eb) his mythic agency in emanating and controlling (e&eb) the universe through Shakti. Shakti = divided universe = Power. Thus in the sexual ritual a man realizes himself as (=) the possessor of Shakti within his partner. In more frequent internalized "theosophical" contemplations one realizes oneself as (=) the possessor of Shakti in all her immanent modalities with the aid of circular diagrams of cosmogenesis (mandalas) and mantras. An expression of this same process was (=) the production by monistic Shaiva Brahmans of increasingly systematic manuals of doctrines and practices on the model of Sanskrit scholastic texts (shāstras). This creation of what may be described as a religious mission to the educated elites also led (eb) to the increasing consolidation of the various streams of monistic Shaivism. This development began in the ninth century with Vasugupta's transmission of the manual Shiva Sūtra, ostensibly revealed to him by Shiva (Pure consciousness; cessation of all dualities space=time, order=chaos sound=silence) himself Further systematization of its teachings by either Vasugupta or his disciple Kallata in the Spanda Kārikā. These two works and their commentaries form the core texts of the "Spanda system" of monistic Shaivism, known for its interpretation of Shakti (manifested and divided universe attributable to mind: Neuron DNA see earlier models) as (=) Spanda, "cosmic pulsation." We shall take this as the pulsation or throbbing of strings as in string theory. The tradition of monistic Shaivism called "Trika" (referring to its emphasis on various triads of modalities of Shakti and cosmic levels) produced the first work of full-fledged scholastic philosophy. This was the Shiva drishti, "Cognition of Shiva," by Somānanda (c. 900-950 C.E.). (See the summary of themes of the Shiva drishti below.) Utpaladeva, a student of Somānanda, wrote a commentary on the Shiva drishti, the Shivadrishtivritti. He also wrote several other works interpreting and furthering the work of Somānanda with much greater sophistication. Those texts are the foundational works of the Pratyabhijnā philosophy of focus in this article. The most comprehensive of these texts are the Īshvarapratyabhijnākārikā, "Verses on the Recognition of the Lord," and two commentaries on the Verses, the short Īshvarapratyabhijnākārikāvritti, and the more detailed Īshvarapratyabhijnāvivriti. (The latter text has been accessible to contemporary scholars only in fragments.) Utpaladeva also wrote a trilogy of more specialized philosophical studies, the Siddhitrayī, "Three Proofs"—Īshvarasiddhi, "Proof of the Lord;" Ajadapramātrisiddhi, "Proof of a Subject who is not Insentient;" and Sambandhasiddhi, "Proof of Relation." Abhinavagupta, widely recognized as one of the greatest philosophers of South Asia, was a disciple of a disciple of Utpaladeva. Abhinava profoundly

elaborated and augmented Utpaladeva‘s arguments in long commentaries, one directly on the Verses, the Īshvarapratyabhijnāvimarshinī; and the other on Utpaladeva‘s longer auto commentary, the Īshvarapratyabhijnāvivritivimarshinī. While Abhinavagupta’s Pratyabhijnā commentaries are of paramount philosophical importance, this thinker’s greatest significance in the history of tantrism is probably his effort, in his monumental Tantrāloka and numerous other works, to systematize and provide (eb) a critical philosophical structure to non-philosophical tantric theology. Abhinava utilized categories from the Pratyabhijnā philosophy to interpret and organize (e&eb) the diverse aspects of doctrine and practice and Shaiva symbolism from the “Trika” sub-tradition; and he synthesized under the rubric of this philosophically rationalized Trika Shaivism an enormous range of symbolism and practice from other Shaiva and Shākta traditions as well. Abhinavagupta is also renowned for his works on Sanskrit poetics—in which he interpreted aesthetic experience as homologous to, and practically approaching the monistic Shaiva Soteriological realization. While the focus of this article is on Utpaladeva‘s and Abhinavagupta’s Pratyabhijnā philosophy, mention should be made of some of the basic themes of Somānanda’s precursory Shiva drishti. Somānanda’s broadest concern is to explain how Shiva through (e&eb) the various modalities of his Shakti (phenomenological manifestation of the universe) emanate (eb) a real universe that remains identical with himself.(Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger)). In establishing the Shaiva doctrine he refutes (e) a number of alternative views on ultimate reality, the self, God and the metaphysical status of the world. He devotes the greatest polemical efforts against the theories of the 4th-6th century Vaiyākarana (or “Grammarian”) philosopher Bhartrihari. According to Bhartrihari, the ultimate reality is (=) the Word Absolute (shabdabrahman)—a super-linguistic plenum, which fragments and emanates into (eb) the multiplicity of forms of expressive speech and referents of that speech. Somānanda repudiates the view that a linguistic entity could be the ultimate reality, while at the same time identifying the true source of **language as (=) the Sound (nāda) integral to Shiva’s (Recall Assets=Liabilities example creative power**; here gratification = deprivation). Somānanda takes a less polemical approach towards Shāktism. He argues that there is ultimately no difference between Shakti and Shiva, who is (=) the possessor of Shakti. This means that impure consciousness is a part of pure consciousness and emanated from (e) pure consciousness. He supports this contention with the analogy of the inseparability of heat from fire, which is the possessor of heat. Nevertheless, he asserts that it is more proper to refer to the ultimate reality as Shiva rather than Shakti. Other Hindu schools criticized by Somānanda include the Pancarātra as well as the Vedānta, Sāmkhya and Nyāya-Vaisheshika systems. Somānanda briefly adduces some considerations against the Buddhist theory of momentariness, which were directly picked up and elaborated by Utpaladeva and Abhinavagupta. The most important of these was his advertence to the experience of recognition (Pratyabhijnā) as evidence both for the continuity of

entities from the past through the present, and for the self that connects the past and (e&eb) present experiences of those entities.(Note we have taken self as witness consciousness the record keeper Neuron DNA).Somānanda's claims that synthetic categories or universals are (=) more primitive than particulars, and his invocation of Sanskrit syntax to explain Shiva's agency likewise had an important impact on Utpaladeva and Abhinavagupta. (See below.) .Also noteworthy is Somānanda's (S Theories) advocacy of a "panpsychist" theory that all things, which emanate from (e) the consciousness of Shiva, have their own consciousness and agency. Somānanda additionally engages in reflecting on the contemplations that lead to (eb) the realization of identity with Shiva. Note here that when shiva himself contemplates on his own identity the universe emanates. It is something like reflecting on zero which we write with much flourish and preen with every equation and all the transactions in the general ledger including qualitative which has been quantified. Utpaladeva (U Theories) and Abhinavagupta (A Theories) ambitiously conceive the Pratyabhijnā system as (=) both a philosophical apologetics (which follows Sanskritic standards of scholastic argument) and an internalized form of tantric ritual that leads students directly to identification with Shiva **(Pure consciousness)**. They explain the basic means by which the system conveys Shiva-identity according to the same basic ritual pattern described above, as (=) shaktyāvishkarana, "the revealing of Shakti." We must understand here that when Shiva, the pure consciousness reflects on his own self (zero) then Shakti is manifested, namely the phenomenal universe is manifested in all its dualities and conflicts. The Pratyabhijnā philosophers, however, also frame Shakti as (=) the reason of a publicly assessable inference, or "inference for the sake of others" (parārthānumāna). According to the scholastic logic, the reason identifies (e&eb) a quality in the inferential subject "I" known to be invariably concomitant (e&eb) with the predicate, "Shiva." Thus I am Shiva because (e) I have his quality, that is, Shakti, the capacity of emanating and controlling the universe. This is most important and the is interpreted as Satchidananda (mine I = cosmic consciousness). Terrestial Brahman Anti Brahman or celestial Brahman-AntiBrahman has the quality but they are marginalized by Para Brahman. That is why you see destruction of powerful empires, Tsunamis, earthquakes, and other natural disasters like supernovae, gamma rays etc, In order to address debates on epistemology that were then current, Utpaladeva and Abhinavagupta further explain the mythic and ritual pattern of Shiva and Shakti in terms of recognition. The specific problem the writers address had been formulated by the Buddhist logic school of Dignāga and Dharmakīrti, which flourished in medieval Kashmir. Contemporary interpreters have characterized the philosophy of Buddhist logic as (=) a species of phenomenalism akin to that of David Hume (D Theories). According to this school, the foundation of knowledge is (=) a series of momentary and discrete perceptual data (svalakshana). There are no grounds in those data for the recognitions of any enduring entities through ostensible cognitions utilizing (e) linguistic or conceptual interpretation (savikalpaka jñāna). In debates over several centuries, the Buddhist (B-Theories) logicians had propounded arguments attacking (e) many concepts that seemed commonsensical and were religiously significant to the various orthodox Hindu philosophical schools—such as ideas of external objects, ordinary and ritual action, an enduring Self, God, and revelation. The

Pratyabhijnā philosophers' response to the problematic posed by Buddhist logic revolutionized earlier approaches of the Nyaya philosophers, the Shaiva Siddhāntin Sadyojyoti and even Utpaladeva's teacher Somānanda, and may be characterized (eb) as a form of transcendental argumentation. Utpaladeva and Abhinavagupta **(U-A Theories) interpret (eb) their central myth of Shiva's emanation and control of the universe through (e&eb) Shakti as (=) itself an act of self-recognition (ahampratyavamarsha, Pratyabhijnā).** Furthermore, abjuring Somānanda's agonistic stance towards Bhartrihari (B Theories) they also equate Shiva's self-recognition (Shakti) with (=) the principle of Supreme Speech (parāvāk), which they derive from the Grammarian. This I **interpret as ultimate expression "Om".** They thereby appropriate the Grammarian's explanation of creation as (=) linguistic in nature. Thus the Kashmiri Shaiva philosophers ascribe (e) to Speech a primordial status, denied by the Buddhist logicians. As ritual recapitulates myth, the Pratyabhijnā system endeavors to lead (eb) the student to participate in the recognition "I am Shiva," by demonstrating that all experiences and contents of experience are (=) expressions of the recognition that "I am Shiva." I interpret here again this means that I= cosmic consciousness the cosmic general ledger and by participation I am contributing and infact everyone even a insentient stone is contributing to "Grand design". The paradox of the Pratyabhijnā formulation of the inference for the sake of others is that the self-recognition "I am Shiva," as an interpretation of Shakti, becomes (=) in effect both the conclusion and the reason. There is some confusion here. I clarify that Shiva is pure consciousness from which impure consciousness Shakti emanates in division of phenomenal universe. Like there are transactions in zero, Shakti exists inside shiva. This is not the reason, but the very conservation law like thermodynamical or assets =- liabilities. However we continue with the assumed conclusion and reason theory further . This circularity of conclusion and reason is a consequence (e) of the Kashmiri Shaiva monism. From the intratraditional perspective, there is no fact that can be adduced in support of (eb) another separate fact, as everything is (=) always the same in essential nature. From the intertraditional perspective of philosophical debate, however, the circularity is (=) not necessarily destructive. The Shaiva technical studies of various topics of epistemology and ontology in effect provide (eb) further ostensible justification for this apparent circularity. Utpaladeva's (U Theories) and Abhinavagupta's (A Theories) epistemology may best be illustrated by (e) its approach to perceptual cognition.The Pratyabhijnā arguments on this subject may be divided into those centered around two sets of terms: prakashan; and vimarsha and cognates such as pratyavamarsha and parāmarsha. **Prakāsha is the "bare subjective awareness" that validates (eb) each cognition, so that one knows (e) that one knows. The thrust of the arguments about Prakāsha is analogous to (=) George Berkeley's thesis (G Theories) of idealism that esse est percipio. The Shaivas contend that, as no object is known without (e) validating awareness, this awareness actually constitutes (e) all objects. There is no ground even for (e) a "representationalist" inference of objects external to awareness that causes (eb) its diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here**

.Furthermore; the Kashmiri Shaivas argue that there cannot be another subject outside (e) of one's own awareness. They conclude, however, not with solipsism as usually understood in the West, but a conception of a universal awareness. All sentient and insentient beings are (=) essentially one awareness. **Vimarsha a**nd its cognates have (e) the significance of apprehension or judgment with a recognitive structure, and may be glossed as (=) "recognitive apprehension."The recognitive is (=) the act of recognizing or an awareness that something perceived (e) has been perceived before. Utpaladeva's (U Theories) and Abhinavagupta's (A Theories) arguments centering on these terms develop earlier considerations of Bhartrihari (B Theories) on the linguistic nature of experience. Utpaladeva (U Theories) and Abhinavagupta (A Theories) refute (e) the Buddhist contention that recognition is (=) a contingent reaction to direct experience by (e) claiming that it is (=) integral or transcendental to all experience. Some of the considerations they adduce to support these claims are the following: that children must build upon a subtle, innate form of (e) linguistic apprehension in their learning of conventional language. There must be a recognitive ordering of (e) our most basic experiences of situations and movements in order to account for (e) our ability to perform (eb) rapid behaviors. Some form of subtle application of language in all experiences is (=) necessary in order to account for our ability to remember them. The two phases of argument operate together. The idealistic Prakāsha arguments make the recognition shown by (e) the vimarsha arguments to be (=) integral to all epistemic processes, constitutive (e) of them and their objects. Moreover, on the radical logic of the Kashmiri Shaiva idealism, the recognition generating all things belongs (e) to one subject. It must therefore be (=) his self-recognition. As it is through the monistic subject's self-recognition that all phenomena are (=) created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation through (e) Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.** The student, by coming to see this self-recognition as (=) the inner reality of all that is experienced, is led (eb) to full participation in it. Also noteworthy is the Kashmiri Shaiva theory of what may be called "semantic exclusion" (apoha). This concept had originally been formulated by the Buddhist logicians to explain (eb) a nonepistemic "coordination" (sārūpya) **between languages and (e&eb) momentary perceptual data** as the basis (e) for successful reference in communication and behaviors. According to the Buddhists, words have (e) no isomorphism with (e&eb) the sense data, but only exclude (e) other words that would not (e) lead to successful behavior. The only reference of the word "cow" to a perceived particular is (=) that it excludes non-cows, for example, a horse, a car, and so on. The Buddhist theory has an interesting point of agreement with (=) contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is not formulated like the latter (that is, on the basis of considerations about the systematicity of entire languages).Utpaladeva and Abhinavagupta argue that exclusion itself depends upon (=) a comparative synthesis, or recognition, of what does and does not fit within particular categories. We recognize that the cow is (=) not a non-cow such as a horse. The Pratyabhijnā theorists thus in effect explain difference itself as (=) a kind of similarity. Difference is identified (e&eb) in

various circumstances like other forms of similarity. According to the Shaivas such difference-identification is (=) one of the principal expressions of Shiva's emanating self-recognition. **The Pratyabhijnā Ontology: The Syntax of Empowered Identity** Just as Utpaladeva and Abhinavagupta appropriate Bhartrihari in equating self-recognition with (=) Supreme Speech and thereby interpreting recognitive apprehension as (=) linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā). The Grammarian view itself originated in Brahmanic interpretations of the Veda as (=) expressing injunctions for sacrifice. The Kashmiri Shaivas further agree (=) with much of **Vedic exegetics in conceiving being as both narrative and recapitulatory ritual action.** Following the account above, it is **Shiva's mythic action through Shakti as self-recognition that constitutes (eb) all experience and objects of experience, and that is reenacted by philosophical discourse**. The Pratyabhijnā thinkers propound their philosophy of Shiva's action to explain (eb) a wide range of topics of ontology. One of their concerns is to describe how Shiva's action generates (eb) a multiplicity of relationships (sambandha) or universals (sāmānya) as (=) the referents of discrete instances of recognitive apprehension. With this theory they attempt to subvert (e) the Buddhist logicians' contention that evanescent particulars are (=) ontologically fundamental. For the Shaivas, **categories are (=) primitive, and particulars are formed out (e) of syntheses of those categories. This is what is being attempted to The Theory of everything by classification and equational dovetail.** Most illustrative of the Pratyabhijnā thinkers' "mythico-ritual approach" to ontology is their use (e) of theories of Sanskrit syntax to explain Shiva's action. Again reflecting the Vedic roots of South Asian philosophies, many schools of Hinduism and Buddhism—even those which do not (e) view all existence as action—frequently advert to considerations of action syntax in treating (e&eb) ontological or metaphysical topics. The relevant considerations pertain to how verbs articulating action relate to (e&eb) declined nouns indicating the concomitants of action (kārakas)—in English, roughly, the agent, object, instrument, purpose, source and location. Now, most Sanskritic philosophies, Hindu as well as Buddhist, have tended to delimit (e) the syntactic role of the agent **(kartri kāraka)—**to different degrees, but sometimes quite strongly. The explicit and implicit reasons for this tendency are complex. At one level it evidently reflects the orthodox Brahmanic norms that subordinate the individual's agency to the order of objective ritual behavior—pertaining to sacrifice, caste, life cycle, and so on. It also seems more broadly to reflect both Hindu and Buddhist concepts of the agent's bondage to (e) the process of action and result (karma) extending across rebirths (see Gerow 1982). The mainstream Buddhist philosophies completely deny (e) the existence of a self in the dependent origination **(pratityasamutpada**) of karma. Developing suggestions of Somānanda, the Pratyabhijnā philosophers expound a distinctive theory of agency to **rationalize their tantric mythic and ritual drama of omnipotence**. In their theory they take up several earlier understandings of the positive albeit delimited role of the agent and radicalize them. According to the Kashmiri Shaivas, all causal processes and other relationships constituting (eb) the universe are synthesized and impelled by (e&eb) the mythic agency of Shiva in his act of self-recognition.

Shiva's agency encompasses (e&eb) the actions of sentient beings as well as the motions and transformations of insentient beings. See earlier models of projection, axes, dimensions, rotations, foldings, concurrences, shocks, encounters, etc. The Kashmiri Shaivas ultimately reduce (e) the entire action of existence to agency. As Abhinavagupta (A Theories) explains, **"Being is (=) the agency of the act of becoming, which is, the freedom characteristic of (e) an agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)."** Again, this theory of omnificent syntactic agency is ritually axiomatic as well as mythical. Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy, in a manner homologous to (=) the epistemology of recognition, as leading to (eb) salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design". Abhinavagupta (A Theories) likewise, in his explanation of the preliminary ceremonies of the tantric ritual, identifies (e&eb) various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases**. He explains that the aspirant's goal in the ritual action is (=) identification with Shiva as agent of all the cases. **Tantra and Kashmiri Shaivism (Wikipedia) for models see sections from (1) to (12)**

Spacelessness is not lack ofspace but cancellation of all positive and negative space and so with time

Only present is positive; past is negative and future is imaginary: a t (1) –b t (2) + i (ct (3)

The spirit soul bewildered by the influence of false ego thinks himself as the doer of activities that are actually carried out by rajas, tamas and Sattva, which are my manifestation in the material world (Geeta: Karma yoga: Text 27, Page 193, Prabhupada: Geeta as it is)

***@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@
@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@@***

Acknowledgements: I have made concerted efforts, sustained and protracted endeavours for the inclusion of every one and every source either in references or cross references. In the eventuality of any act of omission and commission, I make a sincere entreat, earnest beseech and fervent appeal to kindly pardon me. By nature, the work called for browsing and collating and coordinating thousands of various documents. Please excuse me in case of any prevarication from rules, or miss. Classification of the protagonist or antagonist laws is based on the characterstics and pen chance, predilection, proclivity and propensities of the systems under investigation. We acknowledge in unmistakable and unambiguous terms the help of Stanford encyclopedia, Kants writings, and Deleuze's Logic Of

Sense, Penrose and Hawking's nature of space and time, Penrose's Emperor's New Mind, Shadows of mind, Ken Wilber's spectrum of consciousness etc., great men seem to be endless….google search and Wikipedia. Concerted and orchestrated efforts are made to put on recordial evidence the names of all people either under references list or mostly by cross references so that no one is missed. If there be any act of omission or commission, it is my sincere entreat, earnest beseech, fervent appeal to kindly pardon me and the error is absolutely inadvertent and in deliberate. Let not any sensibilities, susceptibilities, and sentimentalities be hurt. I want to put on record with humble gratefulness the help by American Physical society, nature and other Noetic institutes of US who sent lot of alerts for my reference which was very valuable, and could not have been found by me despite assiduous and fervent search. Most important type is that concerning the stability of solutions near to a point of equilibrium. This may be discussed by the theory of Lyapunov. In simple terms, if all solutions of the dynamical system that start out near an equilibrium point x_e stay near x_e forever, then x_e is Lyapunov stable. More strongly, if x_e is Lyapunov stable and all solutions that start out near x_e converge to x_e, then x_e is asymptotically stable. The notion of exponential stability guarantees a minimal rate of decay, i.e., an estimate of how quickly the solutions converge. The idea of Lyapunov stability can be extended to infinite-dimensional manifolds, where it is known as structural stability, which concerns the behavior of different but "nearby" solutions to differential equations. Input-to-state stability (ISS) applies Lyapunov notions to systems with inputs. von Neumann stability is necessary and sufficient for stability in the sense of Lax–Richtmyer (as used in the Lax equivalence theorem): The PDE and the finite difference scheme models are linear; the PDE is constant-coefficient with periodic boundary conditions and have only two independent variables; and the scheme uses no more than two time levels (See Wikipedia) Von Neumann stability is necessary in a much wider variety of cases. It is often used in place of a more detailed stability analysis to provide a good guess at the restrictions (if any) on the step sizes used in the scheme because of its relative simplicity. Albeit forwarded in nine module systematizations, the entire gamut is to be seen in a single shot, and the presentation of nine schedule twenty seven storey models is to circumvent typing of hundreds of superscripts and subscripts. In fact the statement is made inclusive of all previous models, and the variables are definitely different for each schedule, which again is reinstated due to typing of corresponding variables millions of systems, a fastidious and fussy work again. I beg pardon for any inconvenience caused to the readers due to such utilization of convention. I put on **recordial** evidence and acknowledge my heartfelt thanks for the contribution of dear **Professor Gnanendra Prabhu** for sharing his elephantine, phenomenal, monumental and versatile knowledge about, differential topology, functional analysis, complex analysis and philosophy**.** I am grateful to Professor Chadralekha MD PhD. Tagore medical College, Chennai for deliberations and discussions on Medicine. To discussions on Physics topics credit goes to Dr. A.S. Krishna Prasad, Former Director DRDO Chapter, and Bangalore Chapter. Prof. Sunita MSc. PhD., of MS Ramaiah University helped me with valuable suggestions on Aerodynamics and propellant chemistry. Sir KVB Pantulu, former Chairman of NALCO, ESSAR Steels helped in formatting process and project management advices.

Note: Here we talk of the characterstics of systems which satisfy the condition of cosmological constant. or any other for that matter or variable or axiomatic predication. . There are lots of zeroes corresponding and concomitant to the second law of black holes. Infact as many as that of extant and existential blackholes exist. At the outset, it is to be stated that there are hadrons in every system. Supersymmetry between forces and matter, with both open and closed strings; no tachyon; group symmetry is SO (32) and its axiomatic predications, predicational anteriorities, character constitution shall be extant and existential in very many systems, and the characterstics are taken in to consideration in the classification scheme. Many systems have such fundamental instabilities like that of quantum gravity and characterstics of those systems form the citadel and fulcrum, bulwark and manor, mainstay and reinforcement, alcazar and chateau theory has a fundamental instability on which the classification schémas are valid. There are lots of systems which follow the axioms of string theory It is the characterstics of this system which are taken in to consideration in the classification scheme. There are various systems that have the same bastion, support system, stylobate and sentinel as that of the Deleuzean terms and predications and phenomenological methodologies systemized. Each and every system has electrons, neutrons and protons and for that matter quarks. There shall be strong nuclear force and weak nuclear force. There are many systems that satisfy the criterion

specified by the equation, principle or statement in question. Characterstics of the investigating systems form the bastion for the classification scheme and doxa thereof. Systemic differentiation is conducted. Despite gravity being constant, there exists gravity between two objects, and this could be taken as a system. Depending upon some parametric representationalities, functionalities, advantageousness, appropriateness, benefit, facilitation, fittingness, helpfulness, instrumentality, merit, practicality, serviceability, suitability, and usefulness, utility, these systems could be classified in to various categories. In respect of an equation, there shall be many systems that satisfy the given equation. Equations themselves could be by the utilization of the model solved term by term as has been exemplified and illustrated many time in the previous papers. There is lot of systems that could be brought in to the orbit of and gamut of the theory in question which the investigatable systems satisfy the axiomatic predications and postulation alcovishness of the systems in question. Towards the end of classificational consummation, consolidation, corporation and concatenation we take the characterstics of the systems, the predicational interiorities, ontological consonance and primordial exactitude, accolytish representations, functional topology, apocryphal aneurism and atrophied asseveration, event at contracted points, and other parameters as the bastion and stylobate of the stratification purposes. Such totalistic entities would have easy paradigm of relational content, differentiated system of expressly oriented actions with primary focus and locus of homologues receptiveness and differentially instrumental activity, variable universalism and particularism, imperative compatibilities and structural variabilities, interactional dynamical orientation, institutionalization and internalisation of pattern variables common attitudinal orientation of constituionalisation of internalized dispositions, and a qualitative gradient of structural differentiation and ascribed particularistic solidarity abstraction or interactional dynamics, internal differentiation, structural morphology, formal characterization, concept formulation, phenomenological methodologies, constituent structure, transformational minimal condition, paradigmatic feasibilities, programmatic plausibilities, comparative variability, normative aspect of expectational prediction, projection and prognostication as consideration of the investigatory systems. Any scale can be used that is convenient to the classification scheme. It is important to note that the scheme of classification and the stratification doxa must not be adversely antagonistic, inexorably irreconcilable, antithetically antipodal, diametrically opposed, repugnantly retrogressive, inimically inverse, violatively unsimilar, diversely dissimilar, antipathetically anti thetical, conflictingly combative, obstructively pugnacious, inimically obstructive, repellently restrictive, disputatiously gainsaying or conformingly pugnacious to the axiomatic predications and postulation alcovishness of the theory in question or the equation representative or constitutive thereof. Sole intention, main objective and primary aim is twofold. One is towards the end of circumvention of the extra equation and the concomitant and corresponding variable therein. Second is avoidance of clustered congest, swarmed huddle, mustered pack, sardine squash, and swamp throng in the scheme of classification. Only thing that is sought out is the consonance in the entire diaspora and body fabric of the systems under study. This statement is true and holds unmistakably true for all the papers and I sincerely entreat readers to remember the statement and read the paper against this background. When we write A+B we mean by that B is being added to A or vice versa. It is like adding milk to water and water to milk. There may or may not be a time gap. As said earlier there may be many systems that satisfy the conditions of the equations and those systems that are investigatory or investigatable are taken in to consideration based on their characterstics in the classification doxa. When there are more than two entities, we can take logarithm and find the value of that factor to be taken with anti log to obtain prediction and projected values of the model. In case of A-B, we are removing B from A and that means B is eating up A. These factors are taken in to consideration in the application of the model to equations. Log (ab) and Log (a+b) is well defined. Model stands out as universal testament and template for application to each sentence and equation what with the quantification process done and the correlations well defined. In the eventuality of non existence of any connection at some phase, the model would have the accentuation and attrition coefficients and detritions coefficients as zero rendering the equations of concatenation simpler. Projection formula which incorporates in its diaspora the initial values provide autheticative determination, unimpeachable validation, incontrovertible establishment , apodictic evidence, reliable roll out, unfailing cinquecento quattrocento trecento, incontrovertible indication of the final finale, notwithstanding appellation, appellative, brand, cognomen, compellation, designation, flag, handle, identification, label, moniker, nomen, slot, style, surname, tab, tag, term, title designation, appellation, appellative, class, classification, denomination, description, epithet, of the classification scepter, scenario, scimitar,

schottische. There is pure and impure consciousness in every one. Gratification producing and deprivation producing one's can be easily classified from individual general ledgers. Similar analogy holds for collective general ledger and cosmic general ledger or nature's general ledger. It is also to be noted that while dealing with equations towards the end of consummation of the measure, it is necessary that the two variables are to be classified in to three sections and each one would have the adventitious and decidedly stated relationship whereby the fundamental equations are drawn up and the analysis made. All the parametric representationalities, conditionalities, orientationalities remain unequivocal as stated in the variables stated in to consideration section and are different from module to module. In essence the paper is to be read as holistic one with the sole intention, primary objective and raison d' etre to build a TOE. Towards the end of circumvention of typing hundreds of superscripts and subscripts which would be a sardine squash and the concomitant operational difficulties, model is presented in piece meal of nine modules. Logarithms are to be taken in respect of those which incorporate more than one variable in bra-ket. Values of log (ab) and log (a+b) are readily available. Anti logarithm shall be taken at the end to predict, project, prognosticate the value of the variable. This is true for tensors, vectors and other variables too. Affirmational assertion, and explanation justification, statement of vindication, annotational commentary, and explicational glossary for each and every system changes and physical interpretation of results is one thing that is to be with earnest endeavour and feverish and febrile expediency. Editions never ever mean the same and identicality of parameters and this has been explicitly and unmistakably stated in the model apriori itself. Akin and analogous, cognate and concurrent, correspondent and congruent, comparable and complementary, synonymous and duple, tantamount and agnate, commensurate and correlative representation is only to highlight the importance and subterfuge the replication of work. It is my fervent solicitation to kindly bear with me for any lapses, notwithstanding the orchestrated efforts for a paper without any minor errors. Postulation predication, conclusive presumption, differential presuppositions, underscored decidedly axiomatic statement of the statement, equation form the bye word or the watch word in the aggrandizement-amplification, caricature and crock, understatement-unembellishment, elocution-emphasis, enunciation-inflection, announcement-argument, articulation, assertion, asseveration, choice of words, commentary and communication, declaration and definition, delivery-diction, elucidation-emphasis, enunciation-execution, explanation, exposition-formulation, idiom, interpretation and intonation tone and tenor of stratification. Now, how do find the reaction of systems to these singularities. You do the same thing a boss does for you. "Problematize" the events and see how you behave. I will resort to "pressure tactics". "intimidation of deriding report", or "cut in the increment" to make you undergo trials, travails and tribulations. I am happy to see if you improve your work; but may or may not be sad if you succumb to it and hang yourself! We do the same thing with systems. systems show conducive response, felicitous reciprocation or behave erratically with inner roil, eponymous radicality without and with blitzy conviction say like a solipsist nature of bellicose and blustering particles, or for that matter coruscation, trepidiational motion in fluid flows, or seemingly perfidious incendiaries in gormandizing fellow elementary particles, abnormal ebullitions, surcharges calumniations and unwarranted(you think so but the system idoes not!) unrighteous fulminations. So the point that is made here is "like we problematize the "events" to understand the human behaviour we have to "problematize" the events of systems to understand their behaviour. This statement is made in connection to the fact that there shall be creation or destruction of particles or complete obliteration of the system (blackhole evaporation) or obfuscation of results. Some systems are like "inside traders" they will not put signature atoll! How do you find they did it! Anyway, there are possibilities of a CIA finding out as they recently did! So we can do the same thing witn systems to. This is accentuation, corroboration, fortification, .fomentatory note to explain the various coefficients we have used in the model as also the dissipations called door. In the bank example we have clarified that various systems are individually conservative, and their conservativeness extends holisticallytoo.that one law is universal does not mean there is complete adjudication of nonexistence of totality or global or holistic figure. Total always exists and "individual" systems always exists, if we donot bring Kant in to picture! For the time being let us not! Equations would become more ensorcelled and frenzied..... philosophy merges with ontology; ontology merges with univocity of being; analogy has always a theological vision; not a philosophical vision; one becomes adapted to the forms of god; self and world; the univocity of being does not mean that there is one and the same being; on the contrary, beings are multiple and different they are always produced by disjunctive synthesis; and they themselves

are disintegrated and disjoint and divergent; membra disjuncta.the univocity of being signifies that that being is a voice that is said and it is said in one and the same "consciousness". Everything about which consciousness is spoken about being is the same for everything for which it is said like gravity; it occurs therefore as an unique event for everything; for everything for which it happens; eventum tantum; it is the ultimate form for all of the forms; and all these forms are disjointed; it brings about resonance and ramification of its disjunction; the univocity of being merges with the positive use of the disjunctive synthesis, and this is the highest affirmation of its univocity like gravity; it is the eternal resurrection or a return itself, the affirmation of all chance in a single moment, the unique cast for all throws; a simple rejoinder for Einstein's god does not play dice; one being, one consciousness, for all forms and all times. a single instance for all that exists, a single phantom for all the living; a single voice for very hum of voices; or a single silence for all the silences; a single vacuum for all the vacuums; consciousness should not be said without occurring; if consciousness is one unique event in which all the events communicate with each other; univocity refers both to what occurs to what it is said. This is attributable to all states of bodies and states of affairs and the expressible of every proposition. So univocity of consciousness means the identity of the noematic attribute and that which is expressed linguistically and sense fully. Univocity means that it does not allow consciousness to be subsisting in a quasi state and but expresses in all pervading reality; CONSCIOUSNESS AND ITS UNIQUENESS June 12, 2012 at 8:40pm (See Deleuze Logic of sense, Wikipedia and Stanford encyclopedia for more details) Atrocious contrivance and device, gratificational primogeniture, calamitous dodge and expedient, depraved, destructive, disastrous, execrable gambit and gimmick, great solace and succor, iniquitous, injurious, loathsome, low, maleficent, malevolent machination and maneuver, promethaleon of candor, frankness, honesty, honor, ingenuousness, innocence, openness, reality, sincerity, truthfulness, spiteful, stinking, ugly, unpleasant, unpropitious play and ploy, progenitor of jurisprudence and circumspection, racket and ruse, savvy, scam, stratagem, subterfuge, tactic and wile, forthrightness, honesty, truthfulness embodied and personified is the structural predisposition and dispensation of the complex. Nobody need have to take the Brahman-Anti Brahman agency to be a recalcitrant, repugnant, and refractory proposition. God gives and also takes away. That is the point made. And of course lesser gods, and God doth follow them and they follow God. That is why they are lesser gods. And wield the power on lesser mortals like us. When we mean gratification and deprivation we mean celestial Brahman Anti Brahman and Terrestial Brahman and Anti Brahman agencies producing such happiness or sadness by any means such as (blending- amalgamation, adulterant-adulteration, amalgam-amalgamation, blend-combination, composite- compound, debasement-denaturant, fusion-hybrid, intermixture-reduction, amalgam-mixture, composite- compound, fusion-mishmash, aggregate- alloy, amalgamation-blend, commixture-composite, composition-compost, conglomerate-fusion, goulash-medley, mishmash-stew, blend-coalition, commixture-compound, federation-heating, integration-junction, melting-merger, synthesis-unification, assortment- combination, adulteration-alloy, assimilation-association, batter-blend, brewed combine, composite compound, concoction confection, conglomeration cross of money, orientation(dis), penury creation, murder, mayhem plunder, pillage, apocalypse, Armageddon in various forms and permutation and combination of Manichaean artifice, agathokakological malevolence, agley chicane fourberie, fraud, furtiveness, gambit, awry and bad ,flagitious mechanizations, cacodemonic gambit, hanky-panky, intrigue, machination, deprecatory and diabolic underhandedness and wiles, energumenical skullduggery, goetic chicanery, lenocinant stratagem, malefic and maleficent strategy, malominous sell, sellout, sham subterfuge, peccable, humbug, imposture, misrepresentation peccant fraud, fraudulence, hoax, qued , scelerate pretense, ruse, sell, sellout, sham stratagems' of maladroit manouevures , sinistral ,tortious and venal sophistry, contingency enterprise, endangerment emprise, jeopardizing adventure, jejune jeremiad, avoidable inertia, latent passiveness: with each action accounted and showing balance in the nature's general ledger as also actionlessness:e). All these are recorded in Nature's general ledger. Neuron DNA encapsulates all the actions passions, actions, interactions, transactions or the lack of it at various levels of individual, collective and cosmic levels. In celestial Brahman Anti Brahman ledger there is cessation of dualities. One more point to be made is if 5=3+2, then we can say 5 gormandizes 3 or 5 gobbles up 2. Quintessentially such a statement arises out of the difference of LHS and RHS term wise. It must be remembered despite all theoretical abstractions and generalizations be it proton mass or Higgs Boson, or topology of a space, it is notwithstanding the fundamentality of nature is just a number. Just everything in Physics and mathematics is just a number. Let us

assume I buy rice in Bangalore (District) worth 5k.g. I can always divide 5k.g.s of rice in categories of Bangalore (South), Bangalore (west), Bangalore (East) and Bangalore (South). On similar basis, there are billions of neutrons, protons and concomitant interactions which would satisfy the given numerical value. Example of gravity above makes things clear. Everything atleast is a function of time and someplace or some other coordinates, subordinates or superordinates. Mass of proton is 2.47 does not mean total mass of all protons is 2.47. It is this argument that is used in the classification scheme. By the very nature, self the unmonitoring but the witness consciousness agency identifies itself with something, be it a profession, an actor on the screen, when it does not have a identity. Such examples are legion, a Charlie Chaplin in "The Kid" or Raj kapoor in "Jagte Raho". These are people who just wanted to live and living without being put under surveillance every moment is the greatest part of their life. This is "Absolute Subjectivity" of Shiva. Such a state is always craving for identification for in pure consciousness all identities are lost. It is this "identitylessness" or having very minimal identity is what pervades most in India and this is due to expansion of individual consciousness. When that happens, be it education, or wealth, the "identity' becomes so well entrenched, they become the exact opposite of "pure consciousness"," impure consciousness" (Shakti). This is a state of "Absolute or relativistic objectivity". There was this case of a lady who went to such an extent of planning, she was sure that whatever she does must work out. When it did not, she lapsed in to "identity crisis". Such thing as this might happen in well developed countries where the competition is extremal and to be achieved at any cost. Realisation that one is not the doer but the witness consciousness is just like watching a movie in which one has acted. It is in this sense "advaita", "non duality' seems to be of cardinal and preeminent importance for the analytical treat ment of "identity' and "identity lessness", both happen in space time. "Multiverses" in this sense is also like a different film of an actor released at different theaters and it is in this sense we state that that every object in space time is a multiverse. You must have totally immersed in doing a "string theory" problem or watching a "film", oblivious of surroundings. Essentially creation (space time) also is what is happening in mind and honed in to you billion times. Identification of anything with a designation or signification is one that leads to association within spacetime, which would only have pathological consequences and detrimental ramification as stated above or positive implications of "happiness" because of the identity itself. Search or 'show" has to go on! It gives a sense of consummation. Like the discussion of psephologists now doing about electoral results in India!

@@
@@
@@

Introduction:

In excited reveries: 27th February 4.15 Am early morning

It is like Gate way of India!

Black hole with unknown internal differentiation, comparative variability and structural morphology!

What?

Men may come and men may go but it stays on forever!

I am doll a doll

Marionette cup cake of all

I am gimcrack moppet assigned by God,

I am fantoccini, a Don's moll!

I cannot see what flowers are at my feet,

Nor what soft incense hangs upon the boughs,

But, in embalmed darkness, guess each sweet

Wherewith the seasonable month endows

Young Tam came up an' eyed me quick

With reddened cheek --

Braw Tam was daffed like a chick --

He could na speak --

The grass, the thicket, and the fruit-tree wild;

White hawthorn, and the pastoral eglantine;

Fast fading violets cover'd up in leaves;

And mid-May's eldest child,

The coming musk-rose, full of dewy wine,

The murmurous haunt of flies on summer eves

(Keats: ode to Nightingale, A Galloway song: I have mixed up!)

If only we accept that the quantum state is no more than a representation of the information we have, then the spontaneous change of the state upon observation, the so-called collapse or reduction of the wave packet, is just a very natural consequence of the fact that, upon observation, our information changes and therefore we have to change our representation of the information, that is, the quantum state. (Bell's inequalities1999, p. S291)

Pratyabhijñā Vimarśinī!

Quantum mechanics predicts non-locality unless position and momentum were (=) simultaneous "real" properties of a particle!

Nature also is!

No deterministic Local Realistic Theory is consistent with the Quantum Mechanical predictions of the entangled state!

Hamlet!

Quantum mechanics is non-local!

Strange bedfellows!

Two wrongs don't make a right."!

"The squeaky wheel gets the grease."!

Somebody said “robber” and he touches his shoulders!

"Keep your friends close and your enemies closer."!

"A watched pot never boils."!

Live coward and a dead hero!

With many a curve my banks I fret,

By many a field and fallow,

And many a fairy foreland set,

With willow- weed and mallow

I chatter, chatter as I flow,

To join the brimming river,

For men may come and men may go,

But I go on forever (Wikipedia)

What the hell are you talking about?

. I wind about, and in and out

With here a blossom sailing,

And here and there, a busty trout,

And here and there a grayling!

In what context…. ?

And here and there a foamy flake,

Upon me, as I travel,

With many silvery water break,

Above the golden gravel! (Alfred Tennyson Wikipedia)

Impatient imprecations, Blunt- brusque mutterings, Crude-crusty snarls, Curt- direct groans, Discourteous- gruff sputter, impetuous- impolite croak, matter-of-fact- rough grouch, short- snappy grunt, snippy- uncivil rumble , ungracious rumble calm, deliberate –expansive grouse , Gradual-unkind mutter, Abrupt- bluff babble, Candid-crusty blubber, Rude-short gibberish, Trenchant- blunt patter, Brusque- Compendious jabber, Pithy- Sharp burble, Small- Succinct cackle, Gruff- Hasty tattle, Impolite-sharp patter, Tart-Terse burble, Short- Snappy- Snippy grumble! What is that solecism of Richard III!

Ugh!

Stop it!

I was talking about self!

Le coeur a ses raisons, que la raison ne connaît point. [The heart has its reasons which reason is not acquainted with.] (Kierkegaard)

[Man is but a reed, the frailest thing in nature; but he is a thinking reed. The entire universe need not arm itself to crush him. A mist or drop of water is sufficient to kill him. But should the universe crush him, man would still be nobler than what kills him, because he knows that he dies; but the universe knows nothing of the advantage it has over him.] (Kierkegaard)

EXISTENCE precedes (e) ESSENCE. Forms do not (e) determine existence to be what it is. Existence fortuitously becomes (=) and is whatever it becomes and is, and that existence then makes up its "essence". 2) An individual has (e) no essential nature, no self-identity other than that involved (e&eb) in the act of choosing. 3) Truth (=) is subjectivity. 4) Abstractions can never grasp (e) nor communicate the reality of individual existence. 5) Philosophy must concern itself with (e&eb) the human predicament and inner states such as alienation, anxiety, inauthenticity, dread, sense of nothingness, anticipation of death. 6) The universe has (e) no rational direction or scheme. It is (=) meaningless and absurd. 7) The universe does not provide (eb) moral rules. Moral principles are constructed by (e) humans in the context of being responsible for (e) their actions and for the actions of others. 8) Individual actions are (=) unpredictable. 9) Individuals have (e) complete freedom of the will. 10) Individuals cannot (e) help but make choices. 11) An individual can become (=) completely other than what he is (Dictionary of Philosophy, p. 88). **A Primer on Kierkegaardian Motifs Anthony storm Models are given in the next paper attributable and ascribable to spatial constraints**

You said it is a record keeper and a non monitoring agency a Neuron DNA!

I was wondering whether there is a difference between the anagrammatic mind acting on information and a is dynamic self possible!

Dynamic self and pulsating and throbbing self is possible evolutionarily!

What do you mean?

Individual consciousnesses (individual general ledgers) when expanded lead to cosmic consciousness the cosmic general ledger!

There is a view of life which conceives that where the crowd is, (eb) there is also the truth, and that in truth itself there is (=) need of having the crowd on its side. There is another view of life which conceives that wherever there is a crowd (eb) there is untruth, so that (to consider for a moment the extreme case), even if every individual, each for himself in private, were to be in possession of the truth, (eb) yet in case they were all to get together in a crowd—a crowd to which any decisive significance is attributed, a voting, noisy, audible crowd—(eb) untruth would at once be in evidence (The Single Individual, p. 110). (ibid) **Models are given in the next paper attributable and ascribable to spatial constraints**

OK! That is when the big crunch comes!

Why?

You have become ParaBrahaman!

Not Brahman Anti Brahman?

No! They are lesser Gods!

Is mind a curse?

In order to clarify the divergence of objective and subjective reflection, I shall now describe subjective reflection in (eb) its search back and inward into inwardness. At its highest, inwardness in an existing subject is (=) passion; truth as a paradox corresponds (e&eb) to passion, and that truth becomes (=) a paradox is grounded precisely in its relation (e&eb) to an existing subject. In this way the one corresponds to the other. In forgetting that one is an existing subject, one loses passion... [And] the knowing subject shifts from being human to (eb) being a fantastical something.... When the question about truth is asked objectively (eb) truth is reflected upon objectively as (=) an object to which the knower relates (e&eb) himself. What is reflected upon is not the relation but that what he relates himself to is the truth, the true. If only that to which he relates himself is the truth, the true, then the subject is in the truth. When the question about truth is asked subjectively, the individual's relation is reflected upon subjectively. If only the how of this relation is in truth, the individual is in truth, even if he in this way were to relate himself to untruth (**Concluding Unscientific Postscript, p. 198f.).(ibid) Models are given in the next paper attributable and ascribable to spatial constraints**

Why do they hide!

When sin is posited in the particular individual by the qualitative leap, the difference between good and evil is also posited. We have nowhere been guilty of the foolishness that holds that man must sin; on the contrary, we have always protested against all merely imaginatively constructed knowledge. We have said what we again repeat, that sin presupposes itself, just as freedom presupposes itself, and sin cannot be explained by anything antecedent to it, anymore than can freedom. To maintain that freedom begins as liberum arbitrium [free will]...that can choose good just as well as evil inevitably makes every explanation impossible. To speak of good and evil as the objects of freedom finitizes both freedom and the concepts of good and evil. Freedom is infinite and arises out of nothing. Therefore, to want to say that man sins by necessity makes the circle of the leap into a straight line (The Concept of Anxiety, p. 112).(ibid) **Models are given in the next paper attributable and ascribable to spatial constraints**

I told you about the “moon” problem, “cat” problem and “uncertainty” problems!

An individual in despair despairs over something. So it seems for a moment, but only for a moment; in the same moment the true despair or despair in its true form shows (eb) itself. In despairing over something, he really despaired over (eb) himself, and now he wants to get rid of (e) himself. For example, when the ambitious man whose slogan is "Either Caesar or nothing" does not get to be Caesar, he despairs over (eb) it. But this also means something else: precisely because he did not get to be Caesar, he now cannot (e) bear to be himself. Consequently he does not despair because (e) he did not get to be Caesar but despairs over himself because (e) he did not get to be Caesar.... Consequently, to despair over something is (=) still not despair proper.... To despair over oneself, in despair to will to be rid of (e) oneself—this is the formula for all despair (The Sickness unto Death, p. 19f.).(ibid). What Deleuze forgets here is that in wanting to be Ceasar, a truly ambitious man, would have inculcated all the properties and characterstics of ceaser and would have become the Ceasar for himself! Not to the world!

Denosjour pensioner with desolate desideratum, burlesque moccasin, arrondissiment arrogate, disputatious diabolic disport, rinderpest canine distemper, flibbertigibbet frivolous fiddle dude, temerarious tempestuous template, essetialistic mordant, schizostylis schlep, schizocarp schizoid, how many times have I told you to state the subject matter in the caption?

But you know!

Others’ do not!

World is suffering from “Moon” problem, “measurement” problem, and “indistinguishability”!

The subjective existing thinker is (=) aware of the dialectic of communication. Whereas objective thinking is (=) indifferent to the thinking subject and his existence, the subjective thinker as (=) existing is essentially interested in his own thinking, is existing in it. Therefore, his thinking has another kind of reflection, specifically, that of inwardness, of possession, whereby it belongs to the subject and to no one else.... 2. In his existence-relation to the truth, the existing subjective thinker is just as negative as positive, has just as much of the comic as he essentially has of pathos, and is continually in a process of becoming, that is, striving.... In the domain of thinking, the positive can be classed in the following categories: sensate certainty, historical knowledge, speculative result. But this positive is precisely the untrue. Sensate certainty is a delusion (see Greek skepticism...); historical knowledge is an illusion (since it is approximation-knowledge); and the speculative result is a phantom. That is, all of this positive fails to express the state of the knowing subject in existence.... 3.... Lessing has said that contingent historical truths can never become a demonstration of eternal truths of reason, also that the transition whereby one will build an eternal truth on historical reports is a leap.... 4. Lessing has said If God held all truth enclosed in his right hand, and in his left hand the one and only ever-striving drive for truth, even with the corollary of erring forever and ever, and if he were to say to me: Choose!—I would humbly fall down to him at his left hand and say: Father, give! Pure truth is indeed only for you alone! (Concluding Unscientific Postscript, p. 72f., 80f., 93, 106).(ibid)

So?

Others must know what you is your head!

I donot think there is much any way and I am scared others' might come to know!

Where was I?

I think Quantum coherence and coherent thinking!

Leibniz also follows Aristotle (cf. Metaphysics IV.3), in placing great emphasis on the Principle of Identity or the Principle of Contradiction (PC). PC states simply that "a proposition cannot (e) be true and false at the same time, and that therefore A is A and cannot be not A." (G VI 355/AG 321) According to Leibniz, the primary truths of his metaphysical system are identities, but, in a striking move, he combines PC with PIN and asserts in Primary Truths that "all remaining truths are reduced to primary truths with the help of definitions, that is, through the resolution of notions." (A VI iv 1644/AG 31) Furthermore, the combination of PC and (e&eb) PIN will mean that, since in any true proposition the predicate is contained (e) explicitly or implicitly within the subject, this is so for all affirmative truths, whether they be (=) universal or particular, necessary or contingent. Leibniz will use (e) this seemingly innocuous principle to draw profoundly strong metaphysical conclusions about the nature of substance and modality. (Stanford Encyclopedia)

What do you think?

OK1 let us assume a proposition is both true and false at the same time and see what happens to the equations

(Psi) = (psi) (true) e (true) in Dirac brackets + (psi) (false) e (false) in Dirac brackets!

Like Schrödinger's cat statement by Penrose?

Yes!

Now apply the model (see Section)

Empirical subjectivity is (=) constituted in the mind under the influence of (e&eb) the principles affecting it; the mind therefore does not (e) have the characteristics of a preexisting subject. (ES 29) These associations account not only for (e) experience in the basic sense, but up to the highest level of social and cultural life: this is the basis for

Hume's rejection (e) of a social contract model of society (such as Hobbes'), in favour of convention alone. Morals, feelings, bodily comportment, all of these elements of subjectivity are explained, not by (e) transcendental structures, such as Kant will propose, but by (e) the immanent activity of association. Once this habitual structure of the self is (=) in place, Deleuze suggests, the Humean concept of belief comes into play (eb), which is resolutely a central part of human nature. It describes (eb) the particularly human way of going beyond (e) the given. When we expect the sun to come up tomorrow, we do not do so because (e) we know that it will, but because of a belief based (e) on a habit. This in turn reverses (e&eb) the hierarchy of knowledge and belief, and results, for Deleuze, in a, "great conversion (e&eb) of theory to practice." (PI 36) Every act of belief is (=) a practical application of habit, without (e) any reference to an a priori ability to judge. Not only is (=) the human being thus habitual, on Deleuze's reading, but also creative (eb) even in the most mundane moments of life. Finally, Deleuze insists that one of Hume's greatest contributions to modern philosophy is (=) his insistence that all relations are (=) external to their terms: this is the essence of Hume's anti-transcendental stance. Human nature cannot (e) unite itself, there is no 'I' which stands before experience, but only moments of experience themselves, unattached and meaningless without (e) any necessary relation to each other. A **flash of red, a movement, a gust of wind, dashing of a woman against you, obscene gestures, spitting at you, talking extempore about you (my addition) and these elements must be externally related (e&eb) to each other to create (eb) the sensation of a tree in autumn. In the social world, this externality attests (eb) to the always-already interested nature of life: no relation is necessary, or governed by neutral laws, so every relation has (e) a localised and passional motive. The ways in which habits are formed attests to the desires at the heart of our social milieu. Subjectivity, as Deleuze describes it through his reading of Hume, is a practical, passional, empiricist concept, immediately located at the heart of the conventional, which is to say the social.**

Somehow the empiricism itself is unpalatable!

Prayer does not change God, it changes the one who prays!

Yeah!

Same with contemplation, introspection you mentioned!

It makes saints out of sinners, or saints out of sinners' society!

Whoever fights monsters should see to it that in the process he does not become a monster. And if you gaze long enough into an abyss, the abyss will gaze back into you. Nietzsche

That may not be true!

Dānañca, peyyavajjañca,

atthacariyā ca yā idha,

samānattatā ca dhammesu,

tattha tattha yathārahaṃ;

ete kho saṅgahā loke

rathassāṇīva yāyato.

Listen

Generosity, kind words,

doing a good turn for others,

and treating all people alike:

these bonds of sympathy are to the world

what the linchpin is to the chariot wheel. (Buddha)

May be the reference is in different context!

Both are talking about the crowds

Advaita is a singularity?

Singularity is aberration!

Advaita is non causal!

Dvaita talks of dependent reality!

Is not collective truth Nietzsche speaks of a dependent reality?

Yes!

Is dependent reality and illusion?

I donot know!

Think!

Science is a process, and spiritualism is involution!

Like jumping a job?

Joining a job you mean?

Yes!

Yes!

Experience?

Same as they ask for in recruitment adverts!

All are different?

Yes!

Common factor?

Zyphera!

Deductive reasoning is used to find the answers!

Each step is gormandized by the previous one?

Yes!

That is what I am doing!

Inductive method is Absolute subjectivity?

Looks so!

Pure consciousness?

Absolute subjectivity!

You wanted all the hype?

Yes!

Why are you angry then?

Helplessness to come up to expectations!

Consciousness waves?

Well! String Theory talks of information waves I think!

Vibration V (1) (e&eb) Vibration (2)…… and so on

On one hand you say escape from sound and sight , and on the other you talk of melodies consummating with each other!

Space time and Nirvikalpa samadhi?

Death like experience! Yes!

Neuron DNA of the universe?

Property of space and time!

But you classified it?

Yes! Like silent zones, Rajas dominated zones, violent zones and so on……..

You ean you can have an infinite column AL model for String Theory?

Yes!

Total Vibrations- Total frequencies-Total melodies…….!

I sometimes doubt Napier's rule itself!

Why so?

Mass=energy

OK!

Mass- energy is zero!

OK!

Then energy gobbles up mass!

Is it not a contradiction?

………………………………………..

……………………………..

Aw! I have to think over; reading, writing, speaking, and thinking at the same time is bit impossible!

@@
@@
@@

SECTION ONE

Shiva = Pure Consciousness (Here All Dualities Cease: See Earlier Models. Being= Becoming, Self=Nonself Order = Chaos Etc.,); Shakti =Impure Consciousness (Mother Divine: Responsible For The Phenomenal Manifestation)

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

While tantrism is a complex and controversial subject, one of its most definitive characteristics for contemporary classifications—if not its most definitive one—is the pursuit of power.

(1) Tantric traditions are (=) thus those that aim at increasing (eb) the power of the practitioner.
(2) The theological designation for the essence of such power is (=) Shakti (the female counterpart to the male divine principle, whose **essence is power: impure consciousness, one that is responsible for the manifestation of the phenomenal universe. We shall take this as string theory postulate with strings responsible for creation**).
(3) The manifestations of Shakti that the practitioner of tantra aspire after vary greatly, from relatively limited magical proficiencies (siddhis or vibhūtis), through royal power, to (eb) the deindividualized and liberated saint's omnipotence to the performance of God's cosmic acts.
(4) What is God or Para Brahman's act? What is Karma? Take my case for example. I was a professional banker who took VRS, an aspiring star, and an academician who wanted to just make it to the top irrespective of the fact whether I had the abilities and capacities. I got in to a Chakravyuha like Abhimanyu, and lead a life of trials and tribulations and travails for a decade. What is that I did to be punished for this or still being punished? I will assume that God (Parà Brahman will take in to account (e) what I had done in my earlier incarnations (I donot think I did anything like that in this incarnation) wherein I could have tormented or produced a sense of deprivation balance either equivalent to or less than what I am going through now. Or God must

have added interest and given me (eb) the punishment. I will assume both and give models. One with the deprivation balance equivalent to what I had done, and the other deprivation balance which includes the interest for the period between my act of commission and this incarnation. I will also take in to account the third possibility of Brahman-AntiBrahman who like spider weave the web, see the prey undergoing tormentations in the web and increase it just for "play"

(5) In his seminal essay, "Purity and Power among the Brahmans of Kashmir," the Oxford historian Alexis Sanderson elucidates that the tantric pursuit of such power transgresses (e) orthodox, mainstream Hindu norms that delimit human agency for the sake of symbolic and ritual purity (shuddhi) (Sanderson 1985).

(6) **Violating prescriptions regarding** caste, sexuality, diet and death, many of the tantric rites were originally performed in (eb) cremation grounds.

(7) Whereas in Shākta tantrism, Shakti as a Goddess is herself the ultimate deity, in monistic Kashmiri Shaivism she is incorporated into (eb) the metaphysical essence of the God Shiva.

(8) Shiva = pure consciousness (Here all dualities cease: See earlier models. Being= Becoming, self=Nonself order = Chaos etc.,)

(9) Shakti =Impure consciousness (Mother divine: responsible for the manifestation)

NOTATION

Module One

Tantric traditions are (=) thus those that aim at increasing (eb) the power of the practitioner (physical, mental awareness, retention capacity, concentration. "feeling good " about one self etc.,)

G_{13} : Category one of Tantric traditions

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of power of the practitioner

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Theological designation for the essence of such power is (=) Shakti (the female counterpart to the male divine principle, whose **essence is power: impure consciousness, one that is responsible for the manifestation of the phenomenal universe. We shall take this as string theory postulate with strings responsible for creation**).

G_{16} : Category one of Theological designation for the essence of such power

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Shakti (the female counterpart to the male divine principle, whose **essence is power: impure consciousness, one that is responsible for the manifestation of the phenomenal universe. We shall take this as string theory postulate with strings responsible for creation**).

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

manifestations of Shakti that the practitioner of tantra aspire after vary greatly, from relatively limited magical proficiencies (siddhis or vibhūtis), through royal power, to (eb) the deindividualized and liberated saint's omnipotence to the performance of God's cosmic acts(this is not para brahman; It is the knowledge and wisdom of the cosmic general ledger: Shakti is strings)

G_{20} : Category one of manifestations of Shakti that the practitioner of tantra aspire after vary greatly, from relatively limited magical proficiencies (siddhis or vibhūtis), through royal power

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of deindividualized and liberated saint's omnipotence to the performance of God's cosmic acts(this is not para brahman; It is the knowledge and wisdom of the cosmic general ledger: Shakti is strings)

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

What is God or Para Brahman's act? What is Karma? Take my case for example. I was a professional banker who took VRS, an aspiring star, and an academician who wanted to just make it to the top irrespective of the fact whether I had the abilities and capacities. I got in to a Chakravyuha like Abhimanyu, and lead a life of trials and tribulations and travails for a decade. What is that I did to be punished for this or still being punished? I will assume that God (Parà Brahman will take in to account (e) what I had done in my earlier incarnations (I donot think I did anything like that in this incarnation) wherein I could have tormented or produced a sense of deprivation balance either equivalent to or less than what I am going through now. Or God must have added interest and given me (eb) the punishment. I will assume both and give models. One with the deprivation balance equivalent to what I had done, and the other deprivation balance which includes the interest for the period between my act of commission and this incarnation. I will also take in to account the third possibility of Brahman-AntiBrahman who like spider weave the web, see the prey undergoing tormentations in the web and increase it just for "play"

G_{24} : Category one of actions done in my earlier incarnations (I donot think I did anything like that in this incarnation) wherein I could have tormented or produced a sense of deprivation balance either equivalent to or less than what I am going through now

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Parà Brahman

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

G_{28} : Category one of Parà Brahman adds interest(rate is simple, compounded, and astronomical)

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of punishment which is meeting of people who become responsible for increase in deprivation balance

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Violating prescriptions regarding caste, sexuality, diet and death, many of the tantric rites were originally performed in (eb) cremation grounds

G_{32} : Category one of prevarication of prescribed principles

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of consummation of given objective

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Whereas in Shākta tantrism, Shakti as a Goddess is herself the ultimate deity, in monistic Kashmiri Shaivism she is incorporated into (eb) the metaphysical essence of the God Shiva. Like zero contains all the transactions of the general ledger

G_{36} : Category one of shakti

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of shiva

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Shiva = pure consciousness (Here all dualities cease: See earlier models. Being= Becoming, self=Nonself order = Chaos etc.,)

G_{40} : Category one of pure consciousness

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of cessation of dualities'; space=time, being=non being, self=Nonself, chaos=order, gratification=deprivation

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

G_{44} : Category one of transcendence of sight and sound and sensory feelings

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Satchidananda or for that matter UN Satchidananda (they are the same)

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\ \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right] & \left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right] & \left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right] & \left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ \boxed{-(b_{24}'')^{(4)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{28}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{33}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{34}'')^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & \boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \qquad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & \boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \qquad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & \boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \qquad 81$$

$Where\ \boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b_{28}')^{(5)} & -(b_{28}'')^{(5)}(G_{31},t) & -(b_{24}'')^{(4,4,)}(G_{27},t) & -(b_{32}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{13}'')^{(1,1,1,1,1)}(G,t) & -(b_{16}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b_{29}')^{(5)} & -(b_{29}'')^{(5)}(G_{31},t) & -(b_{25}'')^{(4,4,)}(G_{27},t) & -(b_{33}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{14}'')^{(1,1,1,1,1)}(G,t) & -(b_{17}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{21}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b_{30}')^{(5)} & -(b_{30}'')^{(5)}(G_{31},t) & -(b_{26}'')^{(4,4,)}(G_{27},t) & -(b_{34}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{15}'')^{(1,1,1,1,1,)}(G,t) & -(b_{18}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{22}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b_{28}'')^{(5)}(G_{31},t)$, $-(b_{29}'')^{(5)}(G_{31},t)$, $-(b_{30}'')^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b_{24}'')^{(4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b_{32}'')^{(6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$-\left[\begin{matrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{matrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{matrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{matrix}(b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}\end{matrix}\right]T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{matrix}(b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}\end{matrix}\right]T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(A) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(B) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(C) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(D)

$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(E) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(F) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\big((G_{43}), t\big) = (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(a) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)},\ if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, if\ (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$

and $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)},\ if\ (\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad (378)$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad (379)$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad (380)$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- (381)

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 (382)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad (479)$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad (480)$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad (481)$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad (482)$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad (483)$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad (484)$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad (484A)$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: (485)

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: (486)

(a) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: (487)

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: (488)

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWO

Vijnāna-Bhairava Tantra

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) The basic pattern of spiritual practice, which also reflects the appropriation of Goddess worship (Shaktism) by Shaivism, is the approach to Shiva through (e&eb) Shakti. As the Shaiva scripture **Vijnāna-Bhairava** proclaims, Shakti is the **door**.
(2) The adept pursues the realization of identity with the omnipotent Shiva by assuming (eb) his mythic agency in emanating and controlling (e&eb) the universe through Shakti. Shakti = divided universe = Power
(3) Thus in the ritual a man realizes himself as (=) the possessor of Shakti within his partner.
(4) In more frequent internalized "theosophical" contemplations one realizes oneself as (=) the possessor of Shakti in all her immanent modalities with the aid of circular diagrams of cosmogenesis (mandalas) and mantras.
(5) An expression of this same process was (=) the production by monistic Shaiva Brahmans of increasingly systematic manuals of doctrines and practices on the model of Sanskrit scholastic texts (shāstras).
(6) This creation of what may be described as a religious mission to the educated elites also led (eb) to the increasing consolidation of the various streams of monistic Shaivism.
(7) This development began in the ninth century with Vasugupta's transmission of the manual Shiva Sūtra, ostensibly revealed to him by Shiva (Pure consciousness; cessation of all dualities space=time, order=chaos sound=silence) himself
(8) Further systematization of its teachings by either Vasugupta or his disciple Kallata in the Spanda Kārikā. These two works and their commentaries form the core texts of the "Spanda system" of monistic Shaivism, known for its interpretation of Shakti (manifested and divided universe attributable to mind: Neuron DNA see earlier models) as (=) Spanda, "cosmic pulsation." We shall take this as the pulsation or throbbing of strings as in string theory.

The tradition of monistic Shaivism called "Trika" (referring to its emphasis on various triads of modalities of Shakti and cosmic levels) produced the first work of full-fledged scholastic philosophy. This was the Shiva drishti, "Cognition of Shiva," by Somānanda (c. 900-950 C.E.). (See the summary of themes of the Shiva drishti below.) Utpaladeva, a student of Somānanda, wrote a commentary on the Shiva drishti, the Shivadrishtivritti. He also wrote several other works interpreting and furthering the work of Somānanda with much greater sophistication. Those texts are the foundational works of the Pratyabhijnā philosophy of focus in this article. The most comprehensive of these texts are the Īshvarapratyabhijnākārikā, "Verses on the Recognition of the Lord," and two commentaries on the Verses, the short Īshvarapratyabhijnākārikāvritti, and the more detailed Īshvarapratyabhijnāvivriti. (The latter text has been accessible to contemporary scholars only in fragments.) Utpaladeva also wrote a trilogy of more specialized philosophical studies, the Siddhitrayī, "Three Proofs"—Īshvarasiddhi, "Proof of the Lord;" Ajadapramātrisiddhi, "Proof of a Subject who is not Insentient;" and Sambandhasiddhi, "Proof of Relation." Abhinavagupta, widely recognized as one of the greatest philosophers of South Asia, was a disciple of a disciple of Utpaladeva. Abhinava profoundly elaborated and augmented Utpaladeva's arguments in long commentaries, one directly on the Verses, the Īshvarapratyabhijnāvimarshinī; and the other on Utpaladeva's longer auto commentary, the Īshvarapratyabhijnāvivritivimarshinī.

(9) While Abhinavagupta's Pratyabhijnā commentaries are of paramount philosophical importance, this thinker's greatest significance in the history of tantrism is probably his effort, in his monumental Tantrāloka and numerous other works, to systematize and provide (eb) a critical philosophical structure to non-philosophical tantric theology.

NOTATION

Module One

The basic pattern of spiritual practice, which also reflects the appropriation of Goddess worship (Shaktism) by Shaivism, is the approach to Shiva through (e&eb) Shakti. As the Shaiva scripture **Vijnāna-Bhairava** proclaims, Shakti is the **door**.

Pure consciousness is reached through impure consciousness

Paramaarthika(the ultimate transaction) is reached through arthika or loukika

G_{13} : Category one of Pure consciousness

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of reached through impure consciousness

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

adepts pursues the realization of identity with the omnipotent Shiva by assuming (eb) his mythic agency in emanating and controlling (e&eb) the universe through Shakti. Shakti = divided universe = Power

G_{16} : Category one of adepts pursues the realization of identity with the omnipotent Shiva by assuming (eb) his mythic agency in emanating and controlling; universe through Shakti. Shakti = divided universe = Power

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of universe through Shakti. Shakti = divided universe = Power ;adepts pursues the realization of identity with the omnipotent Shiva by assuming (eb) his mythic agency in emanating and controlling

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three.

G_{20} : Category one of Shakti

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of strings

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Internalized “theosophical” contemplations one realizes oneself as (=) the possessor of Shakti in all her immanent modalities with the aid of circular diagrams of cosmogenesis (mandalas) and mantras.

G_{24} : Category one of Internalized "theosophical" contemplations one realizes oneself

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of possessor of Shakti in all her immanent modalities with the aid of circular diagrams of cosmogenesis (mandalas) and mantras. **Man is essentially the flame of God. He is moving towards becoming one. Such an evolutionary state of consciousness and no mind condition is stated here**

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Expression of this same process was (=) the production by monistic Shaiva Brahmans of increasingly systematic manuals of doctrines and practices on the model of Sanskrit scholastic texts (shāstras).

G_{28} : Category one of Expression of this same process

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of manuals of doctrines and practices on the model of Sanskrit scholastic texts (shāstras) which is still evolving(my addition)

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Creation of what may be described as a religious mission to the educated elites also led (eb) to the increasing consolidation of the various streams of monistic Shaivism.

G_{32} : Category one of Creation of what may be described as a religious mission to the educated elites

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of consolidation of the various streams of monistic Shaivism. What is being tried is the path that would be followed by history had such an accentuation done in the holistic sense.

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Shiva (Pure consciousness; cessation of all dualities space=time, order=chaos sound=silence) himself

G_{36} : Category one of Pure consciousness (Shiva)

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of cessation of all dualities space=time, order=chaos sound=silence

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Further systematization of its teachings by either Vasugupta or his disciple Kallata in the Spanda Kārikā. These two works and their commentaries form the core texts of the "Spanda system" of monistic Shaivism, known for its interpretation of Shakti (manifested and divided universe attributable to mind: Neuron DNA see earlier models) as (=) Spanda, "cosmic pulsation." We shall take this as the pulsation or throbbing of strings as in string theory

G_{40} : Category one of Neuron DNA

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **Spanda, "cosmic pulsation." We shall take this as the pulsation or throbbing of strings as in string theory**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

While Abhinavagupta's Pratyabhijnā commentaries are of paramount philosophical importance, this thinker's greatest significance in the history of tantrism is probably his effort, in his monumental Tantrāloka and numerous other works, to systematize and provide (eb) a critical philosophical structure to non-philosophical tantric theology.

G_{44} : Category one of monumental Tantrāloka and numerous other works systematize and provide

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of critical philosophical structure to non-philosophical tantric theology

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} +(a_{16}'')^{(2)}(T_{17},t) +(a_{13}'')^{(1,1,)}(T_{14},t) +(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t) +(a_{40}'')^{(8,8,8)}(T_{41},t) +(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} +(a_{17}'')^{(2)}(T_{17},t) +(a_{14}'')^{(1,1,)}(T_{14},t) +(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t) +(a_{41}'')^{(8,8,8)}(T_{41},t) +(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} +(a_{18}'')^{(2)}(T_{17},t) +(a_{15}'')^{(1,1,)}(T_{14},t) +(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t) +(a_{42}'')^{(8,8,8)}(T_{41},t) +(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $\left[-(b_{44}'')^{(9)}(G_{47},t)\right]$, $\left[-(b_{45}'')^{(9)}(G_{47},t)\right]$, $\left[-(b_{46}'')^{(9)}(G_{47},t)\right]$ are first detrition coefficients for category 1, 2 and 3
$\left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)\right]$, $\left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)\right]$, $\left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)\right]$ are second detrition coefficients for category 1, 2 and 3
$\left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)\right]$, $\left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)\right]$, $\left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)\right]$ are third detrition coefficients for category 1, 2 and 3
$\left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)\right]$, $\left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)\right]$, $\left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)\right]$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(G) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(H) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(I) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(J)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(K) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(L) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \le (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)} =$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)} =$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(b) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}\ , G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(b) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THREE

Language As (=) The Sound (Nāda) Integral To Shiva's (Recall Assets=Liabilities Example Creative Power

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) Abhinava utilized categories from the Pratyabhijnā philosophy to interpret and organize (e&eb) the diverse aspects of doctrine and practice and Shaiva symbolism from the "Trika" sub-tradition; and he synthesized under the rubric of this philosophically rationalized Trika Shaivism an enormous range of symbolism and practice from other Shaiva and Shākta traditions as well. Abhinavagupta is also renowned for his works on Sanskrit poetics—in which he interpreted aesthetic experience as homologous to, and practically approaching the monistic Shaiva Soteriological realization.

(2) While the focus of this article is on Utpaladeva's and Abhinavagupta's Pratyabhijnā philosophy, mention should be made of some of the basic themes of Somānanda's precursory Shiva drishti. Somānanda's broadest concern is to explain how Shiva (Pure consciousness) through (e&eb) the various modalities of his Shakti (phenomenological manifestation of the universe) emanate (eb) a real universe that remains identical with himself.(Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger))

(3) In establishing the Shaiva doctrine he refutes (e) a number of alternative views on ultimate reality, the self, God and the metaphysical status of the world. He devotes the greatest polemical efforts against the theories of the 4th-6th century Vaiyākarana (or "Grammarian") philosopher Bhartrihari.

(4) According to Bhartrihari, the ultimate reality is (=) the Word Absolute (shabdabrahman)—a super-linguistic plenum, which fragments and emanates into (eb) the multiplicity of forms of expressive

speech and referents of that speech.

(5) Somānanda (S Theories) repudiates the view that a linguistic entity could be the ultimate reality, while at the same time identifying the true source of **language as (=) the Sound (nāda) integral to Shiva's (Recall Assets=Liabilities example creative power**; here gratification = deprivation)

(6) Somānanda takes a less polemical approach towards Shāktism. He argues that there is ultimately no difference between Shakti and Shiva, who is (=) the possessor of Shakti. This means that impure consciousness is a part of pure consciousness and emanated from (e) pure consciousness

(7) He supports this contention with the analogy of the inseparability of heat from fire, which is the possessor of heat. Nevertheless, he asserts that it is more proper to refer to the ultimate reality as Shiva rather than Shakti. Other Hindu schools criticized by Somānanda include the Pancarātra as well as the Vedānta, Sāmkhya and Nyāya-Vaisheshika systems.

(8) Somānanda briefly adduces some considerations against the Buddhist theory of momentariness, which were directly picked up and elaborated by Utpaladeva and Abhinavagupta. The most important of these was his advertence to the experience of recognition (Pratyabhijnā) as evidence both for the continuity of entities from the past through the present, and for the self that connects the past and (e&eb) present experiences of those entities.(Note we have taken self as witness consciousness the record keeper Neuron DNA)

(9) Somānanda's claims that synthetic categories or universals are (=) more primitive than particulars, and his invocation of Sanskrit syntax to explain Shiva's agency likewise had an important impact on Utpaladeva and Abhinavagupta. (See below.)

NOTATION

Module One

Abhinava (A Theories) utilized categories from the Pratyabhijnā philosophy to interpret and organize (e&eb) the diverse aspects of doctrine and practice and Shaiva symbolism from the "Trika" sub-tradition; and he synthesized under the rubric of this philosophically rationalized Trika Shaivism an enormous range of symbolism and practice from other Shaiva and Shākta traditions as well. Abhinavagupta is also renowned for his works on Sanskrit poetics—in which he interpreted aesthetic experience as homologous to, and practically approaching the monistic Shaiva Soteriological realization

G_{13} : Category one of A Theories utilized categories from the Pratyabhijnā philosophy to interpret and organize; diverse aspects of doctrine and practice and Shaiva symbolism from the "Trika" sub-tradition

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of diverse aspects of doctrine and practice and Shaiva symbolism from the "Trika" sub-tradition ;A Theories utilized categories from the Pratyabhijnā philosophy to interpret and organize

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

While the focus of this article is on Utpaladeva's and Abhinavagupta's Pratyabhijnā philosophy, mention should be made of some of the basic themes of Somānanda's precursory **Shiva drishti.** Somānanda's broadest concern is to explain how Shiva (Pure consciousness) through (e&eb) the various modalities of his Shakti (phenomenological manifestation of the universe) emanate (eb) a real universe that remains identical with himself.(Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or

giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger)

G_{16} : Category one of Shiva (Pure consciousness); various modalities of his Shakti (phenomenological manifestation of the universe) emanate (eb) a real universe that remains identical with himself.(Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger)

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of various modalities of his Shakti (phenomenological manifestation of the universe) emanate (eb) a real universe that remains identical with himself.(Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger); Shiva (Pure consciousness)

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

While the focus of this article is on Utpaladeva‘s and Abhinavagupta’s Pratyabhijnā philosophy, mention should be made of some of the basic themes of Somānanda’s precursory **Shiva drishti.** Somānanda’s broadest concern is to explain how Shiva (Pure consciousness) through the various modalities of his Shakti (phenomenological manifestation of the universe) emanate (eb) a real universe that remains identical with himself.(Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger)

G_{20} : Category one of Shiva (Pure consciousness) through the various modalities of his Shakti (phenomenological manifestation of the universe: set of string resonances as in string theory)

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of real universe that remains identical with himself. (Note Pure consciousness is then becoming from being in impure consciousness. Shakti like mother plays with her children, and when they get bored enters them, either relieving them of this world or giving them the developmental pure consciousness state as is in the case of Kundalini. process of birth and death goes on till you realise that I= cosmic consciousness (cosmic general ledger)

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

According to Bhartrihari (B Theories) the ultimate reality is the Word Absolute (shabdabrahman)—a super-linguistic plenum, which fragments and emanates into (eb) the multiplicity of forms of expressive speech and referents of that speech.

G_{24} : Category one of Bhartrihari (B Theories) the ultimate reality

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Word Absolute (shabdabrahman)—a super-linguistic plenum, which fragments and emanates into (eb) the multiplicity of forms of expressive speech and referents of that speech.

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Bhartrihari (B Theories) the ultimate reality is the Word Absolute (shabdabrahman)—a super-linguistic plenum, which fragments and emanates into (eb) the multiplicity of forms of expressive speech and referents of that speech

G_{28} : Category one of (B Theories) the ultimate reality is the Word Absolute (shabdabrahman)—a super-linguistic plenum, which fragments and emanates

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of multiplicity of forms of expressive speech and referents of that speech

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Somānanda (S Theories) repudiates the view that a linguistic entity could be the ultimate reality, while at the same time identifying the true source of **language as (=) the Sound (nāda) integral to Shiva's (Recall Assets=Liabilities example creative power; here gratification = deprivation)**

G_{32} : Category one of(S Theories) repudiates the view that a linguistic entity could be the ultimate reality, while at the same time identifying the true source of **language**

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of **Sound (nāda) integral to Shiva's (Pure consciousness) (Recall Assets=Liabilities example creative power**; here gratification = deprivation)

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Somānanda (S Theories) takes a less polemical approach towards Shāktism. He argues that there is ultimately no difference between Shakti and Shiva, who is (=) the possessor of Shakti. This means that impure consciousness is a part of pure consciousness and emanated from (e) pure consciousness

G_{36} : Category one of ultimately no difference between Shakti and Shiva

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of possessor of Shakti

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

impure consciousness is a part of pure consciousness and emanated from (e) pure consciousness

G_{40} : Category one of pure consciousness

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of impure consciousness is a part of pure consciousness and emanated

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Somānanda briefly adduces some considerations against the Buddhist theory of momentariness, which were directly picked up and elaborated by Utpaladeva and Abhinavagupta. The most important of these was his advertence to the experience of recognition (Pratyabhijnā) as evidence both for the continuity of entities from the past through the present, and for the self that connects the past and (e&eb) present experiences of those entities.(Note we have taken self as witness consciousness the record keeper Neuron DNA)

G_{44} : Category one of experience of recognition (Pratyabhijnā) as evidence both for the continuity of entities from the past through the present, and for the self connects the past; present experiences of those entities.(Note we have taken self as witness consciousness the record keeper Neuron DNA)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of present experiences of those entities.(Note we have taken self as witness consciousness the record keeper Neuron DNA);experience of recognition (Pratyabhijnā) as evidence both for the continuity of entities from the past through the present, and for the self connects the past

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) \quad +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) \quad +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) \quad +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b_{20}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{16}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{13}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{15}'')^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{36}'')^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b_{38}'')^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{24}')^{(4)} & \boxed{+(a_{24}'')^{(4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a_{25}')^{(4)} & \boxed{+(a_{25}'')^{(4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{33}'')^{(6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{26}')^{(4)} & \boxed{+(a_{26}'')^{(4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{26} \quad (75)$$

$\boxed{(a_{24}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{25}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{26}'')^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)}$ *are fourth augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)}$ *are fifth augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)}$

are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$

are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \tag{76}$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \tag{77}$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{26} \tag{78}$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\begin{bmatrix}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{28} \tag{79}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\,,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients*

$for\ category$ 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ $are\ \ third\ \ detrition\ coefficients$ $for\ category$ 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t)\\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t)\\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t)\\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{34} \qquad 87$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ $are\ \ first\ \ augmentation\ coefficients$ $for\ category$ 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ $are\ second\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ $are\ third\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$
seventh augmentation coefficients
$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$
Eighth augmentation coefficients
$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt}=(b_{32})^{(6)}T_{33}-\begin{bmatrix}(b'_{32})^{(6)}\,-(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t)\\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt}=(b_{33})^{(6)}T_{32}-\begin{bmatrix}(b'_{33})^{(6)}\,-(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t)\\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt}=(b_{34})^{(6)}T_{33}-\begin{bmatrix}(b'_{34})^{(6)}\,-(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t)\\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3
$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3
$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(M) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(N) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(O) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(P)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(Q) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(R) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}$, $T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(c) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t} \right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)} t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t} \right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^{0}}{G_{14}^{0}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(c) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FOUR

Pure Consciousness

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) Also noteworthy is Somānanda's (S Theories) advocacy of a "panpsychist" theory that all things, which emanate from (e) the consciousness of Shiva, have their own consciousness and agency.
(2) Somānanda additionally engages in reflecting on the contemplations that lead to (eb) the realization of identity with Shiva.
(3) Note here that when shiva himself contemplates on his own identity the universe emanates. It is something like reflecting on zero which we write with much flourish and preen with every equation and all the transactions in the general ledger including qualitative which has been quantified.
(4) Utpaladeva (U Theories) and Abhinavagupta (A Theories) ambitiously conceive the Pratyabhijnā system as (=) both a philosophical apologetics (which follows Sanskritic standards of scholastic argument) and an internalized form of tantric ritual that leads students directly to identification with Shiva **(Pure consciousness)**.
(5) They explain the basic means by which the system conveys Shiva-identity according to the same basic ritual pattern described above, as (=) shaktyāvishkarana, "the revealing of Shakti." We must understand here that when Shiva, the pure consciousness reflects on his own self (zero) then Shakti is manifested, namely the phenomenal universe is manifested in all its dualities and conflicts.
(6) The Pratyabhijnā philosophers, however, also frame Shakti as (=) the reason of a publicly assessable inference, or "inference for the sake of others" (parārthānumāna).
(7) According to the scholastic logic, the reason identifies (e&eb) a quality in the inferential subject "I" known to be invariably concomitant (e&eb) with the predicate, "Shiva."
(8) Thus I am Shiva because (e) I have his quality, that is, Shakti, the capacity of emanating and controlling the universe. This is most important and the is interpreted as Satchidananda(mine I = cosmic consciousness)
(9) Terrestial Brahman Anti Brahman or celestial Brahman-AntiBrahman has the quality but they are marginalized by Para Brahman.

NOTATION

Module One

Also noteworthy is Somānanda's (S Theories) advocacy of a "panpsychist" theory that all things, which emanate from (e) the consciousness of Shiva, have their own consciousness and agency

G_{13} : Category one of consciousness and agency

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of (S Theories) advocacy of a "panpsychist" theory that all things, which emanate from (e) the consciousness of Shiva , the pure consciousness

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Somānanda (S Theories) additionally engages in reflecting on the contemplations that lead to (eb) the realization of identity with Shiva, the pure consciousness where all realities cease. As said, the sole aim is to become God in the real sense of term. That is where we are moving with all flying cars, superconductivity, motion capturing technologies etc., Reflection is essentially a journey in to inner self. When pure consciousness, zero reflects on its own self, then the disintegration takes place.

G_{16} : Category one of(S Theories) additionally engages in reflecting on the contemplations

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of realization of identity with Shiva, the pure consciousness where all realities cease. As said, the sole aim is to become God in the real sense of term. That is where we are moving with all flying cars, superconductivity, motion capturing technologies etc., Reflection is essentially a journey in to inner self. When pure consciousness, zero reflects on its own self, then the disintegration takes place

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Shiva himself contemplates on his own identity the universe emanates. It is something like reflecting on zero which we write with much flourish and preen with every equation and all the transactions in the general ledger including qualitative which has been quantified

G_{20} : Category one of Shiva himself contemplates on his own identity

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of universe emanates. It is something like reflecting on zero which we write with much flourish and preen with every equation and all the transactions in the general ledger including qualitative which has been quantified

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Utpaladeva (U Theories) and Abhinavagupta (A Theories) ambitiously conceive the Pratyabhijnā system as (=) both a philosophical apologetics (which follows Sanskritic standards of scholastic argument) and an internalized form of tantric ritual that leads students directly to identification with Shiva **(Pure consciousness)**.

G_{24} : Category one of Utpaladeva (U Theories) and Abhinavagupta (A Theories) ambitiously conceive the Pratyabhijnā system

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of philosophical apologetics (which follows Sanskritic standards of scholastic argument) and an internalized form of tantric ritual that leads students directly to identification with Shiva **(Pure consciousness)**.

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

They explain the basic means by which the system conveys Shiva-identity according to the same basic ritual pattern described above, as (=) shaktyāvishkarana, "the revealing of Shakti." We must understand here that when Shiva, the pure consciousness reflects on his own self (zero) then Shakti is manifested, namely the phenomenal universe is manifested in all its dualities and conflicts.

G_{28} : Category one of basic means by which the system conveys Shiva-identity according to the same basic ritual pattern described above

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of shaktyāvishkarana, "the revealing of Shakti(the manifestation of phenomenal universe)." We must understand here that when Shiva, the pure consciousness reflects on his own self (zero) then Shakti is manifested, namely the phenomenal universe is manifested in all its dualities and conflicts

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Pratyabhijnā philosophers, however, also frame Shakti as (=) the reason of a publicly assessable inference, or "inference for the sake of others" (parārthānumāna)

G_{32} : Category one of Pratyabhijnā philosophers, however, also frame Shakti

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of reason of a publicly assessable inference, or "inference for the sake of others" (parārthānumāna)

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

According to the scholastic logic, the reason identifies a quality in the inferential subject "I" known to be invariably concomitant (e&eb) with the predicate, "Shiva."

G_{36} : Category one of reason; quality in the inferential subject "I" known to be invariably concomitant (e&eb) with the predicate, "Shiva."

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of quality in the inferential subject "I" known to be invariably concomitant (e&eb) with the predicate, "Shiva.";reason

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Reason identifies a quality in the inferential subject "I" known to be invariably concomitant (e&eb) with the predicate, "Shiva."

G_{40} : Category one of Reason identifies a quality in the inferential subject "I" known to be invariably concomitant; "Shiva."

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of "Shiva."; Reason identifies a quality in the inferential subject "I" known to be invariably concomitant

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Thus I am Shiva because (e) I have his quality, that is, Shakti, the capacity of emanating and controlling the universe. This is most important and the is interpreted as Satchidananda(mine I = cosmic consciousness)

G_{44} : Category one of I have his quality that is, Shakti, the capacity of emanating and controlling the

universe. This is most important and the is interpreted as Satchidananda(mine I = cosmic consciousness)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of I

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition

coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(S) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(T) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(U) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(V)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(W) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(X) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\Big] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(d) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(d) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIVE

(U-A Theories) Interpret Their Central Myth Of Shiva's Emanation And Control Of The Universe Through Shakti As Itself An Act Of Self-Recognition

(Ahampratyavamarsha, Pratyabhijnā).

INTRODUCTION—VARIABLES USED

Phenomenology, developed by Edmund Husserl (1859–1938), can be understood as an outgrowth of the influential ideas of Immanuel Kant (1724-1804). Attempting to resolve some of the key intellectual debates of his era, Kant argued that Noumena (objects of inquiry, investigation or analysis of the workings of the world) must be distinguished from Phenomena (noumenon as experienced and interpreted by the human mind). Kant thus argued that humans can never have direct access to reality, but only to the contents of their minds. Bracketing thus involves setting aside the question of the real existence of a contemplated object, as well as all other questions about the object's physical or objective nature; these questions are left to the natural sciences. For example, the act of seeing a horse qualifies as a mental experience, whether one sees the horse in person, in a dream, or in a hallucination. 'Bracketing' the horse suspends any judgement about the horse as noumenon, and instead analyses the phenomenon of the horse in the human mind. Bracketing has been compared to the instructions given to a jury in a trial to analyze only the evidence presented in court, and to avoid the usual inferences or connections they might make between such evidence and other sources of information. Bracketing can also be better understood in terms of the phenomenological activity it is supposed to make possible: the "unpacking" of phenomena, or, in other words, systematically peeling away their symbolic meanings like layers of an onion until only the thing itself as meant and experienced remains. Thus, one's subjective perception of the bracketed phenomenon is examined and analyzed in its purity.(Wikipedia)

Tantra and Kashmiri Shaivism (Wikipedia)

(1) That is why you see destruction of powerful empires, Tsunamis, earthquakes, and other natural disasters like supernovae, gamma rays etc.,

The Pratyabhijnā Epistemology

(2) In order to address debates on epistemology that were then current, Utpaladeva and Abhinavagupta further explain the mythic and ritual pattern of Shiva and Shakti in terms of recognition. The specific problem the writers address had been formulated by the Buddhist logic school of Dignāga and Dharmakīrti, which flourished in medieval Kashmir. Contemporary interpreters have characterized the philosophy of Buddhist logic as (=) a species of phenomenalism akin to that of David Hume (D Theories).

(3) According to this school, the foundation of knowledge is (=) a series of momentary and discrete perceptual data (svalakshana).

(4) There are no grounds in those data for the recognitions of any enduring entities through ostensible cognitions utilizing (e) linguistic or conceptual interpretation (savikalpaka jñāna).

(5) In debates over several centuries, the Buddhist (B-Theories) logicians had propounded arguments attacking (e) many concepts that seemed commonsensical and were religiously significant to the various orthodox Hindu philosophical schools—such as ideas of external objects, ordinary and ritual action, an enduring Self, God, and revelation.

(6) The Pratyabhijnā philosophers' response to the problematic posed by Buddhist logic revolutionized earlier approaches of the Nyaya philosophers, the Shaiva Siddhāntin Sadyojyoti and even Utpaladeva's teacher Somānanda, and may be characterized (eb) as a form of transcendental argumentation.

(7) Utpaladeva and Abhinavagupta **(U-A Theories) interpret (eb) their central myth of Shiva's emanation and control of the universe through (e&eb) Shakti as (=) itself an act of self-recognition (ahampratyavamarsha, Pratyabhijnā).**

(8) Furthermore, abjuring Somānanda's (S-Theories) agonistic stance towards Bhartrihari (B

Theories) they also equate Shiva's self-recognition (Shakti) with (=) the principle of Supreme Speech (parāvāk), which they derive from the Grammarian. This I interpret as ultimate expression "Om"

(9) They thereby appropriate the Grammarian's explanation of creation as (=) linguistic in nature.

NOTATION

Module One

In order to address debates on epistemology that were then current, Utpaladeva and Abhinavagupta further explain the mythic and ritual pattern of Shiva and Shakti in terms of recognition. The specific problem the writers address had been formulated by the Buddhist logic school of Dignāga and Dharmakīrti, which flourished in medieval Kashmir. Contemporary interpreters have characterized the philosophy of Buddhist logic as (=) a species of phenomenalism akin to that of David Hume (D Theories).

G_{13} : Category one of philosophy of Buddhist logic

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of species of phenomenalism akin to that of David Hume (D Theories).

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Foundation of knowledge is (=) a series of momentary and discrete perceptual data (svalakshana).

G_{16} : Category one of Foundation of knowledge

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of series of momentary and discrete perceptual data (svalakshana).

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

There are no grounds in those data for the recognitions of any enduring entities through ostensible cognitions utilizing (e) linguistic or conceptual interpretation (savikalpaka jñāna)

G_{20} : Category one of linguistic or conceptual interpretation (savikalpaka jñāna)

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of groundlessness in those data for the recognitions of any enduring entities through ostensible cognitions

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

In debates over several centuries, the Buddhist (B-Theories) logicians had propounded arguments attacking (e) many concepts that seemed commonsensical and were religiously significant to the various orthodox Hindu philosophical schools—such as ideas of external objects, ordinary and ritual action, an enduring Self, God, and revelation.

G_{24} : Category one of concepts that seemed commonsensical and were religiously significant to the various orthodox Hindu philosophical schools—such as ideas of external objects, ordinary and ritual action, an enduring Self, God, and revelation

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Buddhist (B-Theories) logicians had propounded arguments

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Pratyabhijnā philosophers' response to the problematic posed by Buddhist logic revolutionized earlier approaches of the Nyaya philosophers, the Shaiva Siddhāntin Sadyojyoti and even Utpaladeva's teacher Somānanda, and may be characterized (eb) as a form of transcendental argumentation.

G_{28} : Category one of Shaiva Siddhāntin Sadyojyoti and even Utpaladeva's teacher Somānanda, and may be characterized

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of transcendental argumentation

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Pratyabhijnā philosophers' response to the problematic posed by Buddhist logic revolutionized earlier approaches of the Nyaya philosophers, the Shaiva Siddhāntin Sadyojyoti and even Utpaladeva's teacher Somānanda, and may be characterized (eb) as a form of transcendental argumentation.

G_{32} : Category one of Pratyabhijnā philosophers' response to the problematic posed by Buddhist logic revolutionized earlier approaches of the Nyaya philosophers, the Shaiva Siddhāntin Sadyojyoti

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of form of transcendental argumentation

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

(U-A Theories) interpret their central myth of Shiva's emanation and control of the universe through Shakti as (=) itself an act of self-recognition (ahampratyavamarsha, Pratyabhijnā).

G_{36} : Category one of **(U-A Theories) interpret their central myth of Shiva's emanation and control of the universe through Shakti**

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of **act of self-recognition (ahampratyavamarsha, Pratyabhijnā).**

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Furthermore, abjuring Somānanda's (S-Theories) agonistic stance towards Bhartrihari (B Theories) they also equate Shiva's self-recognition (Shakti) with (=) the principle of Supreme Speech (parāvāk), which they derive from the Grammarian. This I interpret as ultimate expression "Om"

G_{40} : Category one of Shiva's self-recognition (Shakti)

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of principle of Supreme Speech (parāvāk), which they derive from the Grammarian. This I interpret as ultimate expression "Om"

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

They thereby appropriate the Grammarian's explanation of creation as (=) linguistic in nature.

G_{44} : Category one of Grammarian's explanation of creation

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of linguistic in nature

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a_{20}')^{(3)}\,+(a_{20}'')^{(3)}(T_{21},t)\,+(a_{16}'')^{(2,2,2)}(T_{17},t)\,+(a_{13}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{40}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{44}'')^{(9,9,9)}(T_{45},t) \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a_{21}')^{(3)}\,+(a_{21}'')^{(3)}(T_{21},t)\,+(a_{17}'')^{(2,2,2)}(T_{17},t)\,+(a_{14}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{41}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{45}'')^{(9,9,9)}(T_{45},t) \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a_{22}')^{(3)}\,+(a_{22}'')^{(3)}(T_{21},t)\,+(a_{18}'')^{(2,2,2)}(T_{17},t)\,+(a_{15}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{42}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{46}'')^{(9,9,9)}(T_{45},t) \end{array}\right] G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b_{20}')^{(3)}\,-(b_{20}'')^{(3)}(G_{23},t)\,-(b_{16}'')^{(2,2,2)}(G_{19},t)\,-(b_{13}'')^{(1,1,1,)}(G,t) \\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)\,-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)\,-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t)\,-(b_{40}'')^{(8,8,8,8)}(G_{43},t)\,-(b_{44}'')^{(9,9,9)}(G_{47},t) \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b_{21}')^{(3)}\,-(b_{21}'')^{(3)}(G_{23},t)\,-(b_{17}'')^{(2,2,2)}(G_{19},t)\,-(b_{14}'')^{(1,1,1,)}(G,t) \\ -(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)\,-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)\,-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7,7)}(G_{39},t)\,-(b_{41}'')^{(8,8,8,8)}(G_{43},t)\,-(b_{45}'')^{(9,9,9)}(G_{47},t) \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(Y) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(Z) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(AA) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(BB)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(e) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\leq G_{15}(t)\leq\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\leq T_{15}(t)\leq \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\leq-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\leq-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\leq-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\leq-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$ 318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(e) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIX

Prakāsha Is The "Bare Subjective Awareness" That Validates Each Cognition, So That One Knows That One Knows

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) Thus the Kashmiri Shaiva philosophers ascribe (e) to Speech a primordial status, denied by the Buddhist logicians.

(2) As ritual recapitulates myth, the Pratyabhijnā system endeavors to lead (eb) the student to participate in the recognition "I am Shiva," by demonstrating that all experiences and contents of experience are (=) expressions of the recognition that "I am Shiva." I interpret here again this means that I= cosmic consciousness the cosmic general ledger and by participation I am contributing and infact everyone even a insentient stone is contributing to " Grand design"

(3) The paradox of the Pratyabhijnā formulation of the inference for the sake of others is that the self-recognition "I am Shiva," as an interpretation of Shakti, becomes (=) in effect both the conclusion and the reason. There is some confusion here. I clarify that Shiva is pure consciousness from which impure consciousness Shakti emanates in division of phenomenal universe. Like there are transactions in zero, Shakti exists inside shiva. This is not the reason, but the very conservation law like thermodynamical or assets =- liabilities. However we continue with the assumed conclusion and reason theory further

(4) This circularity of conclusion and reason is a consequence (e) of the Kashmiri Shaiva monism. From the intratraditional perspective, there is no fact that can be adduced in support of (eb) another separate fact, as everything is (=) always the same in essential nature.

(5) From the intertraditional perspective of philosophical debate, however, the circularity is (=) not necessarily destructive.

(6) The Shaiva technical studies of various topics of epistemology and ontology in effect provide (eb) further ostensible justification for this apparent circularity.

(7) Utpaladeva's (U Theories) and Abhinavagupta's (A Theories) epistemology may best be illustrated

by (e) its approach to perceptual cognition.

(8) The Pratyabhijnā arguments on this subject may be divided into those centered around two sets of terms: prakashan; and vimarsha and cognates such as pratyavamarsha and parāmarsha.

(9) Prakāsha is the "bare subjective awareness" that validates (eb) each cognition, so that one knows (e) that one knows.

NOTATION

Module One

Kashmiri Shaiva philosophers ascribe (e) to Speech a primordial status, denied by the Buddhist logicians.

G_{13} : Category one of Speech a primordial status, denied by the Buddhist logicians

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Kashmiri Shaiva philosophers

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Speech a primordial status, denied by the Buddhist logicians

G_{16} : Category one of Speech

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of primordial status

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

As ritual recapitulates myth, the Pratyabhijnā system endeavors to lead (eb) the student to participate in the recognition "I am Shiva," by demonstrating that all experiences and contents of experience are (=) expressions of the recognition that "I am Shiva." I interpret here again this means that I= cosmic consciousness the cosmic general ledger and by participation I am contributing and infact everyone even a insentient stone is contributing to " Grand design"

G_{20} : Category one of I

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of cosmic general ledger

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

G_{24} : Category one of experiences and contents of experience

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of expressions of the recognition that "I am Shiva (Pure consciousness)

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

This circularity of conclusion and reason is a consequence (e) of the Kashmiri Shaiva monism. From the intratraditional perspective, there is no fact that can be adduced in support of (eb) another separate fact, as everything is (=) always the same in essential nature

G_{28} : Category one of Kashmiri Shaiva monism

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of circularity of conclusion and reason

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

everything is (=) always the same in essential nature

G_{32} : Category one of everything

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of same in essential nature

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

The Shaiva technical studies of various topics of epistemology and ontology in effect provide (eb) further ostensible justification for this apparent circularity

G_{36} : Category one of Shaiva technical studies of various topics of epistemology and ontology

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of ostensible justification for this apparent circularity

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Prakāsha is the "bare subjective awareness" that validates (eb) each cognition, so that one knows (e) that one knows.

G_{40} : Category one of **Prakāsha is the "bare subjective awareness"**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **each cognition, so that one knows (e) that one knows.**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Each cognition so that one knows that one knows.

G_{44} : Category one of **so that one knows that one knows.**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **bare subjective awareness"**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)} \ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients

for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)}\,+(a_{16}'')^{(2)}(T_{17},t)\,+(a_{13}'')^{(1,1,)}(T_{14},t)\,+(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)\,+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)\,+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t)\,+(a_{40}'')^{(8,8,8)}(T_{41},t)\,+(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)}\,+(a_{17}'')^{(2)}(T_{17},t)\,+(a_{14}'')^{(1,1,)}(T_{14},t)\,+(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)\,+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)\,+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t)\,+(a_{41}'')^{(8,8,8)}(T_{41},t)\,+(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)}\,+(a_{18}'')^{(2)}(T_{17},t)\,+(a_{15}'')^{(1,1,)}(T_{14},t)\,+(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)\,+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)\,+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t)\,+(a_{42}'')^{(8,8,8)}(T_{41},t)\,+(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for

category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition*

coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$ $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a_{24}')^{(4)}+(a_{24}'')^{(4)}(T_{25},t) & +(a_{28}'')^{(5,5,)}(T_{29},t) & +(a_{32}'')^{(6,6,)}(T_{33},t)\\ +(a_{13}'')^{(1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{36}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9)}(T_{45},t)\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a_{25}')^{(4)}+(a_{25}'')^{(4)}(T_{25},t) & +(a_{29}'')^{(5,5,)}(T_{29},t) & +(a_{33}'')^{(6,6)}(T_{33},t)\\ +(a_{14}'')^{(1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{37}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9)}(T_{45},t)\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a_{26}')^{(4)}+(a_{26}'')^{(4)}(T_{25},t) & +(a_{30}'')^{(5,5,)}(T_{29},t) & +(a_{34}'')^{(6,6,)}(T_{33},t)\\ +(a_{15}'')^{(1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{38}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9)}(T_{45},t)\end{bmatrix}G_{26} \quad (75)$$

$(a_{24}'')^{(4)}(T_{25},t)$, $(a_{25}'')^{(4)}(T_{25},t)$, $(a_{26}'')^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a_{28}'')^{(5,5,)}(T_{29},t)$, $+(a_{29}'')^{(5,5,)}(T_{29},t)$, $+(a_{30}'')^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a_{32}'')^{(6,6,)}(T_{33},t)$, $+(a_{33}'')^{(6,6,)}(T_{33},t)$, $+(a_{34}'')^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a_{13}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{16}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{20}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* $1,2$ *and* 3

$+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{46}'')^{(9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} & \boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} & \boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} & \boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad 84$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5)}(G_{31},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients

for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{matrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{matrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{matrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (92)$$

$\frac{dG_{38}}{dt}$ *93*

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third

augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth
augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth
augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth
augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh
augmentation coefficient for 1,2,3
$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$,$+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$,$+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth
augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth
augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)} & -(b''_{44})^{(9)}(G_{47},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)} & -(b''_{45})^{(9)}(G_{47},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)} & -(b''_{46})^{(9)}(G_{47},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15$ *97*

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$

$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2 \to \infty}(a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)}$ 98

$\lim_{G \to \infty}(b_i'')^{(1)}(G, t) = (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15$,
satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \tag{117}$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \tag{118}$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: (119)

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: (120)

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: (121)

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$

Where we suppose

(EE) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(FF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(GG) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(HH)

$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(II) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(JJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants

$(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\left((G_{43}), t\right) = \ (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\left((G_{43}), t\right)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, \ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , $T_i : \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{13})^{(1)}$, $T_i^0 \leq (\hat{Q}_{13})^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{24})^{(4)}$, $T_i^0 \leq (\hat{Q}_{24})^{(4)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(f) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \tag{180}$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right] \tag{181}$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose $\tag{182}$

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric $\tag{185}$

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\le \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 19 to 24 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\le\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t),t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\le \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)},i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 216

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(3)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3:$ 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\ \ t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b''_{22})^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\le \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right),\quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\le \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 250

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(6)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1,\left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right),\ \ t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying

Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:(\widetilde{(G_{39})},\widetilde{(T_{39})})=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)})d\left(((G_{39})^{(1)},(T_{39})^{(1)};(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 261

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(7)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

268

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{40})^{(8)}\right]\leq(\hat{Q}_{40})^{(8)}$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying
$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\left(u^{(2)}\right)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)}=(\nu_0)^{(2)},(m_1)^{(2)}=(\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)}<(\nu_1)^{(2)}$ 305

$(m_2)^{(2)}=(\nu_1)^{(2)},(m_1)^{(2)}=(\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)}<(\nu_0)^{(2)}<(\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)}=\frac{\mathrm{G}_{16}^{0}}{\mathrm{G}_{17}^{0}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})}\left[e^{((S_1)^{(3)}-(p_{20})^{(3)})t}-e^{-(S_2)^{(3)}t}\right]+G_{22}^{0}e^{-(S_2)^{(3)}t}\le G_{22}(t)\le\right. \quad 324$$

$$\left.\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}\right]+G_{22}^{0}e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le\frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\le T_{22}(t)\le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$

$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$

$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)}\le-(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\le-(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)}\le-(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\le-(\tau_1)^{(4)}$

Definition of $(\nu_1)^{(4)},(\nu_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},\nu^{(4)},u^{(4)}$: 329

By $(\nu_1)^{(4)}>0$, $(\nu_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\left(\nu^{(4)}\right)^2+(\sigma_1)^{(4)}\nu^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{\nu}_1)^{(4)},,(\bar{\nu}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)}>0$, $(\bar{\nu}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\left(\nu^{(4)}\right)^2+(\sigma_2)^{(4)}\nu^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},\ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, ***if*** $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, ***if*** $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, ***if*** $(\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$
$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)},, (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40} e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40} e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}$, $(\mu_2)^{(9)}$, $(v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}$, $(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, $\boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, $\boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$, and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$, and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}\left((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)}\right)}\left[e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} - e^{-(S_2)^{(9)}t}\right] + G_{46}^0 e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}\left((S_1)^{(9)} - (a'_{46})^{(9)}\right)}\left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}\left((R_1)^{(9)} - (b'_{46})^{(9)}\right)}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}\left((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)}\right)}\left[e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \leq \frac{dv^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \leq (\bar{v}_1)^{(1)}$$

386

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(\mathrm{C})^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(\mathrm{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{\mathrm{C}})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(\mathrm{C})^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(\mathrm{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right) - (a''_{25})^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

415

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a_{36}')^{(7)} - (a_{37}')^{(7)} + (a_{36}'')^{(7)}(T_{37}, t)\right) - (a_{37}'')^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition

$(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $v^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq v^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0$ 475

$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0$ 476

$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0$ 477

$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0$ 478

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(f) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T_{14}^{*} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^{*} for which $f(T_{14}^{*}) = 0$. With this value , we obtain from the three first

equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that

there exists a unique T_{37}^{*} for which $f(T_{37}^{*})=0$. With this value , we obtain from the three first equations

$$G_{36}=\frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}=\frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0)<0, f(\infty)>0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*})=0$. With this value , we obtain from the three first equations

$$G_{40}=\frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}=\frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0)<0, f(\infty)>0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*})=0$. With this value , we obtain from the three first equations

$$G_{44}=\frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}=\frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G)=(b'_{13})^{(1)}(b'_{14})^{(1)}-(b_{13})^{(1)}(b_{14})^{(1)}-$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G)+(b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G)=0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0)>0$, $\varphi(\infty)<0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*})=0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19})=(b'_{16})^{(2)}(b'_{17})^{(2)}-(b_{16})^{(2)}(b_{17})^{(2)}-$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19})+(b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19})=0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0)>0$, $\varphi(\infty)<0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*})=0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23})=(b'_{20})^{(3)}(b'_{21})^{(3)}-(b_{20})^{(3)}(b_{21})^{(3)}-$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]}$, $T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]}$ 517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]}$, $G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$\frac{\partial(a_{17}'')^{(2)}}{\partial \mathrm{T}_{17}}(\mathrm{T}_{17}^{*}) = (q_{17})^{(2)}$, $\frac{\partial(b_i'')^{(2)}}{\partial \mathrm{G}_j}(\,(G_{19})^{*}\,) = s_{ij}$ 533

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{\mathrm{d}\mathbb{G}_{16}}{\mathrm{dt}} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}\mathrm{G}_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{\mathrm{d}\mathbb{G}_{17}}{\mathrm{dt}} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}\mathrm{G}_{17}^{*}\mathbb{T}_{17} \quad 535$$

$$\frac{\mathrm{d}\mathbb{G}_{18}}{\mathrm{dt}} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}\mathrm{G}_{18}^{*}\mathbb{T}_{17} \quad 536$$

$$\frac{\mathrm{d}\mathbb{T}_{16}}{\mathrm{dt}} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \textstyle\sum_{j=16}^{18}\big(s_{(16)(j)}\mathrm{T}_{16}^{*}\mathbb{G}_j\big) \quad 537$$

$$\frac{\mathrm{d}\mathbb{T}_{17}}{\mathrm{dt}} = -\big((b_{17}')^{(2)} - (r_{17})^{(2)}\big)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \textstyle\sum_{j=16}^{18}\big(s_{(17)(j)}\mathrm{T}_{17}^{*}\mathbb{G}_j\big) \quad 538$$

$$\frac{\mathrm{d}\mathbb{T}_{18}}{\mathrm{dt}} = -\big((b_{18}')^{(2)} - (r_{18})^{(2)}\big)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \textstyle\sum_{j=16}^{18}\big(s_{(18)(j)}\mathrm{T}_{18}^{*}\mathbb{G}_j\big) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\,\mathbb{R}_{+})$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^{*}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}(\,(G_{23})^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\big((a_{20}')^{(3)} + (p_{20})^{(3)}\big)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^{*}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\big((a_{21}')^{(3)} + (p_{21})^{(3)}\big)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^{*}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\big((a_{22}')^{(3)} + (p_{22})^{(3)}\big)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^{*}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\big((b_{20}')^{(3)} - (r_{20})^{(3)}\big)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \textstyle\sum_{j=20}^{22}\big(s_{(20)(j)}T_{20}^{*}\mathbb{G}_j\big) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\big((b_{21}')^{(3)} - (r_{21})^{(3)}\big)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \textstyle\sum_{j=20}^{22}\big(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\big) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\big((b_{22}')^{(3)} - (r_{22})^{(3)}\big)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \textstyle\sum_{j=20}^{22}\big(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\big) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}\ and\ (b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \ , \ \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}\ and\ (b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^*\right)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left(((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left(((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)})(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(21)}T^{*}_{21} + (b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)})(q_{20})^{(3)}G^{*}_{20} + (a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(20)}T^{*}_{21} + (b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}\right)^{2} + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2} + \left((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2} + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(22)}T^{*}_{21} + (b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVEN

Everything Is Consciousness And Consciousness Is Everything

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) **The thrust of the arguments about Prakāsha is analogous to (=) George Berkeley's thesis (G Theories) of idealism that esse est percipio.**
(2) **The Shaivas contend that, as no object is known without (e) validating awareness, this awareness actually constitutes (e) all objects.**
(3) **There is no ground even for (e) a "representationalist" inference of objects external to awareness that causes (eb) its diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here**
(4) **Furthermore, the Kashmiri Shaivas argue that there cannot be another subject outside (e) of one's own awareness. They conclude, however, not with solipsism as usually understood in the West, but a conception of a universal awareness.**
(5) **All sentient and insentient beings are (=) essentially one awareness**.
(6) **Vimarsha a**nd its cognates have (e) the significance of apprehension or judgment with a recognitive structure, and may be glossed as (=) "recognitive apprehension."
(7) The recognitive is (=) the act of recognizing or an awareness that something perceived (e) has been perceived before.
(8) Utpaladeva's (U Theories) and Abhinavagupta's (A Theories) arguments centering on these terms develop earlier considerations of Bhartrihari (B Theories) on the linguistic nature of experience.
(9) Utpaladeva (U Theories) and Abhinavagupta (A Theories) refute (e) the Buddhist contention that recognition is (=) a contingent reaction to direct experience by (e) claiming that it is (=) integral or transcendental to all experience.

NOTATION

Module One

Thrust of the arguments about Prakāsha is analogous to (=) George Berkeley's thesis (G Theories) of idealism that esse est percipio

G_{13} : Category one of **Thrust of the arguments about Prakāsha**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **George Berkeley's thesis (G Theories) of idealism that esse est percipio**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Shaivas contend that, as no object is known without validating awareness, this awareness actually constitutes (e) all objects

G_{16} : Category one of **validating awareness, this awareness actually constitutes (e) all objects**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of **no object is known**

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

no object is known without validating awareness, this awareness actually constitutes (e) all objects

G_{20} : Category one of all objects

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of no object is known without validating awareness, this awareness

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Groundlessness for a "representationalist" inference of objects external to awareness that causes (eb) its diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here

G_{24} : Category one of **"representationalist" inference of objects external to awareness that causes (eb) its diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **Groundlessness**

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Groundlessness for a "representationalist" inference of objects external to awareness that causes its diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here

G_{28} : Category one of **Groundlessness for a "representationalist" inference of objects external to awareness**

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of **diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here**

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Groundlessness for a "representationalist" inference of objects external to awareness that causes its diverse contents, because (e) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here

G_{32} : Category one of**) causality can be posited only between phenomena of which one has been aware. This means everything is consciousness and consciousness is everything there is no conclusion and reason circularity here**

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of **Groundlessness for a "representationalist" inference of objects external to awareness that causes its diverse contents**

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

All sentient and insentient beings are (=) essentially one awareness

G_{36} : Category one of **All sentient and insentient beings**

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of **one awareness**

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Vimarsha and its cognates have the significance of apprehension or judgment with a recognitive structure, and may be glossed as (=) "recognitive apprehension."
The recognitive is the act of recognizing or an awareness that something perceived (e) has been perceived before

G_{40} : Category one of significance of apprehension or judgment with a recognitive structure, and may be glossed as (=) "recognitive apprehension." ; recognitive; perceived before

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **Vimarsha a**nd its cognates; act of recognizing or an awareness that something perceived (e) has been perceived before ; recognitive is the act of recognizing or an awareness that something perceived

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Multiple modules care while reading dovetailing mathematical appraisal

Vimarsha and its cognates have the significance of apprehension or judgment with a recognitive structure, and may be glossed as (=) "recognitive apprehension."

Utpaladeva (U Theories) and Abhinavagupta (A Theories) refute the Buddhist contention that recognition is a contingent reaction to direct experience by (e) claiming that it is (=) integral or transcendental to all experience

G_{44} : Category one of **Vimarsha a**nd its cognates have the significance of apprehension or judgment with a recognitive structure, and may be glossed; Buddhist contention that recognition is (=) a contingent reaction to direct experience by (e) claiming that it is (=) integral or transcendental to all experience; recognition; claiming that it is (=) integral or transcendental to all experience

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of recognitive apprehension." ; Utpaladeva (U Theories) and Abhinavagupta (A Theories) ; contingent reaction to direct experience by (e) claiming that it is (=) integral or transcendental to all experience; recognition is a contingent reaction to direct experience

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**
Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \big[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\big]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \big[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\big]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \big[(b'_{32})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big)\big]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \big[(b'_{33})^{(6)} - (b''_{33})^{(6)}\big((G_{35}),t\big)\big]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \big[(b'_{34})^{(6)} - (b''_{34})^{(6)}\big((G_{35}),t\big)\big]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**
Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \big[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\big]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \big[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\big]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \big[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\big]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \big[(b'_{36})^{(7)} - (b''_{36})^{(7)}\big((G_{39}),t\big)\big]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \big[(b'_{37})^{(7)} - (b''_{37})^{(7)}\big((G_{39}),t\big)\big]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \big[(b'_{38})^{(7)} - (b''_{38})^{(7)}\big((G_{39}),t\big)\big]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**
Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \big[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\big]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \big[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\big]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \big[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\big]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \big[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}),t\big)\big]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \big[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}),t\big)\big]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \big[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}),t\big)\big]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \big[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\big]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \big[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\big]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \big[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\big]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \big[(b'_{44})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big)\big]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \big[(b'_{45})^{(9)} - (b''_{45})^{(9)}\big((G_{47}),t\big)\big]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a_{20}')^{(3)}\,+(a_{20}'')^{(3)}(T_{21},t)\,+(a_{16}'')^{(2,2,2)}(T_{17},t)\,+(a_{13}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{40}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{44}'')^{(9,9,9)}(T_{45},t) \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a_{21}')^{(3)}\,+(a_{21}'')^{(3)}(T_{21},t)\,+(a_{17}'')^{(2,2,2)}(T_{17},t)\,+(a_{14}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{41}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{45}'')^{(9,9,9)}(T_{45},t) \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a_{22}')^{(3)}\,+(a_{22}'')^{(3)}(T_{21},t)\,+(a_{18}'')^{(2,2,2)}(T_{17},t)\,+(a_{15}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)\,+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)\,+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)\,+(a_{42}'')^{(8,8,8,8)}(T_{41},t)\,+(a_{46}'')^{(9,9,9)}(T_{45},t) \end{array}\right] G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b_{20}')^{(3)}\,-(b_{20}'')^{(3)}(G_{23},t)\,-(b_{16}'')^{(2,2,2)}(G_{19},t)\,-(b_{13}'')^{(1,1,1,)}(G,t) \\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)\,-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)\,-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t)\,-(b_{40}'')^{(8,8,8,8)}(G_{43},t)\,-(b_{44}'')^{(9,9,9)}(G_{47},t) \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b_{21}')^{(3)}\,-(b_{21}'')^{(3)}(G_{23},t)\,-(b_{17}'')^{(2,2,2)}(G_{19},t)\,-(b_{14}'')^{(1,1,1,)}(G,t) \\ -(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)\,-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)\,-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7,7)}(G_{39},t)\,-(b_{41}'')^{(8,8,8,8)}(G_{43},t)\,-(b_{45}'')^{(9,9,9)}(G_{47},t) \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{34} \qquad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(KK) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(LL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(MM) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(NN)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(OO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(PP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(g) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\leq G_{15}(t)\leq\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\leq T_{15}(t)\leq \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\leq-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\leq-(\sigma_1)^{(2)}$ (293)

$-(\tau_2)^{(2)}\leq-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\leq-(\tau_1)^{(2)}$ (294)

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)},(\nu_0)^{(7)}$:-

$$\text{For } 0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq v^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)}+(C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45},t)\right) - (a_{45}'')^{(9)}(T_{45},t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(g) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION EIGHT

"Semantic Exclusion" (Apoha) Nonepistemic "Coordination" (Sārūpya) Between Languages And Momentary Perceptual Data As The Basis For Successful Reference In Communication And Behaviors.

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) Some of the considerations they adduce to support this claim are the following: that children must build upon a subtle, innate form of (e) linguistic apprehension in their learning of conventional language

(2) there must be a recognitive ordering of(e) our most basic experiences of situations and movements in order to account for (e) our ability to perform (eb) rapid behaviors

(3) Some form of subtle application of language in all experiences is (=) necessary in order to account for our ability to remember them.

(4) The two phases of argument operate together. The idealistic Prakāsha arguments make the recognition shown by (e) the vimarsha arguments to be (=) integral to all epistemic processes, constitutive (e) of them and their objects.

(5) Moreover, on the radical logic of the Kashmiri Shaiva idealism, the recognition generating all things belongs (e) to one subject.

NOTATION

Module One

Some of the considerations they adduce to support this claim are the following: that children must build upon a subtle, innate form of (e) linguistic apprehension in their learning of conventional language

G_{13} : Category one of linguistic apprehension in their learning of conventional language

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of children must build upon a subtle, innate form

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Dual module care

recognitive ordering of our most basic experiences of situations and movements in order to account for (e) our ability to perform (eb) rapid behaviors

G_{16} : Category one of basic experiences of situations and movements in order to account for (e) our ability to perform (eb) rapid behaviors; ability to perform (eb) rapid behaviors

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of recognitive ordering; recognitive ordering of our most basic experiences of situations and movements in order to account

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Some form of subtle application of language in all experiences is (=) necessary in order to account for our ability to remember them

G_{20} : Category one of subtle application of language in all experiences

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **necessary in order to account for our ability to remember them**

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be (=) integral to all epistemic processes, constitutive (e) of them and their objects.

G_{24} : Category one of vimarsha arguments to be (=) integral to all epistemic processes, constitutive (e) of them and their objects.

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Idealistic Prakāsha arguments make the recognition shown

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be integral to all epistemic processes, constitutive (e) of them and their objects.

G_{28} : Category one of Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of all epistemic processes, constitutive (e) of them and their objects

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be integral to all epistemic processes, constitutive (e) of them and their objects

G_{32} : Category one of them and their objects

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be integral to all epistemic processes

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Moreover, on the radical logic of the Kashmiri Shaiva idealism, the recognition generating all things belongs (e) to one subject

G_{36} : Category one of one subject

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of radical logic of the Kashmiri Shaiva idealism, the recognition generating all things

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

As it is through the monistic subject's self-recognition that all phenomena are created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation through (e) Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

G_{40} : Category one of through the monistic subject's self-recognition that all phenomena

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation through (e) Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Through the monistic subject's self-recognition that all phenomena are created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's (pure consciousness) emanation through (e) Shakti (phenomenal manifested universe: cosmic general ledger) in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

G_{44} : Category one of Shakti (phenomenal manifested universe: cosmic general ledger) in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Through the monistic subject's self-recognition that all phenomena are created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's (pure consciousness) emanation

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth
augmentation coefficient for 1,2,3
$\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth
augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$
Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(QQ) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(RR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(SS) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(TT)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(UU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(VV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(h) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\hat{A}_{36})^{(7)}+(\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\hat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^{0}e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}\left((R_1)^{(9)}-(b'_{46})^{(9)}\right)}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}\left((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)}\right)}\left[e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^{0}}{G_{14}^{0}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)},(\nu_0)^{(7)}$:-

$$\text{For } 0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq v^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(h) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION EIGHT A

All Phenomena Are Created Through The Monistic Subject's Self-Recognition

INTRODUCTION—VARIABLES USED

(1) As it is through the monistic subject's self-recognition that all phenomena are (=) created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation through (e) Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

(2) The student, by coming to see this self-recognition as (=) the inner reality of all that is experienced, is led (eb) to full participation in it.

(3) Also noteworthy is the Kashmiri Shaiva theory of what may be called "semantic exclusion" (apoha). This concept had originally been formulated by the Buddhist logicians to explain (eb) a nonepistemic "coordination" (sārūpya) **between languages and (e&eb) momentary perceptual data** as the basis (e) for successful reference in communication and behaviors.

NOTATION

Module One

As it is through the monistic subject's self-recognition that all phenomena are created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation through (e) Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

G_{13} : Category one of all phenomena are created; monistic subject's self-recognition

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of monistic subject's self-recognition ;all phenomena are created

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

As it is through the monistic subject's self-recognition that all phenomena are created, the Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation through Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

G_{16} : Category one of Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation; Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Shakti in terms of self-recognition. **Note that the dependence of zero and the phenomenal universe or zero and transactional universe are reciprocal like in a bank.**;Pratyabhijnā thinkers have ostensibly demonstrated their cosmogonic myth of Shiva's emanation

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

G_{20} : Category one of zero

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of all transactions in individual, collective and cosmic general ledgers

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Student, by coming to see this self-recognition as the inner reality of all that is experienced, is led (eb) to full participation in it

G_{24} : Category one of self-recognition

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of inner reality of all that is experienced, is led (eb) to full participation in it

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

self-recognition as the inner reality of all that is experienced, is led (eb) to full participation in it

G_{28} : Category one of self-recognition as the inner reality of all that is experienced

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of full participation in it

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be integral to all epistemic processes, constitutive (e) of them and their objects.

G_{32} : Category one of Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of integral to all epistemic processes, constitutive (e) of them and their objects.

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be integral to all epistemic processes, constitutive (e) of them and their objects.

G_{36} : Category one of them and their objects

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Idealistic Prakāsha arguments make the recognition shown by the vimarsha arguments to be integral to all epistemic processes

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Kashmiri Shaiva theory (theory of vibration) of what may be called "semantic exclusion" (apoha). This concept had originally been formulated by the Buddhist logicians to explain (eb) a nonepistemic "coordination" (sārūpya) **between languages and (e&eb) momentary perceptual data** as the basis (e) for successful reference in communication and behaviors

G_{40} : Category one of Kashmiri Shaiva theory (theory of vibration)

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of "semantic exclusion" (apoha).

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Multiple modules care

concept had originally been formulated by the Buddhist logicians to explain a nonepistemic "coordination" (sārūpya) **between languages and momentary perceptual data** as the basis (e) for successful reference in communication and behaviors

G_{44} : Category one of concept had originally been formulated by the Buddhist logicians; concept had originally been formulated by the Buddhist logicians to explain a nonepistemic "coordination" (sārūpya) **between languages; momentary perceptual data** as the basis (e) for successful reference in communication and behaviors; successful reference in communication and behaviors

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of nonepistemic "coordination" (sārūpya) **between languages and (e&eb) momentary perceptual data** as the basis (e) for successful reference in communication and behaviors; **momentary perceptual data** as the basis (e) for successful reference in communication and behaviors; concept had originally been formulated by the Buddhist logicians to explain a nonepistemic "coordination" (sārūpya) **between languages;** concept had originally been formulated by the Buddhist logicians to explain a nonepistemic "coordination" (sārūpya) **between languages and momentary perceptual data**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\,+(a''_{20})^{(3)}(T_{21},t)\,+(a''_{16})^{(2,2,2)}(T_{17},t)\,+(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\,+(a''_{40})^{(8,8,8,8)}(T_{41},t)\,+(a''_{44})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\,+(a''_{21})^{(3)}(T_{21},t)\,+(a''_{17})^{(2,2,2)}(T_{17},t)\,+(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\,+(a''_{41})^{(8,8,8,8)}(T_{41},t)\,+(a''_{45})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\,+(a''_{22})^{(3)}(T_{21},t)\,+(a''_{18})^{(2,2,2)}(T_{17},t)\,+(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\,+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\,+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\,+(a''_{42})^{(8,8,8,8)}(T_{41},t)\,+(a''_{46})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b'_{20})^{(3)}\,-(b''_{20})^{(3)}(G_{23},t)\,-(b''_{16})^{(2,2,2)}(G_{19},t)\,-(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\,-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\,-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t)\,-(b''_{40})^{(8,8,8,8)}(G_{43},t)\,-(b''_{44})^{(9,9,9)}(G_{47},t) \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b'_{21})^{(3)}\,-(b''_{21})^{(3)}(G_{23},t)\,-(b''_{17})^{(2,2,2)}(G_{19},t)\,-(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\,-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\,-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t)\,-(b''_{41})^{(8,8,8,8)}(G_{43},t)\,-(b''_{45})^{(9,9,9)}(G_{47},t) \end{array}\right] T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b_{28}'')^{(5)}(G_{31},t)}\ ,\boxed{-(b_{29}'')^{(5)}(G_{31},t)}\ ,\boxed{-(b_{30}'')^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{24}'')^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b_{25}'')^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b_{26}'')^{(4,4,)}(G_{27},t)}\ \ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b_{33}'')^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b_{34}'')^{(6,6,6)}(G_{35},t)}\ \ are\ \ third\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b_{14}'')^{(1,1,1,1,1)}(G,t)}\ \ ,\boxed{-(b_{15}'')^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$
$$-\left[\begin{matrix}(a_{32}')^{(6)}\boxed{+(a_{32}'')^{(6)}(T_{33},t)}\ \boxed{+(a_{28}'')^{(5,5,5)}(T_{29},t)}\ \boxed{+(a_{24}'')^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\left[\begin{matrix}(a_{33}')^{(6)}\boxed{+(a_{33}'')^{(6)}(T_{33},t)}\ \boxed{+(a_{29}'')^{(5,5,5)}(T_{29},t)}\ \boxed{+(a_{25}'')^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\left[\begin{matrix}(a_{34}')^{(6)}\boxed{+(a_{34}'')^{(6)}(T_{33},t)}\ \boxed{+(a_{30}'')^{(5,5,5)}(T_{29},t)}\ \boxed{+(a_{26}'')^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \qquad 87$$

$\boxed{+(a_{32}'')^{(6)}(T_{33},t)}\ ,\boxed{+(a_{33}'')^{(6)}(T_{33},t)}\ ,\boxed{+(a_{34}'')^{(6)}(T_{33},t)}\ \ are\ \ first\ \ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{28}'')^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a_{29}'')^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a_{30}'')^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{24}'')^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a_{25}'')^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a_{26}'')^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(WW) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(XX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(YY) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(ZZ)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(AAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(BBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$

Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\,(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+\ (\hat{P}_{40})^{(8)}\,(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+\ (\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}$, $T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(i) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right| \; e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(i) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION NINE

Pratyabhijnā Ontology: The Syntax Of Empowered Identity

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) According to the Buddhists, words have (e) no isomorphism with (e&eb) the sense data, but only exclude (e) other words that would not (e) lead to successful behavior.

(2) The only reference of the word "cow" to a perceived particular is (=) that it excludes non-cows, for example, a horse, a car, and so on.

NOTATION

Module One

According to the Buddhists, words have no isomorphism with (e&eb) the sense data, but only exclude (e) other words that would not (e) lead to successful behavior

G_{13} : Category one of no isomorphism with (e&eb) the sense data, but only exclude (e) other words that would not (e) lead to successful behavior

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of words

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

words have no isomorphism with the sense data, but only exclude (e) other words that would not (e) lead to

successful behavior

G_{16} : Category one of words have no isomorphism; sense data, but only exclude (e) other words that would not (e) lead to successful behavior

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of sense data, but only exclude (e) other words that would not (e) lead to successful behavior; words have no isomorphism

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

words have no isomorphism with the sense data, but only exclude other words that would not (e) lead to successful behavior

G_{20} : Category one of other words that would not (e) lead to successful behavior

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of words have no isomorphism with the sense data, but only exclude

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

words have no isomorphism with the sense data, but only exclude other words that would not (e) lead to successful behavior

G_{24} : Category one of lead to successful behavior

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of words have no isomorphism with the sense data, but only exclude other words

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

only reference of the word “cow” to a perceived particular is that it excludes non-cows, for example, a horse, a car, and so on

G_{28} : Category one of only reference of the word “cow” to a perceived particular

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of " reference to the word" cow' excludes non-cows, for example, a horse, a car, and so on

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

reference of the word "cow" to a perceived particular is that it excludes non-cows, for example, a horse, a car, and so on

G_{32} : Category one of perceived particular is that it excludes non-cows, for example, a horse, a car, and so on

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of reference of the word "cow"

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is not formulated like the latter (that is, on the basis of considerations about the systematicity of entire languages).

G_{36} : Category one of Buddhist theory (B theories) has an interesting point of agreement

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is not formulated like the latter (that is, on the basis of considerations about the systematicity of entire languages).

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is not

formulated like the latter that is, on the basis (e) of considerations about the systematicity of entire languages

G_{40} : Category one of considerations about the systematicity of entire languages

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is not formulated like the latter

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is not formulated like the latter that is, on the basis (e) of considerations about the systematicity of entire languages

G_{44} : Category one of Buddhist theory (B theories); structuralist and poststructuralist conceptions of the determination of linguistic value by difference

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of structuralist and poststructuralist conceptions of the determination of linguistic value by difference ;Buddhist theory (B theories)

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}&+(a''_{16})^{(2)}(T_{17},t)&+(a''_{13})^{(1,1,)}(T_{14},t)&+(a''_{20})^{(3,3,3)}(T_{21},t)\\+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{36})^{(7,7,7)}(T_{37},t)&+(a''_{40})^{(8,8,8)}(T_{41},t)&+(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)}&+(a''_{17})^{(2)}(T_{17},t)&+(a''_{14})^{(1,1,)}(T_{14},t)&+(a''_{21})^{(3,3,3)}(T_{21},t)\\+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{37})^{(7,7,7)}(T_{37},t)&+(a''_{41})^{(8,8,8)}(T_{41},t)&+(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)}&+(a''_{18})^{(2)}(T_{17},t)&+(a''_{15})^{(1,1,)}(T_{14},t)&+(a''_{22})^{(3,3,3)}(T_{21},t)\\+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{38})^{(7,7,7)}(T_{37},t)&+(a''_{42})^{(8,8,8)}(T_{41},t)&+(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)}&-(b''_{16})^{(2)}(G_{19},t)&-(b''_{13})^{(1,1,)}(G,t)&-(b''_{20})^{(3,3,3,)}(G_{23},t)\\-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)&-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)&-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\-(b''_{36})^{(7,7,7)}(G_{39},t)&-(b''_{40})^{(8,8,8)}(G_{43},t)&-(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt}=(b_{17})^{(2)}T_{16}-\begin{bmatrix}(b'_{17})^{(2)}&-(b''_{17})^{(2)}(G_{19},t)&-(b''_{14})^{(1,1,)}(G,t)&-(b''_{21})^{(3,3,3,)}(G_{23},t)\\-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)&-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)&-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)\\-(b''_{37})^{(7,7,7)}(G_{39},t)&-(b''_{41})^{(8,8,8)}(G_{43},t)&-(b''_{45})^{(9,9)}(G_{47},t)\end{bmatrix}T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(CCC) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(DDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(EEE) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(FFF)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(GGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(HHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(j) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(j) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION NINE A

Exclusion Itself Depends Upon A Comparative Synthesis, Or Recognition, Of What Does And Does Not Fit Within Particular Categories

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) Buddhist theory has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by (e) difference, although it is (=) not formulated like the latter that is, on the basis of considerations about the systematicity of (e) entire languages
(2) Utpaladeva (U theories) and Abhinavagupta (A theories) argue that exclusion itself depends upon (=) a comparative synthesis, or recognition, of what does and does not (e&eb) fit within particular categories.
(3) We recognize that the cow is (=) not a non-cow such as a horse.
(4) The Pratyabhijnā theorists thus in effect explain difference itself as (=) a kind of similarity.
(5) Difference is identified (e&eb) in various circumstances like other forms of similarity.
(6) According to the Shaivas such difference-identification is (=) one of the principal expressions of Shiva's emanating self-recognition.
(7) Just as Utpaladeva and Abhinavagupta appropriate Bhartrihari in equating self-recognition with (=) Supreme Speech and thereby interpreting recognitive apprehension as (=) linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā).

NOTATION

Module One

Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is (=) not formulated like the latter that is, on the basis of considerations about the systematicity of (e) entire languages

G_{13} : Category one of difference, although it is (=) not formulated like the latter that is, on the basis of considerations about the systematicity of (e) entire languages

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is (=) not formulated like the latter that is, on the basis of considerations about the systematicity of (e) entire languages

G_{16} : Category one of structuralist; poststructuralist conceptions of the determination of linguistic value by difference

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of poststructuralist conceptions of the determination of linguistic value by difference ;structuralist

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is (=) not formulated like the latter that is, on the basis of considerations about the systematicity of (e) entire languages

G_{20} : Category one of entire languages

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Buddhist theory (B theories) has an interesting point of agreement with contemporary structuralist and poststructuralist conceptions of the determination of linguistic value by difference, although it is (=) not formulated like the latter that is, on the basis of considerations about the systematicity

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Utpaladeva (U theories) and Abhinavagupta (A theories) argue that exclusion itself depends upon a comparative synthesis, or recognition, of what does and does not (e&eb) fit within particular categories.

G_{24} : Category one of Utpaladeva (U theories) and Abhinavagupta (A theories) argue that exclusion itself depends upon

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of comparative synthesis, or recognition, of what does and does not (e&eb) fit within particular categories

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Utpaladeva (U theories) and Abhinavagupta (A theories) argue that exclusion itself depends upon a comparative synthesis, or recognition, of what does and does not (e&eb) fit within particular categories

G_{28} : Category one of Utpaladeva (U theories) and Abhinavagupta (A theories) argue that exclusion itself depends upon a comparative synthesis, or recognition; fit within particular categories

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of fit within particular categories ;Utpaladeva (U theories) and Abhinavagupta (A theories) argue that exclusion itself depends upon a comparative synthesis, or recognition

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

cow is (=) not a non-cow such as a horse

G_{32} : Category one of cow

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of not a non-cow such as a horse

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Pratyabhijnā theorists thus in effect explain difference itself as (=) a kind of similarity

G_{36} : Category one of Pratyabhijnā theorists thus in effect explain difference

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of kind of similarity

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Difference is identified (e&eb) in various circumstances like other forms of similarity

G_{40} : Category one of Difference; various circumstances like other forms of similarity

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of various circumstances like other forms of similarity; Difference

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

According to the Shaivas such difference-identification is (=) one of the principal expressions of Shiva's (pure consciousness) emanating self-recognition

G_{44} : Category one of difference-identification

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of one of the principal expressions of Shiva's (pure consciousness) emanating self-recognition

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\left[+(a''_{20})^{(3)}(T_{21},t)\right]\left[+(a''_{16})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{13})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{40})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{44})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\left[+(a''_{21})^{(3)}(T_{21},t)\right]\left[+(a''_{17})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{14})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{37})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{41})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{45})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\left[+(a''_{22})^{(3)}(T_{21},t)\right]\left[+(a''_{18})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{15})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{38})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{42})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{46})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & \boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & \boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & \boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a_{40}'')^{(8,8\,,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \begin{bmatrix} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \begin{bmatrix} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, \ (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(III) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(JJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(KKK) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(LLL)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(MMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(NNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(k) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},\ \boldsymbol{if}\ (v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(k) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TEN

Shiva's Mythic Action Through Shakti As Self-Recognition That Constitutes (Eb) All Experience And Objects Of Experience

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) The Grammarian view itself originated in Brahmanic interpretations of the Veda as (=) expressing injunctions for sacrifice.

(2) The Kashmiri Shaivas further agree (=) with much of **Vedic exegetics in conceiving being as both narrative and recapitulatory ritual action.** .

NOTATION

Module One

Just as Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in equating self-recognition with Supreme Speech and thereby interpreting recognitive apprehension as (=) linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā).

G_{13} : Category one of self-recognition

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Supreme Speech

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Just as Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in equating self-recognition with Supreme Speech and thereby interpreting recognitive apprehension as (=) linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā).

G_{16} : Category one of Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in equating self-recognition with Supreme Speech

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of recognitive apprehension as (=) linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā).

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Just as Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in equating self-recognition with Supreme Speech and thereby interpreting recognitive apprehension as linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā

G_{20} : Category one of Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in

equating self-recognition with Supreme Speech and thereby interpreting recognitive apprehension

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in equating self-recognition with Supreme Speech and thereby interpreting recognitive apprehension as linguistic in nature, they also follow the Grammarian school in interpreting being or existence (sattā) (the generic referent of language) as (=) action (kriyā)

G_{24} : Category one of Utpaladeva (U theories) and Abhinavagupta (A theories) appropriate Bhartrihari in equating self-recognition with Supreme Speech and thereby interpreting recognitive apprehension as linguistic in nature, they also follow the Grammarian school in interpreting **being or existence (sattā) (the generic referent of language)**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of action (kriyā)

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

existence (sattā) is the generic referent of language

G_{28} : Category one of existence (sattā)

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of generic referent of language

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Grammarian view itself originated in Brahmanic interpretations of the Veda as (=) expressing injunctions for

sacrifice

G_{32} : Category one of Brahmanic interpretations of the Veda as (=) expressing injunctions for sacrifice

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Grammarian view

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Grammarian view itself originated in Brahmanic interpretations of the Veda as (=) expressing injunctions for sacrifice

G_{36} : Category one of Grammarian view itself originated in Brahmanic interpretations of the Veda

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of expressing injunctions for sacrifice

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Kashmiri Shaivas further agree (=) with much of **Vedic exegetics in conceiving being as both narrative and recapitulatory ritual action.**

G_{40} : Category one of **Vedic exegetics conceive being**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **both narrative and recapitulatory ritual action.**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Following the account above, it is **Shiva's (Pure consciousness wherein all dualities like space =time, being =no being, chaos =order, initial conditions for the creation of the universe mythic action through Shakti as (=) self-recognition that constitutes (eb) all experience and objects of experience, and that is reenacted by philosophical discourse**.

G_{44} : Category one of **Shiva's (Pure consciousness wherein all dualities like space =time, being =no being, chaos =order, initial conditions for the creation of the universe mythic action through Shakti**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **self-recognition that constitutes (eb) all experience and objects of experience, and that is reenacted by philosophical discourse**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18}$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) & \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) & \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) & \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition

coefficients for category 1, 2, and 3
$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition
coefficients for category 1, 2, and 3
$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2, and 3
$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition
coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for
category 1, 2 and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation
coefficient for category 1, 2 and 3
$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation
coefficient for category 1, 2 and 3
$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh
augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$
are eighth augmentation coefficient for 1,2,3
$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} & \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} & \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} & \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \quad \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(OOO) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(PPP) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(QQQ) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(RRR)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(SSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(TTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{40})^{(8)},\ T_i^0 \le (\hat{Q}_{40})^{(8)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(l) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(l) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^{*}=\frac{(b_{28})^{(5)}T_{29}^{*}}{[(b_{28}')^{(5)}-(b_{28}'')^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*}=\frac{(b_{30})^{(5)}T_{29}^{*}}{[(b_{30}')^{(5)}-(b_{30}'')^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*})=0$, T_{33}^{*} given by $f(T_{33}^{*})=0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a_{32}')^{(6)}+(a_{32}'')^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a_{34}')^{(6)}+(a_{34}'')^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b_{32}')^{(6)}-(b_{32}'')^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b_{34}')^{(6)}-(b_{34}'')^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a_{36}')^{(7)}+(a_{36}'')^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a_{38}')^{(7)}+(a_{38}'')^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b_{36}')^{(7)}-(b_{36}'')^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b_{38}')^{(7)}-(b_{38}'')^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a_{40}')^{(8)}+(a_{40}'')^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a_{42}')^{(8)}+(a_{42}'')^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b_{40}')^{(8)}-(b_{40}'')^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b_{42}')^{(8)}-(b_{42}'')^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a_{44}')^{(9)}+(a_{44}'')^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a_{46}')^{(9)}+(a_{46}'')^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b_{44}')^{(9)}-(b_{44}'')^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b_{46}')^{(9)}-(b_{46}'')^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TEN A

Shiva's Action Through Shakti As Self-Recognition That Constitutes All Experience And Objects Of Experience

INTRODUCTION—VARIABLES USED

(1) Following the account above, it is **Shiva's action through Shakti as self-recognition that constitutes (eb) all experience and objects of experience, and that is reenacted by philosophical discourse**.

(2) The Pratyabhijnā thinkers propound their philosophy of Shiva's action to explain (eb) a wide range of topics of ontology.

(3) One of their concerns is to describe how Shiva's action generates (eb) a multiplicity of relationships (sambandha) or universals (sāmānya) as (=) the referents of discrete instances of recognitive apprehension.

(4) With this theory they attempt to subvert (e) the Buddhist logicians' contention that evanescent particulars are (=) ontologically fundamental.

(5) For the Shaivas, **categories are (=) primitive, and particulars are formed out (e) of syntheses of those categories. This is what is being attempted to The Theory of everything by classification and equational dovetail**

(6) Most illustrative of the Pratyabhijnā thinkers' "mythico-ritual approach" to ontology is their use (e) of theories of Sanskrit syntax to explain Shiva's action.

(7) Again reflecting the Vedic roots of South Asian philosophies, many schools of Hinduism and Buddhism—even those which do not (e) view all existence as action—frequently advert to considerations of action syntax in treating (e&eb) ontological or metaphysical topics

NOTATION

Module One

Following the account above, it is **Shiva's action through Shakti as self-recognition that constitutes (eb) all experience and objects of experience, and that is reenacted by philosophical discourse**

G_{13} : Category one of **all experience and objects of experience**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **Shiva's action through Shakti as self-recognition that constitutes**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Shiva's action through Shakti as self-recognition

G_{16} : Category one of **Shiva's action through Shakti**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of **self-recognition**

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Pratyabhijnā thinkers propound their philosophy of Shiva's action to explain (eb) a wide range of topics of ontology

G_{20} : Category one of Pratyabhijnā thinkers propound their philosophy of Shiva's action

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of wide range of topics of ontology

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

One of their concerns is to describe how Shiva's action generates a multiplicity of relationships (sambandha) or universals (sāmānya) as (=) the referents of discrete instances of recognitive apprehension

G_{24} : Category one of Shiva's action **(read pure consciousness please: I will not repeat; it is here all dualities cease see earlier models)**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of multiplicity of relationships (sambandha) or universals (sāmānya) as (=) the referents of discrete instances of recognitive apprehension

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Shiva's action generates a multiplicity of relationships (sambandha) or universals (sāmānya) as (=) the referents of discrete instances of recognitive apprehension

G_{28} : Category one of Shiva's action generates a multiplicity of relationships (sambandha) or universals (sāmānya)

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of **referents of discrete instances of recognitive apprehension**

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Categories are (=) primitive, and particulars are formed out (e) of syntheses of those categories. This is what is being attempted to The Theory of everything by classification and equational dovetail

G_{32} : Category one of **Categories**

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of **primitive, and particulars are formed out (e) of syntheses of those categories. This is what is being attempted to The Theory of everything by classification and equational dovetail**

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

One of their concerns is to describe how Shiva's action generates a multiplicity of relationships (sambandha) or universals (sāmānya) as (=) the referents of discrete instances of recognitive apprehension.

G_{36} : Category one of Shiva's action generates a multiplicity of relationships (sambandha) or universals (sāmānya)

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of referents of discrete instances of recognitive apprehension

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Most illustrative of the Pratyabhijnā thinkers' "mythico-ritual approach" to ontology is their use of theories of Sanskrit syntax to explain Shiva's action

G_{40} : Category one of "mythico-ritual approach" to ontology

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of use of theories of Sanskrit syntax to explain Shiva's action

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Again reflecting the Vedic roots of South Asian philosophies, many schools of Hinduism and Buddhism—

even those which do not (e) view all existence as action—frequently advert to considerations of action syntax in treating (e&eb) ontological or metaphysical topics

G_{44} : Category one of view all existence as action—frequently advert to considerations of action syntax in treating (e&eb) ontological or metaphysical topics

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of reflecting the Vedic roots of South Asian philosophies, many schools of Hinduism and Buddhism—

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \qquad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**
Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**
Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**
Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**
Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}+(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}+(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}+(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\,-(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\,-(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\,-(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(UUU) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(VVV) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(WWW) $\lim_{T_2 \to \infty} \ (a_i'')^{(7)}\,(T_{37}, t) = (p_i)^{(7)}$

(XXX)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(YYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(ZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$

$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$

$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(m) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(m) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION ELEVEN

"Being Is The Agency Of The Act Of Becoming, Which Is, The Freedom Characteristic Of An Agent Regarding All Actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, And 1:258-59)."

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) The relevant considerations pertain to how verbs articulating action relate to (e&eb) declined nouns indicating the concomitants of action (kārakas)—in English, roughly, the agent, object, instrument, purpose, source and location.

(2) Now, most Sanskritic philosophies, Hindu as well as Buddhist, have tended to delimit (e) the syntactic role of the agent **(kartri kāraka)—**to different degrees, but sometimes quite strongly.

(3) The explicit and implicit reasons for this tendency are complex. At one level it evidently reflects the orthodox Brahmanic norms that subordinate the individual's agency to the order of objective ritual behavior—pertaining to sacrifice, caste, life cycle, and so on. It also seems more broadly to reflect both Hindu and Buddhist concepts of the agent's bondage to (e) the process of action and result (karma) extending across rebirths (see Gerow 1982).

(4) The mainstream Buddhist philosophies completely deny (e) the existence of a self in the dependent origination **(pratityasamutpada**) of karma.

(5) Developing suggestions of Somānanda, the Pratyabhijnā philosophers expound a distinctive theory of agency to **rationalize their tantric mythic and ritual drama of omnipotence**. In their theory they take up several earlier understandings of the positive albeit delimited role of the agent and radicalize them. According to the Kashmiri Shaivas, all causal processes and other relationships constituting (eb) the universe are synthesized and impelled by (e&eb) the mythic agency of Shiva in his act of self-recognition.

(6) Shiva's agency encompasses (e&eb) the actions of sentient beings as well as the motions and transformations of insentient beings. See earlier models of projection, axes, dimensions, rotations, foldings, concurrences, shocks, encounters, etc.

(7) The Kashmiri Shaivas ultimately reduce (e) the entire action of existence to agency.

(8) As Abhinavagupta (A Theories) explains**, "Being is (=) the agency of the act of becoming, which is, the freedom characteristic of (e) an agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)."**

(9) Again, this theory of omnificent syntactic agency is ritually axiomatic as well as mythical. Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy, in a manner homologous to (=) the epistemology of recognition, as leading to (eb) salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design"

NOTATION

Module One

relevant considerations pertain to how verbs articulating action relate to (e&eb) declined nouns indicating the concomitants of action (kārakas)—in English, roughly, the agent, object, instrument, purpose, source and location

G_{13} : Category one of relevant considerations pertain to how verbs articulating action; declined nouns indicating the concomitants of action (kārakas)—in English, roughly, the agent, object, instrument, purpose, source and location

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of declined nouns indicating the concomitants of action (kārakas)—in English, roughly, the agent, object, instrument, purpose, source and location ;relevant considerations pertain to how verbs

articulating action

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

most Sanskritic philosophies, Hindu as well as Buddhist, have tended to delimit (e) the syntactic role of the agent **(kartri kāraka)**—to different degrees, but sometimes quite strongly

G_{16} : Category one of syntactic role of the agent **(kartri kāraka)**—to different degrees, but sometimes quite strongly

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of most Sanskritic philosophies, Hindu as well as Buddhist thought

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Explicit and implicit reasons for this tendency are complex. At one level it evidently reflects the orthodox Brahmanic norms that subordinate the individual's agency to the order of objective ritual behavior—pertaining to sacrifice, caste, life cycle, and so on. It also seems more broadly to reflect both Hindu and Buddhist concepts of the agent's bondage to (e) the process of action and result (karma) extending across rebirths (see Gerow 1982).

G_{20} : Category one of process of action and result (karma) extending across rebirths (see Gerow 1982).

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Hindu and Buddhist concepts of the agent's bondage

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Mainstream Buddhist philosophies completely deny (e) the existence of a self in the dependent origination **(pratityasamutpada)** of karma

G_{24} : Category one of existence of a self in the dependent origination **(pratityasamutpada)** of karma

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Mainstream Buddhist philosophies

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Developing suggestions of Somānanda, the Pratyabhijnā philosophers expound a distinctive theory of agency to **rationalize their tantric mythic and ritual drama of omnipotence**. In their theory they take up several earlier understandings of the positive albeit delimited role of the agent and radicalize them. According to the Kashmiri Shaivas, all causal processes and other relationships constituting (e) the universe are synthesized and impelled by (e&eb) the mythic agency of Shiva in his act of self-recognition. **Self recognition is contemplation of one's nature, self, and this leads to multifarious differentiation of objects**

G_{28} : Category one of universe is synthesized and impelled by (e&eb) the mythic agency of Shiva in his act of self-recognition. Self recognition is contemplation of one's nature, self, and this leads to multifarious differentiation of objects

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of causal processes and other relationships

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Universe is synthesized and impelled by (e&eb) the mythic agency of Shiva in his act of self-recognition.

G_{32} : Category one of Shiva in his act of self-recognition

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Universe is synthesized and impelled

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Shiva's (pure consciousness) agency encompasses (e&eb) the actions of sentient beings as well as the motions and transformations of insentient beings. See earlier models of projection, axes, dimensions, rotations, foldings, concurrences, shocks, encounters, etc.

G_{36} : Category one of Shiva's (pure consciousness) agencies; actions of sentient beings as well as the motions and transformations of insentient beings. See earlier models of projection, axes, dimensions, rotations, foldings, concurrences, shocks, encounters, etc.

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of actions of sentient beings as well as the motions and transformations of insentient beings. See earlier models of projection, axes, dimensions, rotations, foldings, concurrences, shocks, encounters, etc.; Shiva's (pure consciousness) agency

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Kashmiri Shaivas ultimately reduce the entire action of existence to agency.

G_{40} : Category one of entire action of existence

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of agency of pure consciousness

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

As Abhinavagupta (A Theories) explains**, "Being is (=) the agency of the act of becoming, which is, the freedom characteristic of (e) an agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)."**

G_{44} : Category one of **Being**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **agency of the act of becoming, which is, the freedom characteristic of (e) an agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} +(a_{16}'')^{(2)}(T_{17},t) +(a_{13}'')^{(1,1,)}(T_{14},t) +(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t) +(a_{40}'')^{(8,8,8)}(T_{41},t) +(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} +(a_{17}'')^{(2)}(T_{17},t) +(a_{14}'')^{(1,1,)}(T_{14},t) +(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t) +(a_{41}'')^{(8,8,8)}(T_{41},t) +(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} +(a_{18}'')^{(2)}(T_{17},t) +(a_{15}'')^{(1,1,)}(T_{14},t) +(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t) +(a_{42}'')^{(8,8,8)}(T_{41},t) +(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} \ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(AAAA) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(BBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(CCCC) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(DDDD)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(EEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(FFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$$ 138

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$$ 139

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$$ 140

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$$ 142

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(n) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case,

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)}+(C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

424

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(n) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\,(G_{47})^*\,) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWELVE

INTRODUCTION—VARIABLES USED

Tantra and Kashmiri Shaivism (Wikipedia)

(1) Abhinavagupta (A Theories) likewise, in his explanation of the preliminary ceremonies of the tantric ritual, identifies (e&eb) various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases**.

(2) He explains that the aspirant's goal in the ritual action is (=) identification with Shiva as agent of all the cases.

(3) As Abhinavagupta (A Theories) explains**, "Being is (=) the agency of the act of becoming, which is, the freedom characteristic of (e) an agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)."**

(4) Again, this theory of omnificent syntactic agency is ritually axiomatic as well as mythical. Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy, in a manner homologous to (=) the epistemology of recognition, as leading to (eb) salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design"

NOTATION

Module One

Abhinavagupta (A Theories) likewise, in his explanation of the preliminary ceremonies of the tantric ritual, identifies (e&eb) various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases**

G_{13} : Category one of Abhinavagupta (A Theories)

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of preliminary ceremonies of the tantric ritual, identifies (e&eb) various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the**

Sanskrit grammatical cases

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Abhinavagupta (A Theories) likewise, in his explanation of the preliminary ceremonies of the tantric ritual, identifies (e&eb) various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases**

G_{16} : Category one of Abhinavagupta (A Theories) likewise, in his explanation of the preliminary ceremonies of the tantric ritual; various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases;** Abhinavagupta (A Theories) likewise, in his explanation of the preliminary ceremonies of the tantric ritual

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations—**with the Sanskrit grammatical cases**

G_{20} : Category one of various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations; **Sanskrit grammatical cases**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **Sanskrit grammatical cases** ;various components of the ritual—such as the location, ritual implements and object of sacrifice, flowers, and oblations

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

He explains that the aspirant's goal in the ritual action is (=) identification with Shiva as agent of all the cases.

G_{24} : Category one of aspirant's goal in the ritual action

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of identification with Shiva as agent of all the cases.

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Such identification is through Shakti the manifestation of phenomenal world and is a constitution of strings with resonance. It is the onus on aspirant to transcend the light and sound sight and speech is the basic principle for the achievement of Satchidananda

G_{28} : Category one of identification

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of through Shakti the manifestation of phenomenal world and is a constitution of strings with resonance. It is the onus on aspirant to transcend the light and sound sight and speech is the basic principle for the achievement of Satchidananda

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Such identification is through Shakti the manifestation of phenomenal world and is (=) a constitution of strings with resonance. It is the onus on aspirant to transcend the light and sound sight and speech is the basic principle for the achievement of Satchidananda

G_{32} : Category one of identification is through Shakti the manifestation of phenomenal world

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of constitution of strings with resonance. It is the onus on aspirant to transcend the light and sound sight and speech is the basic principle for the achievement of Satchidananda

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

onus is on aspirant to transcend the light and sound sight and speech is the basic principle for the achievement of Satchidananda

G_{36} : Category one of light and sound sight and speech for the achievement of Satchidananda

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of onus is on aspirant

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

being is the agency of the act of becoming, which is, the freedom characteristic of (e) an agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)

G_{40} : Category one of **agent regarding all actions (Īshvarapratyabhijnāvimarshinī, 1.5.14, and 1:258-59)**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **being is the agency of the act of becoming, which is, the freedom characteristic**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Again, this theory of omnificent syntactic agency is ritually axiomatic as well as mythical. Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy, in a manner homologous to the epistemology of recognition, as leading to (eb) salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design"

G_{44} : Category one of Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of epistemology of recognition, as leading to (eb) salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design"

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} \; +(a''_{16})^{(2)}(T_{17},t) \; +(a''_{13})^{(1,1,)}(T_{14},t) \; +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) \; +(a''_{40})^{(8,8,8)}(T_{41},t) \; +(a''_{44})^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} \; +(a''_{17})^{(2)}(T_{17},t) \; +(a''_{14})^{(1,1,)}(T_{14},t) \; +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) \; +(a''_{41})^{(8,8,8)}(T_{41},t) \; +(a''_{45})^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} \; +(a''_{18})^{(2)}(T_{17},t) \; +(a''_{15})^{(1,1,)}(T_{14},t) \; +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) \; +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) \; +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) \; +(a''_{42})^{(8,8,8)}(T_{41},t) \; +(a''_{46})^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b_{13}'')^{(1,1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$ $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a_{24}')^{(4)} & +(a_{24}'')^{(4)}(T_{25},t) & +(a_{28}'')^{(5,5,)}(T_{29},t) & +(a_{32}'')^{(6,6,)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3)}(T_{21},t) \\ & +(a_{36}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a_{25}')^{(4)} & +(a_{25}'')^{(4)}(T_{25},t) & +(a_{29}'')^{(5,5,)}(T_{29},t) & +(a_{33}'')^{(6,6)}(T_{33},t) \\ & +(a_{14}'')^{(1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3)}(T_{21},t) \\ & +(a_{37}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a_{26}')^{(4)} & +(a_{26}'')^{(4)}(T_{25},t) & +(a_{30}'')^{(5,5,)}(T_{29},t) & +(a_{34}'')^{(6,6,)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3)}(T_{21},t) \\ & +(a_{38}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a_{24}'')^{(4)}(T_{25},t)$, $(a_{25}'')^{(4)}(T_{25},t)$, $(a_{26}'')^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a_{28}'')^{(5,5,)}(T_{29},t)$, $+(a_{29}'')^{(5,5,)}(T_{29},t)$, $+(a_{30}'')^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a_{32}'')^{(6,6,)}(T_{33},t)$, $+(a_{33}'')^{(6,6,)}(T_{33},t)$, $+(a_{34}'')^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a_{13}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a_{16}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a_{20}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)} (T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty} (b_i'')^{(6)} \left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(GGGG) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(HHHH)The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(IIII) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(JJJJ)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(KKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(LLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$$ 138

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$$ 139

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$$ 140

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$$ 142

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(o) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}\ , G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right| \; e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)},, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

$$\text{For } 0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$$

$$v^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq v^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45},t)\right) - (a_{45}'')^{(9)}(T_{45},t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(o) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523

A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b_i'')^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTEEN

Parā, Parā-Aparā And Aparā Śakti

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

(1) The trident (triśūlābija maṇḍalam), symbol and yantra of Parama Shiva, representing the triadic energies of Parā, Parā-aparā and aparā śakti
(2) The trident (triśūlābija maṇḍalam), symbol and yantra of Parama Shiva, representing the triadic energies of Parā, Parā-aparā and aparā śakti
(3) The trident (triśūlābija maṇḍalam), symbol and yantra of Parama Shiva, representing the triadic energies of Parā, Parā-aparā and aparā śakti
(4) In Kashmir Shaivism, all things are (=) a manifestation of this Consciousness.
(5) This means that from the point of view of Kashmir Shaivism the phenomenal world (Śakti) is (=) real, and it exists and has its being in (eb) Consciousness (Chit).
(6) In comparison, Advaita Vedanta holds that Brahman is (=) inactive (niṣkriya) and the phenomenal world is (=) an illusion (māyā).
(7) Thus, the philosophy of Kashmir Shaivism, also called the Trika, can be seen in contrast (e&eb) to Shankara's Advaita. This is disputable. Chit Shakti is one responsible for (e) creation and maya Shakti is illusion which is responsible for (e) producing nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of status of no mind condition). This is my interpretation. There is no conflict between the two.
(8) Thus, the philosophy of Kashmir Shaivism, also called the Trika, can be seen in contrast (e&eb) to Shankara's Advaita. This is disputable. Chit Shakti is one responsible for (e) creation and maya Shakti is illusion which is responsible for (e) producing nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of status of no mind condition). This is my interpretation. There is no conflict between the two.
(9) Thus, the philosophy of Kashmir Shaivism, also called the Trika, can be seen in contrast (e&eb) to Shankara's Advaita. This is disputable. Chit Shakti is one responsible for (e) creation and maya Shakti is illusion which is responsible for (e) producing nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of status of no mind condition or

achievement if the status of I = cosmic general ledger). This is my interpretation. There is no conflict between the two.

NOTATION

Module One

Again, this theory of omnificent syntactic agency is ritually axiomatic as well as mythical. Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy, in a manner homologous to the epistemology of recognition, as leading to (eb) salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design"

G_{13} : Category one of theory of omnificent syntactic agency is ritually axiomatic as well as mythical. Utpaladeva (U Theories) describes the method of the Pratyabhijnā philosophy, in a manner homologous to the epistemology of recognition,

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of salvation through the contemplation of one's status as the agent of the universe. Whatever one does be it that of the top most man or a hang man he is doing so to serve the " Grand design"

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

trident (triśūlābija maṇḍalam), symbol and yantra of Parama Shiva, representing the triadic energies of Parā, Parā-aparā and aparā śakti

G_{16} : Category one of Parā; Parā-aparā

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Parā-aparā; Parā

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Parā-aparā and aparā śakti

G_{20} : Category one of Parā-aparā; aparā śakti

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of aparā śakti; Parā-aparā

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Parā and aparā śakti

G_{24} : Category one of Parā; aparā śakti

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of aparā śakti ;Parā

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

all things are (=) a manifestation of this Consciousness

G_{28} : Category one of all things

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of manifestation of Consciousness

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

This means that from the point of view of Kashmir Shaivism the phenomenal world (Śakti) is real, and it exists and has its being in (eb) Consciousness (Chit)

G_{32} : Category one of phenomenal world (Śakti)

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of real, and it exists and has its being in (eb) Consciousness (Chit)

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

phenomenal world (Śakti) is real, and it exists and has its being in (eb) Consciousness (Chit)

G_{36} : Category one of phenomenal world (Śakti) is real, and it exists and has its being

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Consciousness (Chit)

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Advaita Vedanta holds that Brahman is (=) inactive (niṣkriya) and the phenomenal world is (=) an illusion (māyā).

G_{40} : Category one of Advaita Vedanta holds that Brahman

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of inactive (niṣkriya)

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Phenomenal world is (=) an illusion (māyā).

G_{44} : Category one of phenomenal world

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of illusion (māyā).

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)} \ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$
$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients

for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{16}')^{(2)} & +(a_{16}'')^{(2)}(T_{17},t) & +(a_{13}'')^{(1,1,)}(T_{14},t) \quad +(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a_{17}')^{(2)} & +(a_{17}'')^{(2)}(T_{17},t) & +(a_{14}'')^{(1,1,)}(T_{14},t) \quad +(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{18}')^{(2)} & +(a_{18}'')^{(2)}(T_{17},t) & +(a_{15}'')^{(1,1,)}(T_{14},t) \quad +(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for

category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition*

coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$ $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\left[\begin{array}{ccc}(a_{24}')^{(4)}+(a_{24}'')^{(4)}(T_{25},t) & +(a_{28}'')^{(5,5,)}(T_{29},t) & +(a_{32}'')^{(6,6,)}(T_{33},t)\\ +(a_{13}'')^{(1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{36}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9)}(T_{45},t)\end{array}\right]G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\left[\begin{array}{ccc}(a_{25}')^{(4)}+(a_{25}'')^{(4)}(T_{25},t) & +(a_{29}'')^{(5,5,)}(T_{29},t) & +(a_{33}'')^{(6,6)}(T_{33},t)\\ +(a_{14}'')^{(1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{37}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9)}(T_{45},t)\end{array}\right]G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\left[\begin{array}{ccc}(a_{26}')^{(4)}+(a_{26}'')^{(4)}(T_{25},t) & +(a_{30}'')^{(5,5,)}(T_{29},t) & +(a_{34}'')^{(6,6,)}(T_{33},t)\\ +(a_{15}'')^{(1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3)}(T_{21},t)\\ +(a_{38}'')^{(7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9)}(T_{45},t)\end{array}\right]G_{26} \quad (75)$$

$(a_{24}'')^{(4)}(T_{25},t)$, $(a_{25}'')^{(4)}(T_{25},t)$, $(a_{26}'')^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a_{28}'')^{(5,5,)}(T_{29},t)$, $+(a_{29}'')^{(5,5,)}(T_{29},t)$, $+(a_{30}'')^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a_{32}'')^{(6,6,)}(T_{33},t)$, $+(a_{33}'')^{(6,6,)}(T_{33},t)$, $+(a_{34}'')^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a_{13}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{16}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{20}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* $1,2$ *and* 3

$+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* $1,2$ *and* 3

$+(a_{46}'')^{(9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} & \boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} & \boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} & \boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} \\ \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} & \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} \\ \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad 84$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5)}(G_{31},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients

for category 1,2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)]$, $[-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)]$,$[-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)]$ are seventh detrition coefficients for category 1,2, and 3

$[-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)]$,$[-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)]$,$[-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)]$ are eighth detrition coefficients for category 1,2, and 3

$[-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)]$,$[-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)]$,$[-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)]$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}[+(a''_{32})^{(6)}(T_{33},t)] & [+(a''_{28})^{(5,5,5)}(T_{29},t)] & [+(a''_{24})^{(4,4,4,)}(T_{25},t)]\\ [+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)] & [+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)] & [+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)]\\ [+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)] & [+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)] & [+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)]\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}[+(a''_{33})^{(6)}(T_{33},t)] & [+(a''_{29})^{(5,5,5)}(T_{29},t)] & [+(a''_{25})^{(4,4,4,)}(T_{25},t)]\\ [+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)] & [+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)] & [+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)]\\ [+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)] & [+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)] & [+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)]\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}[+(a''_{34})^{(6)}(T_{33},t)] & [+(a''_{30})^{(5,5,5)}(T_{29},t)] & [+(a''_{26})^{(4,4,4,)}(T_{25},t)]\\ [+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)] & [+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)] & [+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)]\\ [+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)] & [+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)] & [+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)]\end{bmatrix}G_{34} \qquad 87$$

$[+(a''_{32})^{(6)}(T_{33},t)]$,$[+(a''_{33})^{(6)}(T_{33},t)]$,$[+(a''_{34})^{(6)}(T_{33},t)]$ *are first augmentation coefficients for category* 1, 2 *and* 3

$[+(a''_{28})^{(5,5,5)}(T_{29},t)]$,$[+(a''_{29})^{(5,5,5)}(T_{29},t)]$,$[+(a''_{30})^{(5,5,5)}(T_{29},t)]$ *are second augmentation coefficients for category* 1, 2 *and* 3

$[+(a''_{24})^{(4,4,4,)}(T_{25},t)]$,$[+(a''_{25})^{(4,4,4,)}(T_{25},t)]$,$[+(a''_{26})^{(4,4,4,)}(T_{25},t)]$ *are third augmentation coefficients for category* 1, 2 *and* 3

$[+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)]$,$[+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)]$,$[+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)]$ - are fourth augmentation coefficients

$[+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)]$,$[+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)]$,$[+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)]$ - fifth augmentation coefficients

$[+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)]$,$[+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)]$,$[+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)]$ sixth augmentation coefficients

$[+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)]$,$[+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)]$,$[+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)]$ seventh augmentation coefficients

$[+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)]$,$[+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)]$,$[+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)]$ Eighth augmentation coefficients

$[+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)]$,$[+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)]$,$[+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)]$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad (91)$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad (92)$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$\frac{dG_{38}}{dt}$ *93*

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for

category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}+(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}+(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}+(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third

augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$,$+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$,$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients

for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)} & -(b''_{44})^{(9)}(G_{47},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)} & -(b''_{45})^{(9)}(G_{47},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)} & -(b''_{46})^{(9)}(G_{47},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15$ *97*

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$

$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2 \to \infty}(a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)}$ 98

$\lim_{G \to \infty}(b_i'')^{(1)}(G, t) = (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$

Where we suppose

(MMMM) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(NNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(OOOO) $\lim_{T_2 \to \infty} \ (a_i'')^{(7)}\ (T_{37}, t) = (p_i)^{(7)}$

(PPPP)

$\lim_{G \to \infty} (b_i'')^{(7)}\ \big((G_{39}), t\big) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$

$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(QQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(RRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants

$(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\widehat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)} (T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)} \left((G_{43}), t\right) = \ (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\widehat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\left((G_{43}), t\right)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\widehat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\widehat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, \ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}, \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , $T_i : \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{13})^{(1)}$, $T_i^0 \leq (\hat{Q}_{13})^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)},\ T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{24})^{(4)}$, $T_i^0 \leq (\hat{Q}_{24})^{(4)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(p) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \tag{180}$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right] \tag{181}$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose $\tag{182}$

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric $\tag{185}$

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\le \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 19 to 24 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\le\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t),t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\le \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)},i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 216

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(3)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3:$ 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b''_{22})^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\le \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}((G_{31}),(T_{31}))$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right),\quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\le \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 250

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(6)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1,\left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right),\ \ t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying

Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:(\widetilde{(G_{39})},\widetilde{(T_{39})})=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)})d\left(((G_{39})^{(1)},(T_{39})^{(1)};(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 261

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(7)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

268

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{40})^{(8)}\right]\leq(\hat{Q}_{40})^{(8)}$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}\ and\ (\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(8)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying
$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)}=(\nu_0)^{(2)},(m_1)^{(2)}=(\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)}<(\nu_1)^{(2)}$ 305

$(m_2)^{(2)}=(\nu_1)^{(2)},(m_1)^{(2)}=(\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)}<(\nu_0)^{(2)}<(\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)}=\frac{\mathrm{G}_{16}^{0}}{\mathrm{G}_{17}^{0}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})}\left[e^{((S_1)^{(3)}-(p_{20})^{(3)})t}-e^{-(S_2)^{(3)}t}\right]+G_{22}^{0}e^{-(S_2)^{(3)}t}\le G_{22}(t)\le\right. \quad 324$$

$$\left.\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}\right]+G_{22}^{0}e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le\frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\le T_{22}(t)\le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$

$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$

$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)}\le-(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\le-(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)}\le-(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\le-(\tau_1)^{(4)}$

Definition of $(\nu_1)^{(4)},(\nu_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},\nu^{(4)},u^{(4)}$: 329

By $(\nu_1)^{(4)}>0$, $(\nu_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2+(\sigma_1)^{(4)}\nu^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{\nu}_1)^{(4)},,(\bar{\nu}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)}>0$, $(\bar{\nu}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2+(\sigma_2)^{(4)}\nu^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},\ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the

roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$
$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40} e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40} e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}$, $(\mu_2)^{(9)}$, $(v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}$, $(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}\left((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)}\right)}\left[e^{\left((S_1)^{(9)} - (p_{44})^{(9)}\right)t} - e^{-(S_2)^{(9)}t}\right] + G_{46}^0 e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}\left((S_1)^{(9)} - (a'_{46})^{(9)}\right)}\left[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}\left((R_1)^{(9)} - (b'_{46})^{(9)}\right)}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}\left((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)}\right)}\left[e^{\left((R_1)^{(9)} + (r_{44})^{(9)}\right)t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \leq \frac{dv^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \leq (\bar{v}_1)^{(1)}$$

386

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(\mathrm{C})^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(\mathrm{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{\mathrm{C}})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(\mathrm{C})^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(\mathrm{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{\mathrm{C}})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{\mathrm{C}})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \le \frac{d\nu^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \ge \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \le \nu^{(3)}(t) \le (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \le \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \le \nu^{(3)}(t) \le (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \le \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \le \nu^{(3)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \le (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \le (\bar{\nu}_1)^{(3)} \le (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right) - (a''_{25})^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

415

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{32}'')^{(6)} = (a_{33}'')^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b_{32}'')^{(6)} = (b_{33}'')^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a_{36}')^{(7)} - (a_{37}')^{(7)} + (a_{36}'')^{(7)}(T_{37}, t)\right) - (a_{37}'')^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition

$(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

For $0 < \boxed{(v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}$, $\boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $v^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq v^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

with $(p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0$ 475

$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0$ 476

$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0$ 477

$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0$ 478

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(p) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T_{14}^{*} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^{*} for which $f(T_{14}^{*}) = 0$. With this value , we obtain from the three first

equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that

there exists a unique T_{37}^{*} for which $f(T_{37}^{*})=0$. With this value , we obtain from the three first equations

$$G_{36}=\frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}=\frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0)<0, f(\infty)>0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*})=0$. With this value , we obtain from the three first equations

$$G_{40}=\frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}=\frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0)<0, f(\infty)>0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*})=0$. With this value , we obtain from the three first equations

$$G_{44}=\frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}=\frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G)=(b'_{13})^{(1)}(b'_{14})^{(1)}-(b_{13})^{(1)}(b_{14})^{(1)}-$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G)+(b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G)=0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0)>0$, $\varphi(\infty)<0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*})=0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19})=(b'_{16})^{(2)}(b'_{17})^{(2)}-(b_{16})^{(2)}(b_{17})^{(2)}-$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19})+(b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19})=0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0)>0$, $\varphi(\infty)<0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*})=0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23})=(b'_{20})^{(3)}(b'_{21})^{(3)}-(b_{20})^{(3)}(b_{21})^{(3)}-$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^{*})]}$, $T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^{*})]}$ 517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^{*})]}$, $G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^{*})]}$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and A

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$\frac{\partial(a_{17}'')^{(2)}}{\partial \mathrm{T}_{17}}(\mathrm{T}_{17}^{*}) = (q_{17})^{(2)}$, $\frac{\partial(b_i'')^{(2)}}{\partial \mathrm{G}_j}(\,(G_{19})^{*}\,) = s_{ij}$ 533

taking into account equations and neglecting the terms of power 2, we obtain

$\frac{\mathrm{d}\mathbb{G}_{16}}{\mathrm{dt}} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}\mathrm{G}_{16}^{*}\mathbb{T}_{17}$ 534

$\frac{\mathrm{d}\mathbb{G}_{17}}{\mathrm{dt}} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}\mathrm{G}_{17}^{*}\mathbb{T}_{17}$ 535

$\frac{\mathrm{d}\mathbb{G}_{18}}{\mathrm{dt}} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}\mathrm{G}_{18}^{*}\mathbb{T}_{17}$ 536

$\frac{\mathrm{d}\mathbb{T}_{16}}{\mathrm{dt}} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\big(s_{(16)(j)}\mathrm{T}_{16}^{*}\mathbb{G}_j\big)$ 537

$\frac{\mathrm{d}\mathbb{T}_{17}}{\mathrm{dt}} = -\big((b_{17}')^{(2)} - (r_{17})^{(2)}\big)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\big(s_{(17)(j)}\mathrm{T}_{17}^{*}\mathbb{G}_j\big)$ 538

$\frac{\mathrm{d}\mathbb{T}_{18}}{\mathrm{dt}} = -\big((b_{18}')^{(2)} - (r_{18})^{(2)}\big)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\big(s_{(18)(j)}\mathrm{T}_{18}^{*}\mathbb{G}_j\big)$ 539

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\,\mathbb{R}_{+})$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^{*}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}(\,(G_{23})^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$\frac{d\mathbb{G}_{20}}{dt} = -\big((a_{20}')^{(3)} + (p_{20})^{(3)}\big)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^{*}\mathbb{T}_{21}$ 541

$\frac{d\mathbb{G}_{21}}{dt} = -\big((a_{21}')^{(3)} + (p_{21})^{(3)}\big)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^{*}\mathbb{T}_{21}$ 542

$\frac{d\mathbb{G}_{22}}{dt} = -\big((a_{22}')^{(3)} + (p_{22})^{(3)}\big)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^{*}\mathbb{T}_{21}$ 543

$\frac{d\mathbb{T}_{20}}{dt} = -\big((b_{20}')^{(3)} - (r_{20})^{(3)}\big)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\big(s_{(20)(j)}T_{20}^{*}\mathbb{G}_j\big)$ 544

$\frac{d\mathbb{T}_{21}}{dt} = -\big((b_{21}')^{(3)} - (r_{21})^{(3)}\big)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\big(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\big)$ 545

$\frac{d\mathbb{T}_{22}}{dt} = -\big((b_{22}')^{(3)} - (r_{22})^{(3)}\big)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\big(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\big)$ 546

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \qquad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \qquad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \qquad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \qquad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \qquad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \qquad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^*\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\right)\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FOURTEEN

Pure Consciousness (Cessation Of Dualities: Initial Conditions For The Creation Of Universe) And Individual Consciousness (Individual General Ledger)

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

(1) Citi: Universal Consciousness (citi) is (=) the fundamental stuff of the universe. This Consciousness is one and includes the whole. It could also be called God or Shiva. We have adduced this as cosmic general ledger
(2) Mala: Consciousness contracts (e&eb) itself.
(3) The one becomes (=) many.
(4) Shiva becomes (=) the individual (jīva).
(5) This contraction (e) is called mala (impurity).
(6) There are three malas, the mala of individuation (Āṇava mala), the mala of the limited mind (māyīya mala), and the mala of the body (karma mala).
(7) Upāya: An individual caught in the suffering of embodied existence, afflicted by (e&eb) the three malas, and eventually yearns to return to (eb) his or her primordial state of Universal Consciousness.
(8) To attain this, he or she undertakes (e&eb) seemingly difficult sādhana or spiritual practice

NOTATION

Module One

Chit Shakti is one responsible for creation and maya Shakti is illusion which is responsible for producing nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of status of no mind condition). This is my interpretation. There is no conflict between the two

G_{13} : Category one of creation and maya Shakti is illusion which is responsible for (e) producing nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of status of no mind condition). This is my interpretation. **There is no conflict between the two**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Chit Shakti is one responsible

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Chit Shakti is one responsible for creation and maya Shakti is illusion which is responsible for producing nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of status of no mind condition). This is my interpretation. There is no conflict between the two

G_{16} : Category one of nescience (non identification of oneself with the pure consciousness; lack of knowledge or achievement of **status of no mind condition**). This is my interpretation. There is no conflict between the two

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Chit Shakti is one responsible for creation and maya Shakti is illusion

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Citi: Universal Consciousness (citi) is (=) the fundamental stuff of the universe. This Consciousness is one and includes the whole. It could also be called God or Shiva. We have adduced this as cosmic general ledger

G_{20} : Category one of Universal Consciousness (citi)

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of fundamental stuff of the universe

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

G_{24} : Category one of Universal Consciousness (citi)

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of cosmic general ledger

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Mala: Consciousness contracts (e&eb) itself.

G_{28} : Category one of Consciousness

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Consciousness

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Shiva (Pure consciousness) becomes (=) the individual (jīva).

G_{32} : Category one of Shiva (Pure consciousness)

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of individual (jīva).

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Mala of individuation (Āṇava mala), the mala of the limited mind (māyīya mala), and the mala of the body (karma mala)

G_{36} : Category one of Mala of individuation (Āṇava mala); mala of the limited mind (māyīya mala)

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of mala of the limited mind (māyīya mala);Mala of individuation (Āṇava mala);

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

mala of the limited mind (māyīya mala), and the mala of the body (karma mala)

G_{40} : Category one of mala of the limited mind (māyīya mala); mala of the body (karma mala)

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of mala of the body (karma mala);mala of the limited mind (māyīya mala

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Mala of individuation (Āṇava mala), the mala of the limited mind (māyīya mala), and the mala of the body (karma mala)

G_{44} : Category one of Mala of individuation (Āṇava mala); mala of the body (karma mala)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of mala of the body (karma mala);Mala of individuation (Āṇava mala),

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\left[+(a''_{20})^{(3)}(T_{21},t)\right]\left[+(a''_{16})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{13})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{40})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{44})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\left[+(a''_{21})^{(3)}(T_{21},t)\right]\left[+(a''_{17})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{14})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{37})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{41})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{45})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\left[+(a''_{22})^{(3)}(T_{21},t)\right]\left[+(a''_{18})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{15})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{38})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{42})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{46})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & \boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} \quad \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & \boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} \quad \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & \boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} \quad \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(SSSS) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(TTTT) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(UUUU) $\lim_{T_2 \to \infty} \ (a_i'')^{(7)}\,(T_{37}, t) = (p_i)^{(7)}$
(VVVV)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(WWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(XXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(q) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right| \; e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}} \left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(q) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^{*}=\frac{(b_{28})^{(5)}T_{29}^{*}}{[(b_{28}')^{(5)}-(b_{28}'')^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*}=\frac{(b_{30})^{(5)}T_{29}^{*}}{[(b_{30}')^{(5)}-(b_{30}'')^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*})=0$, T_{33}^{*} given by $f(T_{33}^{*})=0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a_{32}')^{(6)}+(a_{32}'')^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a_{34}')^{(6)}+(a_{34}'')^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b_{32}')^{(6)}-(b_{32}'')^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b_{34}')^{(6)}-(b_{34}'')^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a_{36}')^{(7)}+(a_{36}'')^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a_{38}')^{(7)}+(a_{38}'')^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b_{36}')^{(7)}-(b_{36}'')^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b_{38}')^{(7)}-(b_{38}'')^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a_{40}')^{(8)}+(a_{40}'')^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a_{42}')^{(8)}+(a_{42}'')^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b_{40}')^{(8)}-(b_{40}'')^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b_{42}')^{(8)}-(b_{42}'')^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a_{44}')^{(9)}+(a_{44}'')^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a_{46}')^{(9)}+(a_{46}'')^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b_{44}')^{(9)}-(b_{44}'')^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b_{46}')^{(9)}-(b_{46}'')^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTEEN

Anuttara

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

(1) While most other paths observe offering incense and external objects to the deity, this path takes on (e) to offering breaths.
(2) The individual controls (e&eb) his heart and pulse by reducing it significantly.
(3) The final stage is renouncing (e) consumption of food and water.
(4) As a result, he/she connects the state of the supreme in the form of Shiva which results in (eb) purification of the body and generation of **ojas.**
(5) However, Kashmir Shaivism describes four methods (upāya-s): āṇavopāya, the method of the body, śaktopāya, the method of the mind, śāmbhavopāya, the method of Consciousness, and anupāya the 'methodless' method.

NOTATION

Module One

Upāya: An individual caught in the suffering of embodied existence, afflicted by the three malas, and eventually yearns to return to (eb) his or her primordial state of Universal Consciousness

G_{13} : Category one of individual caught in the suffering of embodied existence; three malas, and eventually yearns to return to (eb) his or her primordial state of Universal Consciousness

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of three malas, and eventually yearns to return to (eb) his or her primordial state of Universal Consciousness; individual caught in the suffering of embodied existence

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

individual caught in the suffering of embodied existence, afflicted by the three malas, and eventually yearns to return to (eb) his or her primordial state of Universal Consciousness

G_{16} : Category one of individual caught in the suffering of embodied existence, afflicted by the three malas, and eventually yearns to return

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of his or her primordial state of Universal Consciousness

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

While most other paths observe offering incense and external objects to the deity, this path takes on (e) to offering breaths

G_{20} : Category one of **offering breaths (Breathe properly)**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of most other paths observe offering incense and external objects to the deity, this path takes on

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

individual controls (e&eb) his heart and pulse by reducing it significantly

G_{24} : Category one of individual; heart and pulse by reducing it significantly

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of heart and pulse by reducing it significantly; individual

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

As a result, he/she connects the state of the supreme in the form of Shiva which results in (eb) purification of the body and generation of **ojas.**

G_{28} : Category one of he/she connects the state of the supreme in the form of Shiva

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of purification of the body and generation of **ojas.**

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Kashmir Shaivism describes four methods (upāya-s): āṇavopāya, the method of the body, śaktopāya, the method of the mind, śāmbhavopāya, the method of Consciousness, and anupāya the 'methodless' method.

G_{32} : Category one of āṇavopāya, the method of the body; śaktopāya, the method of the mind, śāmbhavopāya

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of śaktopāya, the method of the mind, śāmbhavopāya; āṇavopāya, the method of the body

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

śāmbhavopāya, the method of Consciousness, and anupāya the 'methodless' method

G_{36} : Category one of śāmbhavopāya, the method of Consciousness; anupāya the 'methodless' method

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of anupāya the 'methodless' method ;śāmbhavopāya, the method of Consciousness

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Āṇavopāya, the method of the body and anupāya the 'methodless' method

G_{40} : Category one of Āṇavopāya, the method of the body; anupāya the 'methodless' method

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of anupāya the 'methodless' method; Āṇavopāya, the method of the body

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Fruit of the individual's sādhana is the attainment of Self-realisation (mokṣa), the cessation if dualities to become Shiva

G_{44} : Category one of Fruit of the individual's sādhana

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of attainment of Self-realisation (mokṣa), the cessation if dualities to become Shiva

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21},t)+(a''_{16})^{(2,2,2)}(T_{17},t)+(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)+(a''_{40})^{(8,8,8,8)}(T_{41},t)+(a''_{44})^{(9,9,9)}(T_{45},t) \end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}+(a''_{21})^{(3)}(T_{21},t)+(a''_{17})^{(2,2,2)}(T_{17},t)+(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)+(a''_{41})^{(8,8,8,8)}(T_{41},t)+(a''_{45})^{(9,9,9)}(T_{45},t) \end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21},t)+(a''_{18})^{(2,2,2)}(T_{17},t)+(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)+(a''_{42})^{(8,8,8,8)}(T_{41},t)+(a''_{46})^{(9,9,9)}(T_{45},t) \end{array}\right]G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{c} (b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23},t)-(b''_{16})^{(2,2,2)}(G_{19},t)-(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t)-(b''_{40})^{(8,8,8,8)}(G_{43},t)-(b''_{44})^{(9,9,9)}(G_{47},t) \end{array}\right]T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{c} (b'_{21})^{(3)}-(b''_{21})^{(3)}(G_{23},t)-(b''_{17})^{(2,2,2)}(G_{19},t)-(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t)-(b''_{41})^{(8,8,8,8)}(G_{43},t)-(b''_{45})^{(9,9,9)}(G_{47},t) \end{array}\right]T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$

$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}\ \ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\ ,\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\left[\begin{matrix}(a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\left[\begin{matrix}(a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\left[\begin{matrix}(a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{34} \qquad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ ,\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ are\ second\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}\ \ are\ third\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation

coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \quad +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \quad -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \quad -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \quad -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(YYYY) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(ZZZZ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(AAAAA) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(BBBBB)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(r) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(r) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b_i'')^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTEEN A

"Unsurpassed Reality"

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

(1) Mokṣa: The fruit of the individual's sādhana is the attainment of Self-realisation (mokṣa). In Kashmir Shaivism, the state of liberation (mukti) is called sahaja samadhi and is characterised by (e) the attainment of unwavering bliss-consciousness while living one's ordinary life. It is stated that such a bliss consciousness also knows what suffering-unconsciousness is and hence the differentiation becomes integration.

(2) Anuttara is the ultimate principle in Kashmir Shaivism, and as such, it is the fundamental reality underneath (e&eb) the whole Universe.

(3) Among the multiple interpretations of Anuttara are: "supreme", "above all" and "unsurpassed reality". In the Sanskrit alphabet Anuttara is associated to the first letter - "A" (in Devanagari "अ"). As the ultimate principle, Anuttara is identified (e&eb) with Śiva, Śakti (as Śakti is identical to Śiva), the supreme consciousness (cit), uncreated light (prakāśa), supreme subject (aham) and atemporal vibration (Spanda).

(4) The practitioner who realizes Anuttara through any means, whether by her own efforts or by direct transmission by the Grace of Shiva/shakti, is liberated and perceives absolutely no difference between herself and (=) the body of the universe. Simple idea of I = cosmic general ledger

NOTATION

Module One

state of liberation (mukti) is called sahaja samadhi

G_{13} : Category one of state of liberation (mukti)

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of sahaja samadhi

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Sahaja samadhi is characterised by (e) the attainment of unwavering bliss-consciousness while living one's ordinary life. It is stated that such a bliss consciousness also knows what suffering-unconsciousness is and hence the differentiation becomes integration

G_{16} : Category one of attainment of unwavering bliss-consciousness while living one's ordinary life. It is stated that such a bliss consciousness also knows what suffering-unconsciousness is and hence the differentiation becomes integration

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Sahaja samadhi is characterised

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

. It is stated that such a bliss consciousness also knows what suffering-unconsciousness is and hence the differentiation becomes integration

G_{20} : Category one of knowing truth

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of knowing untruth

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

G_{24} : Category one of knowing bliss

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of knowing despair

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

G_{28} : Category one of knowing Ananada

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of knowing despondency

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

G_{32} : Category one of knowing Satchidananda

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of knowing what is UN Satchidananda

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Anuttara is the ultimate principle in Kashmir Shaivism, and as such, it is the fundamental reality underneath (e&eb) the whole Universe

G_{36} : Category one of Anuttara

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of ultimate principle in Kashmir Shaivism

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

it is the fundamental reality underneath the whole Universe

G_{40} : Category one of Anuttara; fundamental reality underneath the whole Universe

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of fundamental reality underneath the whole Universe ;Anuttara

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Among the multiple interpretations of Anuttara are: "supreme", "above all" and "unsurpassed reality". In the Sanskrit alphabet Anuttara is associated to the first letter - "A" (in Devanagari "अ"). As the ultimate principle, Anuttara is identified (e&eb) with Śiva, Śakti (as Śakti is identical to Śiva), the supreme consciousness (cit), uncreated light (prakāśa), supreme subject (aham) and atemporal vibration (Spanda).

G_{44} : Category one of Anuttara; Śiva, Śakti (as Śakti is identical to Śiva), the supreme consciousness (cit), uncreated light (prakāśa), supreme subject (aham) and atemporal vibration (Spanda).

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Śiva, Śakti (as Śakti is identical to Śiva), the supreme consciousness (cit), uncreated light (prakāśa), supreme subject (aham) and atemporal vibration (Spanda).;Anuttara

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)}\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)}\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)}\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(EEEEE) $\qquad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(FFFFF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(GGGGG) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(HHHHH)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(IIIII) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(JJJJJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(s) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28}, G_{30} and G_{28}, G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30}. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}} \left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)} (T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(s) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTEEN

Svatantrya, Self-Created Free Will

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

(1) Being and (=) beings become one and the same by virtue of the "erotic friction," whereby subject perceives (e) object and in that act of perception is filled with (e&eb) nondual being/consciousness/bliss.
(2) Being and (=) beings become one and the same by virtue of the "erotic friction," whereby subject perceives (e) object and in that act of perception is filled with (e&eb) nondual being/consciousness/bliss
(3) Being and (=) beings become one and the same by virtue of the "erotic friction," whereby subject perceives (e) object and in that act of perception is filled with (e&eb) nondual being/consciousness/bliss
(4) Anuttara is different from the notion of transcendence in that, even though it is above all, it does not (e) imply a state of separation from the Universe.

Aham

(5) Aham is the concept of supreme reality as (=) heart. It is considered to be a non-dual interior space of Śiva (topological space of pure consciousness where all dualities cease space=time, being=non being, self=Nonself, chaos=order: see earlier models; one analogy is topological space of vacuum energy), support for the entire manifestation, supreme mantra and identical to Śakti (phenomenal manifestation of the divided universe). Note in the final and ultimate analysis the emphasis is on pure consciousness=impure consciousness the initial condition for creation of the universe.
(6) Aham is the concept of supreme reality as (=) heart. It is considered to be a non-dual interior space of Śiva (topological space of pure consciousness where all dualities cease space=time, being=non being, self=Nonself, chaos=order: see earlier models; one analogy is topological space of vacuum energy), support for the entire manifestation, supreme mantra and identical to Śakti (phenomenal manifestation of the divided universe). Note in the final and ultimate analysis the emphasis is on pure consciousness=impure consciousness the initial condition for creation of the universe

Pratyabhijnā

(7) Pratyabhijña is the philosophical articulation of Kashmir Saivism. Pratyabhijña literally means (eb)"spontaneous recognition", as it does not have any upāyas (means), that is, there is (=) nothing to practice; the only thing to do is recognize (e&eb) who you are. This "means" can actually be called anupāya, Sanskrit for "without means". Ksemaraja, the student of Abhinavagupta, uses a mirror analogy to explain Pratyabhijña. Note mirror neurons are detected recently. Look at yourself, say like I am doing through these papers.
(8) Pratyabhijña is the philosophical articulation of Kashmir Saivism. Pratyabhijña literally means (eb)"spontaneous recognition", as it does not have any upāyas (means), that is, there is (=) nothing to practice; the only thing to do is recognize (e&eb) who you are. This "means" can actually be called anupāya, Sanskrit for "without means". Ksemaraja, the student of Abhinavagupta, uses a mirror analogy to explain Pratyabhijña. Note mirror neurons are detected recently. Look at yourself ,say like I am doing through these papers

Svatantrya, self-created free will

(9) The concept of free will plays a central role in Kashmir Shaivism. Known technically as svātantrya it is the cause (eb) of the creation of the universe - a primordial force that stirs up (e&eb) the absolute and manifests the world inside the supreme consciousness of Siva. It is here Pure consciousness (zero) reflects upon itself its nature and divides in to various transactional entities.

NOTATION

Module One

practitioner who realizes Anuttara through any means, whether by her own efforts or by direct transmission by the Grace of Shiva/shakti, is liberated and perceives absolutely no difference between herself and (=) the body of the universe. Simple idea of I = cosmic general ledger

G_{13} : Category one of I

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of cosmic general ledger

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Being and beings become one and the same by virtue of the "erotic friction," whereby subject perceives object and in that act of perception is filled with (e&eb) nondual being/consciousness/bliss

G_{16} : Category one of object and in that act of perception is filled with (e&eb) nondual being/consciousness/bliss

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Being and (=) beings become one and the same by virtue of the "erotic friction," whereby subject perceives

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Being and beings become one and the same by virtue of the "erotic friction," whereby subject perceives object and in that act of perception is filled with (e&eb) nondual being/consciousness/bliss

G_{20} : Category one of Being and beings become one and the same by virtue of the "erotic friction," whereby subject perceives object and in that act of perception; nondual being/consciousness/bliss

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of nondual being/consciousness/bliss ;Being and beings become one and the same by virtue of the "erotic friction," whereby subject perceives object and in that act of perception

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Anuttara is different from the notion of transcendence in that, even though it is above all, it does not (e) imply a state of separation from the Universe

G_{24} : Category one of imply a state of separation from the Universe

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Anuttara is different from the notion of transcendence in that, even though it is above all

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Aham is the concept of supreme reality as (=) heart. It is considered to be a non-dual interior space of Śiva (topological space of pure consciousness where all dualities cease space=time, being=non being, self=Nonself, chaos=order: see earlier models; one analogy is topological space of vacuum energy), support for the entire manifestation, supreme mantra and identical to Śakti (phenomenal manifestation of the divided universe). Note in the final and ultimate analysis the emphasis is on pure consciousness=impure consciousness the initial condition for creation of the universe

G_{28} : Category one of Aham is the concept of supreme reality

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of heart (**non-dual interior space of Śiva (topological space of pure consciousness where all dualities cease space=time, being=non being, self=Nonself**, chaos=order: see earlier models; one analogy is topological space of vacuum energy), support for the entire manifestation, supreme mantra and identical to Śakti (phenomenal manifestation of the divided universe). Note in the final and ultimate analysis the emphasis is on pure consciousness=impure consciousness the initial condition for creation of the universe)

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Aham is the concept of supreme reality as (=) heart. It is considered to be a non-dual interior space of Śiva (topological space of pure consciousness where all dualities cease space=time, being=non being, self=Nonself, chaos=order: see earlier models; one analogy is topological space of vacuum energy), support for the entire manifestation, supreme mantra and identical to Śakti (phenomenal manifestation of the divided universe). Note in the final and ultimate analysis the emphasis is on pure consciousness=impure consciousness the initial condition for creation of the universe

G_{32} : Category one of pure consciousness

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of impure consciousness the initial condition for creation of the universe

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Pratyabhijña is the philosophical articulation of Kashmir Saivism. Pratyabhijña literally means "spontaneous recognition", as it does not have any upāyas (means), that is, there is (=) nothing to practice; the only thing to do is recognize (e&eb) who you are. This "means" can actually be called anupāya, Sanskrit for "without means". Ksemaraja, the student of Abhinavagupta, uses a mirror analogy to explain Pratyabhijña. Note mirror neurons are detected recently. Look at yourself, say like I am doing through these papers.

G_{36} : Category one of Pratyabhijña

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of)"spontaneous recognition", as it does not have any upāyas (means), that is, there is (=) nothing to practice

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Pratyabhijña is the philosophical articulation of Kashmir Saivism. Pratyabhijña literally means "spontaneous recognition", as it does not have any upāyas (means), that is, there is nothing to practice; the only thing to do is recognize (e&eb) who you are. This "means" can actually be called anupāya, Sanskrit for "without means". Ksemaraja, the student of Abhinavagupta, uses a mirror analogy to explain Pratyabhijña. Note mirror neurons are detected recently. Look at yourself, contemplate on yourself, introspect on self say like I am doing through these papers

G_{40} : Category one of Pratyabhijña is the philosophical articulation of Kashmir Saivism. Pratyabhijña literally means "spontaneous recognition", as it does not have any upāyas (means), that is, there is nothing to practice; the only thing to do is recognize; who you are (you are shiva the pure consciousness)

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of who you are (you are shiva the pure consciousness); Pratyabhijña is the philosophical articulation of Kashmir Saivism. Pratyabhijña literally means "spontaneous recognition", as it does not have any upāyas (means), that is, there is nothing to practice; the only thing to do is recognize

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

The concept of free will plays a central role in Kashmir Shaivism. Known technically as svātantrya it is the cause (eb) of the creation of the universe - a primordial force that stirs up (e&eb) the absolute and manifests the world inside the supreme consciousness of Siva. It is here Pure consciousness (zero) reflects upon itself its nature and divides in to various transactional entities.

G_{44} : Category one of **concept of free will plays a central role in Kashmir Shaivism. Known technically as svātantrya it is the cause (we are talking of the freewill of absolute subjectivity, pure consciousness; non dual state;**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **absolute and manifests the world inside the supreme consciousness of Siva.**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}), t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{c} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{array}\right] T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{c} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{array}\right] T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{c} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{array}\right] T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category

1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t)+(a_{16}'')^{(2,2,2)}(T_{17},t)+(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)+(a_{40}'')^{(8,8,8,8)}(T_{41},t)+(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t)+(a_{17}'')^{(2,2,2)}(T_{17},t)+(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)+(a_{41}'')^{(8,8,8,8)}(T_{41},t)+(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t)+(a_{18}'')^{(2,2,2)}(T_{17},t)+(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)+(a_{42}'')^{(8,8,8,8)}(T_{41},t)+(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad (85)$$

$$- \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b_{28}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b_{24}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}, \boxed{(a_{37}'')^{(7)}(T_{37},t)}, \boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\left[-(b''_{36})^{(7)}(G_{39},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\left[-(b''_{37})^{(7)}(G_{39},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\left[-(b''_{38})^{(7)}(G_{39},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(KKKKK) $\qquad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(LLLLL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(MMMMM) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(NNNNN)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(OOOOO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(PPPPP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\left((G_{43}), t\right) = (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\left((G_{43}), t\right)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(t) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0$ for $\mathrm{t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})($and\ not\ on\ \ t$) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(t) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVENTEEN

Function Of Svātantrya Is Self Reflection And Contemplation Of One's Nature

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

(1) In Svātantrya all conscious subjects are (=) co-participant in various degrees to the divine sovereignty. (Co creativity)
(2) Humans have a degree of free will limited by (e) their level of consciousness.
(3) Ultimately, Kashmir Shaivism as a monistic idealist philosophical system views all subjects to be identical - "all are (=) one" - and that one is (=) Śiva, the supreme (Pure) consciousness.
(4) Thus, all subjects have (e) free will and are god/divine but can be ignorant of this.
(5) Ignorance too is (=) a force projected by svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge. Note Krishna telling Arjuna that he is ignorance in ignorant and intelligence of intelligent. he is manifested in three forms rajas, tamas and Sattva
(6) One **function of svātantrya is (=) granting divine grace** - śaktipāt.
(7) In this philosophical system, spiritual liberation IS accessible by mere effort, but can be guided by the will/grace of god (i.e. the liberated, the masters). Thus, if the disciple finds such a master, he need only surrender himself and await divine grace to eliminate (e) the limitations that imprison his consciousness.
(8) **Causality in Kashmir Shaivism is considered to be created by (e) Svātantrya along with the universe**.
(9) Thus there can be no contradiction, limitation or rule to force (eb) Śiva to act one way or the other.
(10) Svātantrya always exists beyond (e) the limiting shield of cosmic illusion, māyā.

NOTATION

Module One

Svātantrya all conscious subjects are co-participant in various degrees to the divine sovereignty. (Co creativity)

G_{13} : Category one of Svātantrya all conscious subjects

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of co-participant in various degrees to the divine sovereignty. (Co creativity)

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Humans have a degree of free will limited by (e) their level of consciousness

G_{16} : Category one of level of consciousness

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Humans have a degree of free will

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Kashmir Shaivism as a monistic idealist philosophical system views all subjects to be identical - "all are (=) one" - and that one is (=) Śiva, the supreme (Pure) consciousness

G_{20} : Category one of monistic idealist philosophical system views all subjects to be identical - "all are

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of one" - and that one is (=) Śiva, the supreme (Pure) consciousness

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

All subjects have (e) free will and are god/divine but can be ignorant of this. Note that the freewill mentioned here is that of Pure consciousness. Absolute subjectivity has freewill and human beings have less choice to remain uncoordinated with!

G_{24} : Category one of free will and are god/divine but can be ignorant of this. Note that the freewill mentioned here is that of Pure consciousness. Absolute subjectivity has freewill and human beings have less choice to remain uncoordinated with!

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of All subjects

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Ignorance too is a force projected by svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge. **Note Krishna telling Arjuna that he is ignorance in ignorant and intelligence of intelligent. He is manifested in three forms rajas, tamas and Sattva** (written out of memory. I donot state to be emphatic or self authoritarian)

G_{28} : Category one of Ignorance

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of force projected by svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge.

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Ignorance too is a force projected by svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge

G_{32} : Category one of Ignorance too is a force projected; svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge; Ignorance too is a force projected

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Ignorance too is a force projected by svātantrya itself upon the creation and can be removed by svātantrya and also by Self-knowledge

G_{36} : Category one of svātantrya and also by Self-knowledge

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Ignorance too is a force projected by svātantrya itself upon the creation

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

function of svātantrya is (=) granting divine grace - śaktipāt.(take this a deterministic future)

G_{40} : Category one of **function of svātantrya**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **granting divine grace** – śaktipāt

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Causality in Kashmir Shaivism is considered to be created by (e) Svātantrya along with the universe

G_{44} : Category one of **Svātantrya along with the universe**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **Causality in Kashmir Shaivism is considered to be created**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18}$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{ccc} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{ccc} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, \ (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(QQQQQ) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(RRRRR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(SSSSS) $\lim_{T_2 \to \infty} \ (a_i'')^{(7)}\,(T_{37}, t) = (p_i)^{(7)}$
(TTTTT)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(UUUUU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(VVVVV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(u) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})($and\ not\ on\ \ t$) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(u) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION EIGHTEEN

Spanda

INTRODUCTION—VARIABLES USED

Kashmir Shaivism from Wikipedia

Spanda

(1) The Spanda system, introduced by Vasugupta (c. 800 AD), is usually described as (=) "vibration/movement of consciousness".
(2) It is just the vibration of (e) the information in the cosmic general ledger (information)
(3) Abhinavagupta uses the expression "some sort of movement" to imply the distinction from physical movement; it is rather a vibration or sound inside the Divine, a throb. The essence of this vibration is (=) the ecstatic self-recurrent consciousness. What are we to make out of self recurrent consciousness? I interpret this as the reflection of one's self. When you reflect then only some theory ensues about yours elf about world. As is happening in these papers.
(4) Association in space and time is (=) dangerous and kills truth; I have been continuously shown colours and even now despite knowing I am being fooled, I sometimes tend to confuse the illusion and "play" by Brahman –Anti Brahman with "reality"(?) (my statement)
(5) The central tenet of this system is (=) "everything is Spanda", both the objective exterior reality and the subjective world. You can say quantum mechanical systems for subjective world. Note this is what exactly string Theory says.
(6) Nothing exists without (e) movement, yet the ultimate movement takes place (e) not in space or time, but inside (eb) the Supreme Consciousness (cit). Cessation of dualities is well the initial condition of the creation of the universe. Death is loss of self, but nor achievement of pure consciousness. so let there be any confusion between the initial conditions for the creation of the universe, and the loss of " self, the non monitoring witness consciousness, which becomes pulsating and throbbing only evolutionarily when the ere is development of individual consciousness as is happening in science (see This is a Debit Credit world; Man is moving towards No mind condition)
(7) Nothing exists without (e) movement, yet the ultimate movement takes place (e) not in space or time, but inside (eb) the Supreme Consciousness (cit).
(8) So, it is a cycle of internalization and (e&eb) externalization of consciousness itself, relating to the most Elevated plane in creation (Śiva-Śakti Tattva). There is nothing to glorify the statement. It is almost like join a job and retiring. Involution and evolution has similar meaning. In essence, concentration is on one's self at a later stage.
(9) In order to describe the connotations of the Spanda concept, a series of equivalent concepts are enumerated, such as: **self recurrent consciousness - vimarśa**, unimpeded will of the Supreme Consciousness (cit) - svātantrya, supreme creative energy - visarga, heart of the divine - hṛdaya and ocean of light-consciousness - cidānanda. The most important texts of the system are Śiva Sutras, Spanda Karika and Vijñāna Bhairava Tantra.

NOTATION

Module One

Svātantrya always exists beyond (e) the limiting shield of cosmic illusion, māyā.

G_{13} : Category one of limiting shield of cosmic illusion, māyā

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Svātantrya always exists

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Spanda system, introduced by Vasugupta (c. 800 AD), is usually described as (=) "vibration/movement of consciousness". (Interpret this as information as in String theory)

G_{16} : Category one of Spanda system

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of "vibration/movement of consciousness". (Interpret this as information as in String theory)

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

It is just the vibration of (e) the information in the cosmic general ledger (information)

G_{20} : Category one of Spanda system

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of vibration of the information in the cosmic general ledger (information)

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Abhinavagupta uses the expression "some sort of movement" to imply the distinction from physical movement; it is rather a vibration or sound inside the Divine, a throb. The essence of this vibration is (=) the ecstatic self-recurrent consciousness. **What are we to make out of self recurrent consciousness? I interpret this as the reflection of one's self. When you reflect then only some theory ensues about yours elf about world. As is happening in these papers.**

G_{24} : Category one of essence of this vibration

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of ecstatic self-recurrent consciousness

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Association in space and time is dangerous and kills truth; I have been continuously shown colours and even now despite knowing I am being fooled, I sometimes tend to confuse the illusion and "play" by Brahman –Anti Brahman with "reality"(?) (my statement)

G_{28} : Category one of kills truth; I have been continuously shown colours and even now despite knowing I am being fooled, I sometimes tend to confuse the illusion and "play" by Brahman –Anti Brahman with "reality"(?) (my statement)

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Association in space and time

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Central tenet of this system is (=) "everything is Spanda", both the objective exterior reality and the subjective world. You can say quantum mechanical systems for subjective world. Note this is what exactly string Theory says

G_{32} : Category one of Central tenet of this system

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of "everything is Spanda", both the objective exterior reality and the subjective world. You can say quantum mechanical systems for subjective world. Note this is what exactly string Theory says

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Nothing exists without movement, yet the ultimate movement takes place (e) not in space or time, but inside (eb) the Supreme Consciousness (cit). Cessation of dualities is well the initial condition of the creation of the universe. Death is loss of self, but nor achievement of pure consciousness. so let there be any confusion between the initial conditions for the creation of the universe, and the loss of " self, the non monitoring witness consciousness, which becomes pulsating and throbbing only evolutionarily when the ere is development of individual consciousness as is happening in science (see This is a Debit Credit world; Man is moving towards No mind condition

G_{36} : Category one of movement, yet the ultimate movement takes place (e) not in space or time, but inside (eb) the Supreme Consciousness (cit).

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Nothing exists

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Nothing exists without movement, yet the ultimate movement takes place not in space or time, but inside (eb) the Supreme Consciousness (cit).

G_{40} : Category one of not in space or time, but inside (eb) the Supreme Consciousness (cit).

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Nothing exists without movement, yet the ultimate movement

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Nothing exists without movement, yet the ultimate movement takes place not in space or time, but inside (eb) the Supreme Consciousness (cit).

G_{44} : Category one of Nothing exists without movement, yet the ultimate movement takes place not in space or time,

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Supreme Consciousness (cit). Cessation of dualities is well the initial condition of the creation of the universe. Death is loss of self, but nor achievement of pure consciousness. so let there be any confusion between the initial conditions for the creation of the universe, and the loss of " self, the non monitoring witness consciousness, which becomes pulsating and throbbing only evolutionarily when the ere is development of individual consciousness as is happening in science (see This is a Debit Credit world; Man is moving towards No mind condition

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)}\ \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(WWWWW) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(XXXXX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(YYYYY) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(ZZZZZ)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(AAAAAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(BBBBBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\ \ G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\ \ T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}$, $T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(v) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}\left((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)}\right)}\left[e^{\left((S_1)^{(3)} - (p_{20})^{(3)}\right)t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}\left((S_1)^{(3)} - (a'_{22})^{(3)}\right)}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}\left((R_1)^{(3)} - (b'_{22})^{(3)}\right)}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}\left((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)}\right)}\left[e^{\left((R_1)^{(3)} + (r_{20})^{(3)}\right)t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right) - (b''_{25})^{(4)}\left((G_{27}), t\right) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad (354)$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad (355)$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad (356)$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^{0}e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}\left((R_1)^{(9)}-(b'_{46})^{(9)}\right)}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}\left((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)}\right)}\left[e^{\left((R_1)^{(9)}+(r_{44})^{(9)}\right)t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^{0}}{G_{14}^{0}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(v) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION NINETEEN

Ultra-Secure Internet

INTRODUCTION—VARIABLES USED

Einstein's 'spooky' theory may lead to ultra-secure internet 7 hours ago Some sentences and images are deleted due to spatial constraints Kindly bear with me for the commission of act of omission.

(1) (Phys.org) —Einstein's scepticism about quantum mechanics may lead to (eb) an ultra-secure internet, suggests a new paper by researchers from Swinburne University of Technology and Peking University.
(2) Associate Professor Margaret Reid from Swinburne's Centre for Quantum and Optical Science said Einstein's reservations about quantum mechanics were highlighted in (eb) a phenomenon known as "'spooky' action at a distance."
(3) In 1935 Einstein and researchers highlighted a 'spooky' theory in quantum mechanics, which is the strange way entangled particles stay connected (e&eb) even when separated by large distances.
(4) "Until now the real application of this has been for messages being shared (e&eb) between two people securely without interception, regardless of the spatial separation between them," Professor Reid said.
(5) "In this paper, we give theoretical proof that such messages can be shared (e&eb) between more than two people and may provide (eb) unprecedented security for a future quantum internet."
(6) In the 1990s, scientists realised you can securely transmit a message through encrypting and using (e) a shared key generated by Einstein's strange entanglement to decode (eb) the message from the sender and receiver.
(7) Using the quantum key meant the message was (=) completely secure from interception during transmission.
(8) Sending Einstein's entanglement to a larger number of people means (eb) the key can be distributed among all the receiving parties, so they must collaborate (e&eb) to decipher the message, which Professor Reid said makes the message even more secure.
(9) "We found that a secure message can be shared by up to three to four people, opening (eb) the possibility to the theory being applicable to (e&eb) secure messages being sent from many too many.

Unconventional Programming Paradigms Lecture Notes in Computer Science Volume 3566, 2005, pp 257-269 Autonomic Computing: An Overview Manish Parashar, Salim Hariri

(1) The increasing scale complexity, heterogeneity and dynamism of networks, systems and applications have made our computational and information infrastructure brittle, unmanageable and insecure. This has necessitated the investigation of an alternate paradigm for system and application design, which is based (e) on strategies used by biological systems to deal with similar challenges – a vision that has been referred to as autonomic computing.

(2) The overarching goal of autonomic computing is (=) to realize computer and software systems and applications that can manage (e&eb) themselves in accordance with high-level guidance from humans.

(3) Meeting the grand challenges of autonomic computing requires (e) scientific and technological advances in a wide variety of fields, as well as new software and system architectures that support (eb) the effective integration of the constituent technologies. This paper presents an introduction to autonomic computing, its challenges, and opportunities.

NOTATION

Module One

(Phys.org) —Einstein's scepticism about quantum mechanics may lead to (eb) an ultra-secure internet, suggests a new paper by researchers from Swinburne University of Technology and Peking University

G_{13} : Category one of Einstein's scepticism about quantum mechanics

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of ultra-secure internet

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Einstein and researchers highlighted a 'spooky' theory in quantum mechanics, which is the strange way entangled particles stay connected (e&eb) even when separated by large distances

G_{16} : Category one of ‘spooky' theory in quantum mechanics, which is the strange way entangled particles stay connected; separated by large distances

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of separated by large distances; 'spooky' theory in quantum mechanics, which is the strange way entangled particles stay connected

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

"Until now the real application of this has been for messages being shared (e&eb) between two people securely without interception, regardless of the spatial separation between them," Professor Reid said

G_{20} : Category one of real application of this has been for messages; two people securely without interception, regardless of the spatial separation between them,"

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of two people securely without interception, regardless of the spatial separation between them,"; real application of this has been for messages

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Messages can be shared between more than two people and may provide (eb) unprecedented security for a future quantum internet."

G_{24} : Category one of Messages can be shared between more than two people

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of unprecedented security for a future quantum internet.

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Message can be securely transmitted through encrypting and using a shared key generated by Einstein's strange entanglement to decode (eb) the message from the sender and receiver.

G_{28} : Category one of shared key generated by Einstein's strange entanglement to decode (eb) the message from the sender and receiver.

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Message can be securely transmitted through encrypting

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Message can be securely transmitted through encrypting and using a shared key generated by Einstein's

strange entanglement to decode (eb) the message from the sender and receiver

G_{32} : Category one of Message can be securely transmitted through encrypting and using a shared key generated by Einstein's strange entanglement to decode

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of message from the sender and receiver

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Using the quantum key meant the message was completely secure from interception during transmission

G_{36} : Category one of completely secure from interception during transmission

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of quantum key meant the message

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

This has necessitated the investigation of an alternate paradigm for system and application design, which is based (e) on strategies used by biological systems to deal with similar challenges – a vision that has been referred to as autonomic computing.

G_{40} : Category one of strategies used by biological systems to deal with similar challenges – a vision that has been referred to as autonomic computing.

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of investigation of an alternate paradigm for system and application design

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Overarching goal of autonomic computing is to realize computer and software systems and applications that can manage (e&eb) themselves in accordance with high-level guidance from humans

G_{44} : Category one of Overarching goal of autonomic computing; Overarching goal of autonomic computing is to realize computer and software systems and applications; themselves in accordance with high-level guidance from humans

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of realize computer and software systems and applications that can manage (e&eb) themselves in accordance with high-level guidance from humans; themselves in accordance with high-level guidance from humans ;Overarching goal of autonomic computing is to realize computer and software systems and applications

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15}$ 6

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16}$ 7

$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17}$ 8

$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18}$ 9

$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16}$ 10

$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17}$ 11

$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18}$ 12

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20}$ 13

$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21}$ 14

$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22}$ 15

$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20}$ 16

$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21}$ 17

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\,+(a''_{20})^{(3)}(T_{21},t)\;\; +(a''_{16})^{(2,2,2)}(T_{17},t)\;\; +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{40})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{44})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\,+(a''_{21})^{(3)}(T_{21},t)\;\; +(a''_{17})^{(2,2,2)}(T_{17},t)\;\; +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{41})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{45})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\,+(a''_{22})^{(3)}(T_{21},t)\;\; +(a''_{18})^{(2,2,2)}(T_{17},t)\;\; +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{42})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{46})^{(9,9,9)}(T_{45},t) \end{array}\right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)}\boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)}\boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)}\boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} & \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

Where $\boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a_{40}'')^{(8,8\,,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}-(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}-(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}-(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)},\ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)},\ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(CCCCCC) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(DDDDDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(EEEEEE) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(FFFFFF)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(GGGGGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(HHHHHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(w) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)},\ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)},\ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(w) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY

Multipartite Einstein-Podolsky-Rosen Steering

INTRODUCTION—VARIABLES USED

Einstein's 'spooky' theory may lead to ultra-secure internet 7 hours ago Some sentences and images

are deleted due to spatial constraints Kindly bear with me for the commission of act of omission

(1) "The message will also remain secure if the devices receiving (e) the message have been tampered (e&eb) with, like if an iPhones were hacked, because (e) of the nature of Einstein's spooky entanglement.
(2) "The message will also remain secure if the devices receiving (e) the message have been tampered (e&eb) with, like if an iPhones were hacked, because (e) of the nature of Einstein's spooky entanglement.
(3) "The message will also remain secure if the devices receiving (e) the message have been tampered (e&eb) with, like if an iPhones were hacked, because (e) of the nature of Einstein's spooky entanglement.
(4) "Discovering that it can be applied to a situation with more parties has the potential to create (eb) a more secure internet – with less messages being intercepted (e&eb) from external parties?" The paper was published in Physical Review Letters.

More information: Genuine Multipartite Einstein-Podolsky-Rosen Steering, Q.Y. He and M. D. Reid, Phys. Rev. Lett. 111, 250403 (2013) link.aps.org/doi/10.1103/PhysRevLett.111.250403 Journal reference: Physical Review Letters Provided by Swinburne University of Technology

(5) Authors develop the concept of (e) genuine N-partite Einstein-Podolsky-Rosen (EPR) steering.
(6) This nonlocality is (=) the natural multipartite extension of (e) the original EPR paradox.
(7) Useful properties emerge that are not (e) guaranteed for genuine multipartite entangled states.
(8) In particular, there is a close link with the (e&eb) task of one-sided, device-independent quantum secret sharing.
(9) Authors derive inequalities to demonstrate (eb) multipartite EPR steering for (e) Greenberger-Horne-Zeilinger and Gaussian continuous variable states in (eb) loophole-free scenarios. © 2013 American Physical Society

NOTATION

Module One

Message will also remain secure if the devices receiving (e) the message have been tampered with, like if an iPhones were hacked, because (e) of the nature of Einstein's spooky entanglement.

G_{13} : Category one of message have been tampered (e&eb) with, like if an iPhones were hacked, because (e) of the nature of Einstein's spooky entanglement.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Message will also remain secure if the devices

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Message will also remain secure if the devices receiving the message have been tampered with, like if an iPhones were hacked, because (e) of the nature of Einstein's spooky entanglement

G_{16} : Category one of nature of Einstein's spooky entanglement

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Message will also remain secure if the devices receiving the message have been tampered with, like if an iPhones were hacked

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Discovering that it can be applied to a situation with more parties has the potential to create a more secure internet – with less messages being intercepted (e&eb) from external parties?" The paper was published in Physical Review Letters

G_{20} : Category one of Discovering that it can be applied to a situation with more parties has the potential

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of more secure internet – with less messages being intercepted (e&eb) from external parties?"

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Discovering that it can be applied to a situation with more parties has the potential to create a more secure internet – with less messages being intercepted (e&eb) from external parties?"

G_{24} : Category one of Discovering that it can be applied to a situation with more parties has the potential to create a more secure internet – with less messages being intercepted; external parties

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of external parties ;Discovering that it can be applied to a situation with more parties has the potential to create a more secure internet – with less messages being intercepted

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Authors develop the concept of (e) genuine N-partite Einstein-Podolsky-Rosen (EPR) steering

G_{28} : Category one of genuine N-partite Einstein-Podolsky-Rosen (EPR) steering

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of concept

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

This nonlocality is the natural multipartite extension of (e) the original EPR paradox.

G_{32} : Category one of nonlocality

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of natural multipartite extension of (e) the original EPR paradox

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

nonlocality is the natural multipartite extension of the original EPR paradox

G_{36} : Category one of original EPR paradox

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of nonlocality is the natural multipartite extension

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

There is a close link with the (e&eb) task of one-sided, device-independent quantum secret sharing.

G_{40} : Category one of natural multipartite extension of the original EPR paradox; task of one-sided, device-independent quantum secret sharing

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of task of one-sided, device-independent quantum secret sharing ;natural multipartite extension of the original EPR paradox

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

multipartite EPR steering for (e) Greenberger-Horne-Zeilinger and Gaussian continuous variable states in (eb) loophole-free scenarios

G_{44} : Category one of Greenberger-Horne-Zeilinger and Gaussian continuous variable states in (eb) loophole-free scenarios

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of multipartite EPR steering

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21},t)\;\; +(a''_{16})^{(2,2,2)}(T_{17},t)\;\; +(a''_{13})^{(1,1,1,)}(T_{14},t)\\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{40})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{44})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a'_{21})^{(3)}+(a''_{21})^{(3)}(T_{21},t)\;\; +(a''_{17})^{(2,2,2)}(T_{17},t)\;\; +(a''_{14})^{(1,1,1,)}(T_{14},t)\\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{41})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{45})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21},t)\;\; +(a''_{18})^{(2,2,2)}(T_{17},t)\;\; +(a''_{15})^{(1,1,1,)}(T_{14},t)\\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\;\; +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\;\; +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\;\; +(a''_{42})^{(8,8,8,8)}(T_{41},t)\;\; +(a''_{46})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\,-(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\,-(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\,-(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ \boxed{-(b_{24}'')^{(4)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{28}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{33}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{34}'')^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a_{28}')^{(5)}\boxed{+(a_{28}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a_{29}')^{(5)}\boxed{+(a_{29}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a_{30}')^{(5)}\boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

$Where\ \boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(IIIIII) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(JJJJJJ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(KKKKKK) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(LLLLLL)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(MMMMMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(NNNNNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(x) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad (378)$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad (379)$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad (380)$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- (381)

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 (382)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(x) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^{*}=\frac{(b_{28})^{(5)}T_{29}^{*}}{[(b_{28}')^{(5)}-(b_{28}'')^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*}=\frac{(b_{30})^{(5)}T_{29}^{*}}{[(b_{30}')^{(5)}-(b_{30}'')^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*})=0$, T_{33}^{*} given by $f(T_{33}^{*})=0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a_{32}')^{(6)}+(a_{32}'')^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a_{34}')^{(6)}+(a_{34}'')^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b_{32}')^{(6)}-(b_{32}'')^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b_{34}')^{(6)}-(b_{34}'')^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a_{36}')^{(7)}+(a_{36}'')^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a_{38}')^{(7)}+(a_{38}'')^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b_{36}')^{(7)}-(b_{36}'')^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b_{38}')^{(7)}-(b_{38}'')^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a_{40}')^{(8)}+(a_{40}'')^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a_{42}')^{(8)}+(a_{42}'')^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b_{40}')^{(8)}-(b_{40}'')^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b_{42}')^{(8)}-(b_{42}'')^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a_{44}')^{(9)}+(a_{44}'')^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a_{46}')^{(9)}+(a_{46}'')^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b_{44}')^{(9)}-(b_{44}'')^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b_{46}')^{(9)}-(b_{46}'')^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY ONE

Einstein-Podolsky-Rosen Steering And Multipartite Quantum Nonlocality

INTRODUCTION—VARIABLES USED

More information: Genuine Multipartite Einstein-Podolsky-Rosen Steering, Q.Y. He and M. D. Reid,

Phys. Rev. Lett. 111, 250403 (2013) link.aps.org/doi/10.1103/PhysRevLett.111.250403 Journal reference: Physical Review Letters Provided by Swinburne University of Technology

(1) Entanglement and nonlocality has been a central issue (eb) in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are (=) correlated quantum states which demonstrate (eb) an inconsistency between the completeness of quantum mechanics and (e&eb)local realism.
(2) Quantum mechanics and the concept of local realism. Schroedinger introduced the term 'steering' to describe (eb) this apparent nonlocality, and pointed out that these states necessarily involve (e&eb) entanglement, that is, they cannot (e) be separated into factorized terms.
(3) Later, Bell's work served to demonstrate that the situation is even more serious. Bell found quantum states whose correlations cannot (e) be explained by any possible local hidden variable (LHV) theory. Thus, Einstein's hope of a local realistic completion of quantum mechanics was not feasible

Unified criteria for multipartite quantum nonlocality Phys. Rev. A 84, 032115 – Published 19 September 2011 E. G. Cavalcanti

(4) Wiseman and co-workers [H. M. Wiseman, S. J. Jones, and A. C. Doherty, Phys. Rev. Lett. 98, 140402, (2007)] proposed a distinction among the nonlocality classes of Bell's nonlocality, Einstein-Podolsky-Rosen (EPR) paradox or steering, and entanglement based (e) on whether or not an overseer trusts each party in a bipartite scenario where they are asked to demonstrate entanglement.
(5) Here authors extend that concept to the multipartite case and derive inequalities that progressively test for (e) those classes of nonlocality, with different thresholds for each level.
(6) This framework includes the three classes of nonlocality above in special cases and introduces (eb) a family of others. DOI: http://dx.doi.org/10.1103/PhysRevA.84.032115

Criteria for generalized macroscopic and mesoscopic quantum coherence Phys. Rev. A 77, 062108 – Published 18 June 2008 E. G. Cavalcanti

(7) Authors consider macroscopic, mesoscopic, and "S-scopic" quantum superpositions of eigenstates of an observable and develop (eb) some signatures for their existence.
(8) They define the extent, or size S of a superposition, with respect to (e&eb) an observable $\hat{x}$, as being the range of outcomes of $\hat{x}$ predicted by (e) that superposition.
(9) Such superpositions are referred to as (=) generalized S-scopic superpositions to distinguish them from (e) the extreme superpositions that superpose (e&eb) only the two states that have (e) a difference S in their prediction for the observable. We also consider generalized S-scopic superpositions of coherent states.

NOTATION

Module One

Entanglement and nonlocality has been a central issue in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are correlated quantum states which demonstrate (eb) an inconsistency between the completeness of quantum mechanics and (e&eb)local realism

G_{13} : Category one of Entanglement and nonlocality has been a central issue (eb) in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of correlated quantum states which demonstrate (eb) an inconsistency between the completeness of quantum mechanics and (e&eb)local realism

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Entanglement and nonlocality has been a central issue in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are correlated quantum states which demonstrate an inconsistency between the completeness of quantum mechanics and (e&eb)local realism

G_{16} : Category one of Entanglement and nonlocality has been a central issue in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are correlated quantum states

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of inconsistency between the completeness of quantum mechanics and (e&eb)local realism

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Entanglement and nonlocality has been a central issue in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are correlated quantum states which demonstrate an inconsistency between the completeness of quantum mechanics and (e&eb)local realism

G_{20} : Category one of Entanglement and nonlocality has been a central issue in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are correlated quantum states which demonstrate an inconsistency between the completeness of quantum mechanics; local realism

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of local realism; Entanglement and nonlocality has been a central issue in quantum mechanics since the Einstein-Podolsky-Rosen (EPR) paradox which brought EPR paradox shows that there are correlated quantum states which demonstrate an inconsistency between the completeness of quantum mechanics

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Schroedinger introduced the term ‘steering’ to describe this apparent nonlocality, and pointed out that these

states necessarily involve (e&eb) entanglement, that is, they cannot (e) be separated into factorized terms

G_{24} : Category one of Schroedinger introduced the term 'steering'

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of apparent nonlocality, and pointed out that these states necessarily involve (e&eb) entanglement, that is, they cannot (e) be separated into factorized terms

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Schroedinger introduced the term 'steering' to describe this apparent nonlocality, and pointed out that these states necessarily involve entanglement, that is, they cannot (e) be separated into factorized terms

G_{28} : Category one of steering 'describes this apparent nonlocality, and pointed out that these states; entanglement, that is, they cannot (e) be separated into factorized terms

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of entanglement, that is, they cannot (e) be separated into factorized terms ;steering' describes this apparent nonlocality, and pointed out that these states

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

'steering' describes this apparent nonlocality, and pointed out that these states necessarily involve entanglement, that is, they cannot (e) be separated into factorized terms

G_{32} : Category one of separated into factorized terms

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of 'steering' describes this apparent nonlocality, and pointed out that these states necessarily involve entanglement

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Quantum states whose correlations cannot (e) be explained by any possible local hidden variable (LHV) theory. Thus, Einstein's hope of a local realistic completion of quantum mechanics was not feasible

G_{36} : Category one of possible local hidden variable (LHV) theory. Thus, Einstein's hope of a local realistic completion of quantum mechanics was not feasible

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Quantum states whose correlations

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Here authors extend that concept to the multipartite case and derive inequalities that progressively test for (e) those classes of nonlocality, with different thresholds for each level.

G_{40} : Category one of classes of nonlocality, with different thresholds for each level.

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of concept to the multipartite case and derive inequalities that progressively test

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Framework includes the three classes of nonlocality above in special cases and introduces (eb) a family of others. DOI: http://dx.doi.org/10.1103/PhysRevA.84.032115

G_{44} : Category one of three classes of nonlocality above in special cases

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of family of others.

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1,2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category $1,2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category $1,2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{c} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{c} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{c} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(OOOOOO) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(PPPPPP) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(QQQQQQ) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(RRRRRR) $\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(SSSSSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(TTTTTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{32})^{(6)},\ T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(y) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

208

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)}$$

209

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)}$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),\left((G_{23})^{(2)},(T_{23})^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

214

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25}\ \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t} \right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)} t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t} \right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0<\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(v_1)^{(7)}<(\bar{v}_1)^{(7)}$

$$v^{(7)}(t)\geq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(v_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(v_2)^{(7)}}}$$

it follows $(v_0)^{(7)}\leq v^{(7)}(t)\leq(v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{v}_1)^{(7)}-(v_0)^{(7)}}{(v_0)^{(7)}-(\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)}\leq v^{(7)}(t)\leq(\bar{v}_1)^{(7)}$

If $0<(v_1)^{(7)}<(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)}\leq\frac{(v_1)^{(7)}+(C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)}-(v_2)^{(7)}\right)t\right]}}\leq v^{(7)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(\bar{v}_1)^{(7)}$$

If $0<(v_1)^{(7)}\leq(\bar{v}_1)^{(7)}\leq\boxed{(v_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(v_1)^{(7)}\leq v^{(7)}(t)\leq\frac{(\bar{v}_1)^{(7)}+(\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)}-(\bar{v}_2)^{(7)}\right)t\right]}}\leq(v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)}\leq v^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{v^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)}+(C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)}-(v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)}+(\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)}-(\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(y) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b_i'')^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY ONE A

S-Scopic Superpositions

INTRODUCTION—VARIABLES USED

Criteria for generalized macroscopic and mesoscopic quantum coherence Phys. Rev. A 77, 062108 – Published 18 June 2008 E. G. Cavalcanti

(1) Such superpositions are referred to as (=) generalized **S-scopic superpositions** to distinguish them from (e) the extreme superpositions that superpose (e&eb) only the two states that have (e) a difference S in their prediction for the observable. We also consider generalized S-scopic superpositions of coherent states.
(2) Authors explore the constraints that are placed on the statistics if we suppose a system to be described by (e) mixtures of superpositions that are restricted in size.
(3) In this way we arrive at experimental criteria that are sufficient to deduce (eb) the existence of a generalized S-scopic superposition.
(4) The signatures developed are useful where one is able to demonstrate (eb) a degree of squeezing.
(5) They also discuss how the signatures enable (eb) a new type of Einstein-Podolsky-Rosengedankenexperiment.DOI: ttp://dx.doi.org/10.1103/PhysRevA.77.062108

Multipartite quantum nonlocality under local decoherence Phys. Rev. A 86, 012108 – Published 16 July 2012 R. Chaves

(6) Authors study the nonlocal properties of two-qubit maximally entangled and (e&eb) N-qubit Greenberger-Horne-Zeilinger states under local decoherence.
(7) They show that the (non)resilience of entanglement under local depolarization or dephasing is (=) not necessarily equivalent to the (non)resilience of Bell-inequality violations.
(8) Apart from entanglement and Bell-inequality violations, they consider also nonlocality as quantified by (e) the nonlocal content of correlations and provide (eb) several examples of anomalous behaviors, both in the bipartite and multipartite cases.
(9) Apart from entanglement and Bell-inequality violations, they consider also nonlocality as quantified by (e) the nonlocal content of correlations and provide (eb) several examples of anomalous behaviors, both in the bipartite and multipartite cases.

NOTATION

Module One

Such superpositions are referred to as (=) generalized **S-scopic superpositions** are different from the extreme superpositions that superpose (e&eb) only the two states that have (e) a difference S in their prediction for the observable. We also consider generalized S-scopic superpositions of coherent states.

G_{13} : Category one of generalized **S-scopic superpositions** is different from the extreme superpositions; two states that have (e) a difference S in their prediction for the observable.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of two states that have (e) a difference S in their prediction for the observable.; generalized **S-scopic superpositions** are different from the extreme superpositions

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Authors explore the constraints that are placed on the statistics if we suppose a system to be described by (e) mixtures of superpositions that are restricted in size

G_{16} : Category one of mixtures of superpositions that are restricted in size

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of constraints that are placed on the statistics

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

In this way we arrive at experimental criteria that are sufficient to deduce (eb) the existence of a generalized S-scopic superposition

G_{20} : Category one of generalized S-scopic superposition

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of existence

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

signatures developed are useful where one is able to demonstrate (eb) a degree of squeezing

G_{24} : Category one of signatures developed are useful

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of degree of squeezing

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Signatures enable (eb) a new type of Einstein-Podolsky-Rosen gedanken experiment.

G_{28} : Category one of Signatures

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of new type of Einstein-Podolsky-Rosen gedanken experiment

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Authors study the nonlocal properties of two-qubit maximally entangled and (e&eb) N-qubit Greenberger-Horne-Zeilinger states under local decoherence

G_{32} : Category one of nonlocal properties of two-qubit maximally entangled; N-qubit Greenberger-Horne-Zeilinger states under local decoherence

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of N-qubit Greenberger-Horne-Zeilinger states under local decoherence ;nonlocal properties of two-qubit maximally entangled

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

They show that the (non)resilience of entanglement under local depolarization or dephasing is (=) not necessarily equivalent to the (non)resilience of Bell-inequality violations

G_{36} : Category one of(non)resilience of entanglement under local depolarization or dephasing

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of not necessarily equivalent to the (non)resilience of Bell-inequality violations

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Apart from entanglement and Bell-inequality violations, they consider also nonlocality as quantified by the nonlocal content of correlations and provide (eb) several examples of anomalous behaviors, both in the bipartite and multipartite cases.

G_{40} : Category one of nonlocality

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of quantified by the nonlocal content of correlations and provide (eb) several examples of anomalous behaviors, both in the bipartite and multipartite cases.

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Apart from entanglement and Bell-inequality violations, they consider also nonlocality as quantified by (e) the nonlocal content of correlations and provide (eb) several examples of anomalous behaviors, both in the bipartite and multipartite cases

G_{44} : Category one of nonlocal content of correlations and provide (eb) several examples of anomalous behaviors, both in the bipartite and multipartite cases

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Apart from entanglement and Bell-inequality violations, they consider also nonlocality as quantified

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**
$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)} + (a_{16}'')^{(2)}(T_{17},t) + (a_{13}'')^{(1,1,)}(T_{14},t) + (a_{20}'')^{(3,3,3)}(T_{21},t) \\ + (a_{24}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{28}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{36}'')^{(7,7,7)}(T_{37},t) + (a_{40}'')^{(8,8,8)}(T_{41},t) + (a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)} + (a_{17}'')^{(2)}(T_{17},t) + (a_{14}'')^{(1,1,)}(T_{14},t) + (a_{21}'')^{(3,3,3)}(T_{21},t) \\ + (a_{25}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{29}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{37}'')^{(7,7,7)}(T_{37},t) + (a_{41}'')^{(8,8,8)}(T_{41},t) + (a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)} + (a_{18}'')^{(2)}(T_{17},t) + (a_{15}'')^{(1,1,)}(T_{14},t) + (a_{22}'')^{(3,3,3)}(T_{21},t) \\ + (a_{26}'')^{(4,4,4,4,4)}(T_{25},t) + (a_{30}'')^{(5,5,5,5,5)}(T_{29},t) + (a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ + (a_{38}'')^{(7,7,7)}(T_{37},t) + (a_{42}'')^{(8,8,8)}(T_{41},t) + (a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for*

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for

category 1,2, and 3

$-(b_{16}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(UUUUUU) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(VVVVVV) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(WWWWWW) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(XXXXXX)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(YYYYYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(ZZZZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\quad ,\quad \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\quad ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(z) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14}\ \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14}\ \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \ \frac{1}{2}$ it results

$T_{14}\ \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\widehat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{36})^{(7)} \tag{256}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{36})^{(7)}\right]\leq(\widehat{Q}_{36})^{(7)} \tag{257}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric (258)

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \tag{259}$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t} \right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)} t} \right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t} \right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(z) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY ONE B

Multipartite Quantum Nonlocality

INTRODUCTION—VARIABLES USED

Multipartite quantum nonlocality under local decoherence Phys. Rev. A 86, 012108 – Published 16 July 2012 R. Chaves

(1) In addition, they study the practical implications of these anomalies on (eb) the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems. DOI: http://dx.doi.org/10.1103/PhysRevA.86.012108
(2) Practical implications of these anomalies on (eb) the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems.

Testing for Multipartite quantum nonlocality under local decoherence Phys. Rev. A 86, 012108 – Published 16 July 2012 R. Chaves

(1) In addition, they study the practical implications of these anomalies on (eb) the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems. DOI: http://dx.doi.org/10.1103/PhysRevA.86.012108
(2) Practical implications of these anomalies on (eb) the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems.

Testing for Multipartite Quantum Nonlocality Using Functional Bell Inequalities Phys. Rev. Lett. 103, 180402 – Published 29 October 2009 Q. Y. He, E. G. Cavalcanti et al

(3) Authors show that arbitrary functions of continuous variables, e.g., position and momentum, can be used (e) to generate tests that distinguish quantum theory from local hidden variable theories.
(4) By optimizing these functions, we obtain (eb) more robust violations of local causality than obtained previously.

(5) Authors analytically calculate the optimal function and include (e) the effect of nonideal detectors and noise, revealing that optimized functional inequalities are (=) resistant to standard forms of decoherence.

(6) Authors analytically calculate the optimal function and include (e) the effect of nonideal detectors and noise, revealing that optimized functional inequalities are (=) resistant to standard forms of decoherence.

(7) These inequalities could allow (eb) a loophole-free Bell test with efficient homodyne detection. DOI: http://dx.doi.org/10.1103/PhysRevLett.103.180402

Quantum Nonlocality and Partial Transposition for Continuous-Variable Systems Phys. Rev. Lett 101, 040404 – Published 25 July 2008 Alejo Salles et al

(8) A continuous-variable Bell inequality, valid for (e) an arbitrary number of observers measuring (eb) observables with an arbitrary number of outcomes, was recently introduced [Cavalcanti et al., Phys. Rev. Lett. 99, 210405 (2007)].

(9) They prove that any n-mode quantum state violating this inequality with quadrature measurements necessarily has (e) a negative partial transposition.

NOTATION

Module One

In addition, they study the practical implications of these anomalies on the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems. DOI: http://dx.doi.org/10.1103/PhysRevA.86.012108

G_{13} : Category one of practical implications of these anomalies; practical implications of these anomalies on the usefulness of noisy Greenberger-Horne-Zeilinger states

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems; resources for nonlocality-based physical protocols given by communication complexity problems.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Practical implications of these anomalies on the usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems

G_{16} : Category one of Practical implications of these anomalies; Greenberger-Horne-Zeilinger states

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of usefulness of noisy Greenberger-Horne-Zeilinger states as (=) resources for nonlocality-based physical protocols given by communication complexity problems; resources for nonlocality-based physical protocols given by communication complexity problems

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Arbitrary functions of continuous variables, e.g., position and momentum can be used (e) to generate tests that distinguish quantum theory from local hidden variable theories.

G_{20} : Category one of generate tests that distinguish quantum theory from local hidden variable theories.

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Arbitrary functions of continuous variables, e.g., position and momentum c

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

By optimizing these functions, we obtain (eb) more robust violations of local causality than obtained previously

G_{24} : Category one of optimisation of these functions

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of robust violations of local causality than obtained previously

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Authors analytically calculate the optimal function and include (e) the effect of nonideal detectors and noise, revealing that optimized functional inequalities are (=) resistant to standard forms of decoherence.

G_{28} : Category one of optimal function and include (e) the effect of nonideal detectors and noise, revealing that optimized functional inequalities

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of resistant to standard forms of decoherence

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Authors analytically calculate the optimal function and include (e) the effect of nonideal detectors and noise, revealing that optimized functional inequalities are resistant to standard forms of decoherence.

G_{32} : Category one of effect of nonideal detectors and noise, revealing that optimized functional inequalities are resistant to standard forms of decoherence

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of optimal function calculation

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

These inequalities could allow (eb) a loophole-free Bell test with efficient homodyne detection

G_{36} : Category one of These inequalities

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of loophole-free Bell test with efficient homodyne detection

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Continuous-variable Bell inequality, valid for an arbitrary number of observers measuring (eb) observables with an arbitrary number of outcomes, was recently introduced [Cavalcanti et al., Phys. Rev. Lett. 99, 210405 (2007)].

G_{40} : Category one of arbitrary number of observers measuring (eb) observables with an arbitrary number of outcomes, was recently introduced [Cavalcanti et al., Phys. Rev. Lett. 99, 210405 (2007)].

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Continuous-variable Bell inequality

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Continuous-variable Bell inequality, valid for an arbitrary number of observers measuring (eb) observables with an arbitrary number of outcomes, was recently introduced [Cavalcanti et al., Phys. Rev. Lett. 99,

210405 (2007)].

G_{44} : Category one of Continuous-variable Bell inequality, valid for an arbitrary number of observers

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of observables with an arbitrary number of outcomes, was recently introduced [Cavalcanti et al., Phys. Rev. Lett. 99, 210405 (2007)].

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18}$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a_{44}'')^{(9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b_{16}')^{(2)} & -(b_{16}'')^{(2)}(G_{19},t) & -(b_{13}'')^{(1,1,)}(G,t) & -(b_{20}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{36}'')^{(7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b_{17}')^{(2)} & -(b_{17}'')^{(2)}(G_{19},t) & -(b_{14}'')^{(1,1,)}(G,t) & -(b_{21}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{37}'')^{(7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b_{18}')^{(2)} & -(b_{18}'')^{(2)}(G_{19},t) & -(b_{15}'')^{(1,1,)}(G,t) & -(b_{22}'')^{(3,3,3,)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6)}(G_{35},t) \\ -(b_{38}'')^{(7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b_{16}'')^{(2)}(G_{19},t)$, $-(b_{17}'')^{(2)}(G_{19},t)$, $-(b_{18}'')^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt}=(a_{20})^{(3)}G_{21}-\left[\begin{array}{ccc}(a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt}=(a_{21})^{(3)}G_{20}-\left[\begin{array}{ccc}(a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt}=(a_{22})^{(3)}G_{21}-\left[\begin{array}{ccc}(a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}\\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}\end{array}\right]G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt}=(b_{20})^{(3)}T_{21}-\left[\begin{array}{ccc}(b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)}\\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}\end{array}\right]T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1,2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1,2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1,2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt}=(a_{29})^{(5)}G_{28}-\left[\begin{array}{ccc}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt}=(a_{30})^{(5)}G_{29}-\left[\begin{array}{ccc}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{array}\right]G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34} \quad (87)$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34} \quad 90$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition

coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation

coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)} \boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} \boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(AAAAAAA) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(BBBBBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(CCCCCCC) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(DDDDDDD)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(EEEEEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(FFFFFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{16})^{(2)}$, $T_i^0 \leq (\hat{Q}_{16})^{(2)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\Big] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(aa) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq-(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq-(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(aa) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO

Entanglement Distillation

INTRODUCTION—VARIABLES USED

Reviving Einstein's spooky action at a distance Mar 18, 2014

(1) Physicists at The University of Queensland and the Australian National University (ANU) have demonstrated a software-based quantum amplifier which has the potential to expand (eb) the use of ultra-secure quantum cryptographic communications.
(2) The development could lead to (eb) new ways to guarantee security of everyday communications, including financial transactions and email.
(3) ANU Research School of Physics and Engineering PhD student Helen Chranowski said this had the potential to (e) guarantee security for any kind of information you want to transmit, from particularly sensitive information, like banking data or electoral results, to personal emails.
(4) Quantum cryptography exploits (e) quantum mechanics to guarantee 100 per cent security between a sender and receiver of information.
(5) Most forms of quantum communication also use (e) a phenomenon known as quantum entanglement, which Albert Einstein described as "spooky action at a distance".
(6) Although powerful, this phenomenon is fragile and is often destroyed (e) before it can be transmitted over large distances.
(7) The usual approach to combating loss - an amplifier – such as those used for radio or television signals, only makes (eb) the situation worse. Researchers, led by Professor Ping Koy Lam at the ANU Research School of Physics and Engineering, have used a sophisticated scheme to work out when an amplified signal works or fails.
(8) They have developed an algorithm which allows (eb) them to verify whether a quantum entanglement signal has been successfully received.
(9) The software allows the physicists to filter out (e&eb) unwanted signal failures.

NOTATION

Module One

Physicists at The University of Queensland and the Australian National University (ANU) have demonstrated a software-based quantum amplifier which has the potential to expand (eb) the use of ultra-secure quantum cryptographic communications

G_{13} : Category one of software-based quantum amplifier which has the potential

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of use of ultra-secure quantum cryptographic communications

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Development could lead to (eb) new ways to guarantee security of everyday communications, including financial transactions and email.

G_{16} : Category one of Development of software-based quantum amplifier which has the potential to expand the use of ultra-secure quantum cryptographic communications

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of new ways to guarantee security of everyday communications, including financial transactions and email

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

G_{20} : Category one of guarantee security for any kind of information you want to transmit, from particularly sensitive information, like banking data or electoral results, to personal emails

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of potential to software-based quantum amplifier which has the potential to expand the use of ultra-secure quantum cryptographic communications

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Quantum cryptography exploits (e) quantum mechanics to guarantee 100 per cent security between a sender and receiver of information.

G_{24} : Category one of quantum mechanics to guarantee 100 per cent security between a sender and receiver of information.

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Quantum cryptography

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Most forms of quantum communication also use (e) a phenomenon known as quantum entanglement, which Albert Einstein described as "spooky action at a distance".

G_{28} : Category one of phenomenon known as quantum entanglement, which Albert Einstein described as "spooky action at a distance".

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Most forms of quantum communication

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Phenomenon albeit powerful is fragile and is often destroyed (e) before it can be transmitted over large distances

G_{32} : Category one of transmitted over large distances

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Phenomenon albeit powerful is fragile

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

usual approach to combating loss - an amplifier – such as those used for radio or television signals, only makes (eb) the situation worse

G_{36} : Category one of usual approach to combating loss - an amplifier – such as those used for radio or television signals

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of situation worse

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

algorithm allows (eb) them to verify whether a quantum entanglement signal has been successfully received

G_{40} : Category one of algorithm

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of verify whether a quantum entanglement signal has been successfully received

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

software allows the physicists to filter out (e&eb) unwanted signal failures

G_{44} : Category one of software; unwanted signal failures

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of unwanted signal failures; software

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition

coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{20})^{(3)}\left[+(a''_{20})^{(3)}(T_{21},t)\right]\left[+(a''_{16})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{13})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{40})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{44})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c} (a'_{21})^{(3)}\left[+(a''_{21})^{(3)}(T_{21},t)\right]\left[+(a''_{17})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{14})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{37})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{41})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{45})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c} (a'_{22})^{(3)}\left[+(a''_{22})^{(3)}(T_{21},t)\right]\left[+(a''_{18})^{(2,2,2)}(T_{17},t)\right]\left[+(a''_{15})^{(1,1,1,)}(T_{14},t)\right] \\ \left[+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\right]\left[+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\right]\left[+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{38})^{(7,7,7,7)}(T_{37},t)\right]\left[+(a''_{42})^{(8,8,8,8)}(T_{41},t)\right]\left[+(a''_{46})^{(9,9,9)}(T_{45},t)\right] \end{array}\right] G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for

category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

are fourth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & \boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & \boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & \boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a_{40}'')^{(8,8\,,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) \\ & -(b''_{32})^{(6,6,6)}(G_{35},t) & \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & \\ & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & \\ & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) & \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & \\ & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & \\ & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) & \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) \\ & -(b''_{33})^{(6,6,6)}(G_{35},t) & \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & \\ & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & \\ & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) & \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & \\ & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & \\ & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) & \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) \\ & -(b''_{34})^{(6,6,6)}(G_{35},t) & \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & \\ & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & \\ & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) & \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & \\ & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & \\ & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) & \end{array}\right] T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) \\ & +(a''_{24})^{(4,4,4,)}(T_{25},t) & \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & \\ & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & \\ & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) & \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & \\ & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & \\ & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) & \end{array}\right] G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) \\ & +(a''_{25})^{(4,4,4,)}(T_{25},t) & \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & \\ & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & \\ & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) & \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & \\ & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & \\ & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) & \end{array}\right] G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{40}}{dt} =$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)} G_{44}$$
$$- \left[\begin{array}{c} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)} G_{45}$$
$$- \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)} T_{45} -$$
$$\left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, \ (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$\lim_{G\to\infty}(b_i'')^{(3)}\,(G_{23},t) = \ (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$

$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and \ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$

$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$

$(b_i'')^{(4)}\big((G_{27}),t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}),t\big) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}\,(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}\,(G_{31}, t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ \ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(GGGGGGG) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(HHHHHHH) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq \ (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(IIIIIII) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(JJJJJJJ)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(KKKKKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(LLLLLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and \ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}$, $(r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}\left(G_{19}(s_{(16)}), s_{(16)}\right)\right) T_{18}(s_{(16)})\Big] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''^{(3)}_{20}\right)\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{20}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{21}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}\left(T_{21}(s_{(20)}), s_{(20)}\right)\right) G_{22}(s_{(20)})\Big] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)},\ T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)},\ T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{36})^{(7)}$, $T_i^0 \leq (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(bb) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28}, G_{30} and G_{28}, G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30}. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b'_i)^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)},\ if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, if\ (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$

and $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)},\ if\ (\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)}$, ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(bb) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO A

Distillation Protocols Are Necessarily Non-Deterministic

INTRODUCTION—VARIABLES USED

Reviving Einstein's spooky action at a distance Mar 18, 2014

(1) It is now possible to use (e) an amplifier to distribute (e&eb) the signals over greater distances, and for the sender and receiver to know if the signal has arrived safely.
(2) Professor Lam said the results showed clear success in resurrecting (eb) the quantum effect that so perturbed Einstein.
(3) UQ's Professor Timothy Ralph said the results would help harness (e) the power of quantum entanglement in communications.
(4) "If we want to use entanglement over larger distances, and harness its power for telecommunication, we need (e) quantum amplifiers that allow us (eb) to resurrect quantum entanglement in a communication line," he said. The results have been published in the latest on-line issue of Nature Photonics.

More information: Measurement-based noiseless linear amplification for quantum communication, Nature Photonics (2014) DOI: 10.1038/nphoton.2014.49 Journal reference: Nature Photonics Provided by University of Queensland

NATURE PHOTONICS | ARTICLE Measurement-based noiseless linear amplification for quantum communication Helen M. Chrzanowski et al Nature Photonics (2014) doi:10.1038/nphoton.2014.49

(5) Entanglement distillation is (=) an indispensable ingredient in extended quantum communication networks.
(6) Distillation protocols are (=) necessarily non-deterministic and require (e) advanced experimental techniques such as noiseless amplification.
(7) Recently, it was shown that the benefits of noiseless amplification could be extracted by (e) performing a post-selective filtering (e&eb) of the measurement record to improve (eb) the performance of quantum key distribution.
(8) Authors apply this protocol to entanglement degraded by (e) transmission loss of up to the equivalent of 100 km of optical fibre.
(9) Authors measure an effective entangled resource stronger than that achievable by even a maximally entangled resource passively transmitted through (e&eb) the same channel.

NOTATION

Module One

It is now possible to use an amplifier to distribute the signals over greater distances, and for the sender and receiver to know if the signal has arrived safely

G_{13} : Category one of distribute the signals over greater distances, and for the sender and receiver to know if the signal has arrived safely

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of amplifier

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Results showed clear success in resurrecting (eb) the quantum effect that so perturbed Einstein.

G_{16} : Category one of Results

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of resurrection quantum effect

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

UQ's Professor Timothy Ralph said the results would help harness (e) the power of quantum entanglement in communications

G_{20} : Category one of power of quantum entanglement in communications

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of results would help

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

"If we want to use entanglement over larger distances, and harness its power for telecommunication, we need (e) quantum amplifiers that allow us (eb) to resurrect quantum entanglement in a communication line," he said. The results have been published in the latest on-line issue of Nature Photonics

G_{24} : Category one of quantum amplifiers that allow us (eb) to resurrect quantum entanglement in a communication line

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of use entanglement over larger distances, and harness its power for telecommunication

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

quantum amplifiers that allow us (eb) to resurrect quantum entanglement in a communication line

G_{28} : Category one of quantum amplifiers

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of resurrect quantum entanglement in a communication line

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Measurement of an effective entangled resource stronger than that achievable by even a maximally entangled resource passively transmitted through (e&eb) the same channel

G_{32} : Category one of Measurement of an effective entangled resource stronger than that achievable by even a maximally entangled resource passively transmitted; same channel

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of same channel ;Measurement of an effective entangled resource stronger than that achievable by even a maximally entangled resource passively transmitted

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Distillation protocols are necessarily non-deterministic and require (e) advanced experimental techniques such as noiseless amplification

G_{36} : Category one of non-deterministic and require (e) advanced experimental techniques such as noiseless amplification

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Distillation protocols

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Distillation protocols are necessarily non-deterministic and require (e) advanced experimental techniques such as noiseless amplification

G_{40} : Category one of advanced experimental techniques such as noiseless amplification

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Distillation protocols are necessarily non-deterministic

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Authors apply this protocol to entanglement degraded by (e) transmission loss of up to the equivalent of 100 km of optical fibre

G_{44} : Category one of transmission loss of up to the equivalent of 100 km of optical fibre

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of application of protocol to entanglement degraded

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & \boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} & \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & \boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} & \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & \boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & \boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} & \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$
$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\widehat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\widehat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)},$ are positive constants 109

$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\widehat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\widehat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(MMMMMMM) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(NNNNNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

(OOOOOOO) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(PPPPPPP)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(QQQQQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(RRRRRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40,41,42$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(cc) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0$ for $\mathrm{t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)},\ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)},\ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(cc) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO B

Ease Of Implementation And Near-Optimal Probability Of Success

INTRODUCTION—VARIABLES USED

NATURE PHOTONICS | ARTICLE Measurement-based noiseless linear amplification for quantum communication Helen M. Chrzanowski et al Nature Photonics (2014) doi:10.1038/nphoton.2014.49

(1) They also provide a proof-of-principle demonstration of secret key extraction from (e) an otherwise insecure regime.
(2) The measurement-based (e) noiseless linear amplifier offers two advantages over its physical counterpart: ease of implementation and near-optimal probability of success.
(3) It should provide an effective and versatile tool for (e) a broad class of entanglement-based quantum communication protocols.

Entanglement distillation (Wikipedia)

(4) Entanglement distillation is the transformation of N copies of an arbitrary entangled state \rho into (eb) some number of approximately pure Bell pairs, using (e) only local operations and classical communication (LOCC).
(5) Entanglement distillation can in this way overcome (e) the degenerative influence of noisy quantum channels by transforming previously shared less entangled pairs into (eb) a smaller number of maximally entangled pairs (Bell states).

Quantifying Entanglement

(6) A two qubit system can be written as (=) a superposition of possible computational basis qubit states: $|00\rangle, |01\rangle, |10\rangle, |11\rangle$, each with an associated complex coefficient α: $|\psi\rangle = \alpha_{00}|00\rangle + \alpha_{01}|01\rangle + \alpha_{10}|10\rangle + \alpha_{11}|11\rangle$
(7) As in the case of a single qubit, the probability of measuring a particular computational basis state $|x\rangle$ is (=) the amplitude of its associated coefficient $|\alpha_x|^2$, subject to the normalization condition $\sum_{x\epsilon 0,1} |\alpha_x|^2 = 1$.
(8) The Bell state is a particularly important example of a two qubit state: $\frac{|00\rangle + |11\rangle}{\sqrt{2}}$
(9) Bell states possess the amazing property that measurement outcomes of a Bell state are (=) correlated.

NOTATION

Module One

They also provide a proof-of-principle demonstration of secret key extraction from (e) an otherwise insecure regime.

G_{13} : Category one of **otherwise** insecure regime.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of proof-of-principle demonstration of secret key extraction

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

measurement-based noiseless linear amplifier offers two advantages over its physical counterpart: ease of implementation and near-optimal probability of success

G_{16} : Category one of noiseless linear amplifier offers two advantages over its physical counterpart: ease of implementation and near-optimal probability of success

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of measurement-based

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

G_{20} : Category one of measurement-based noiseless linear amplifier

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of near-optimal probability of success

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

G_{24} : Category one of measurement-based noiseless linear amplifier

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of ease of implementation

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Entanglement distillation is the transformation of N copies of an arbitrary entangled state \rho into some number of approximately pure Bell pairs, using (e) only local operations and classical communication (LOCC).

G_{28} : Category one of **Entanglement distillation is** the transformation of N copies of an arbitrary entangled state \rho

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of some number of approximately pure Bell pairs, using (e) only local operations and classical communication (LOCC).

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

A two qubit system can be written as (=) a superposition of possible computational basis qubit states: $|00\rangle, |01\rangle, |10\rangle, |11\rangle$, each with an associated complex coefficient α:

$$|\psi\rangle = \alpha_{00}|00\rangle + \alpha_{01}|01\rangle + \alpha_{10}|10\rangle + \alpha_{11}|11\rangle$$

G_{32} : Category one of LHs of $|\psi\rangle = \alpha_{00}|00\rangle + \alpha_{01}|01\rangle + \alpha_{10}|10\rangle + \alpha_{11}|11\rangle$

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of RHS terms of $|\psi\rangle = \alpha_{00}|00\rangle + \alpha_{01}|01\rangle + \alpha_{10}|10\rangle + \alpha_{11}|11\rangle$

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

As in the case of a single qubit, the probability of measuring a particular computational basis state $|x\rangle$ is (=) the amplitude of its associated coefficient $|\alpha_x|^2$, subject to the normalization condition $\sum_{x\epsilon 0,1} |\alpha_x|^2 = 1$.

G_{36} : Category one of probability of measuring a particular computational basis state $|x\rangle$

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of amplitude of its associated coefficient $|\alpha_x|^2$, subject to the normalization condition $\sum_{x\epsilon 0,1} |\alpha_x|^2 = 1$.

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Bell state is a particularly important example of a two qubit state: $\frac{|00\rangle + |11\rangle}{\sqrt{2}}$

G_{40} : Category one of Bell state is a particularly important example of a two qubit state

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of $\frac{|00\rangle + |11\rangle}{\sqrt{2}}$. There are two options. Either take the characterstics of the systems that satisfy the conditionalities of the given proposition or alternatively take all the three categories equivalent.

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Bell states possess the amazing property that measurement outcomes of a Bell state are (=) correlated

G_{44} : Category one of Bell states possess the amazing property that measurement outcomes of a Bell state

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of correlated

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$

$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{c} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{c} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt}=(b_{17})^{(2)}T_{16}-\begin{bmatrix}(b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t)\\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t)\end{bmatrix}T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(SSSSSSS) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(TTTTTTT) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(UUUUUUU) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(VVVVVVV)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(WWWWWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(XXXXXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\left((G_{43}), t\right) = (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\left((G_{43}), t\right)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(dd) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}} \left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad (333)$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad (334)$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad (335)$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad (336)$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(dd) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO C

Bell States Can Be Used To Quantify Entanglement

INTRODUCTION—VARIABLES USED

Entanglement distillation (Wikipedia)

(1) As can be seen from the expression above, the two possible measurement outcomes are (=) zero and one, both with probability of 50%.
(2) As a result, a measurement of the second qubit always gives (eb) the same result as the measurement of the first qubit.
(3) **Bell states can be used (e) to quantify entanglement**.
(4) Let m be the number of high-fidelity copies of a Bell state that can be produced using (e) LOCC.
(5) Given a large number of Bell states the amount of entanglement present in a pure state $|\psi\rangle$ can then be defined as (=) the ratio of n/m, called the distillable entanglement of a particular state $|\phi\rangle$, which gives (eb) a quantified measure of the amount of entanglement present in a given system.
(6) The process of entanglement distillation aims (eb) to saturate this limiting ratio.
(7) The number of copies of a pure state that may be converted into a maximally entangled state is equal to (=) the von Neumann entropy S(p) of the state, which is an extension of the concept of classical entropy for quantum systems.
(8) Mathematically, for a given density matrix p, the von Neumann entropy S(p) is $S(p) = -\mathrm{Tr}(p \ln p)$.
(9) Entanglement can then be quantified as the entropy of entanglement, which is the von Neumann entropy of either p_A or p_B as: $E = -\mathrm{Tr}(p_A \ln p_A) = -\mathrm{Tr}(p_B \ln p_B)$, This ranges from 0 for a product state to 1 for a maximally entangled state

NOTATION

Module One

As can be seen from the expression above, the two possible measurement outcomes are (=) zero and one, both with probability of 50%.

G_{13} : Category one of two possible measurement outcomes

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of zero and one, both with probability of 50%.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

As a result, a measurement of the second qubit always gives (eb) the same result as the measurement of the first qubit

G_{16} : Category one of measurement of the second qubit

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of same result as the measurement of the first qubit

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Bell states can be used (e) to quantify entanglement

G_{20} : Category one of **quantify entanglement**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **Bell states**

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Let m be the number of high-fidelity copies of a Bell state that can be produced using (e) LOCC

G_{24} : Category one of LOCC

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of m be the number of high-fidelity copies of a Bell state that can be produced

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Given a large number of Bell states the amount of entanglement present in a pure state $|\psi\rangle$ can then be defined as the ratio of n/m, called the distillable entanglement of a particular state $|\phi\rangle$, which gives (eb) a quantified measure of the amount of entanglement present in a given system.

G_{28} : Category one of Given a large number of Bell states the amount of entanglement present in a pure state $|\psi\rangle$

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of ratio of n/m, called the distillable entanglement of a particular state $|\phi\rangle$, which gives (eb) a quantified measure of the amount of entanglement present in a given system.

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Given a large number of Bell states the amount of entanglement present in a pure state $|\psi\rangle$ can then be defined as the ratio of n/m, called the distillable entanglement of a particular state $|\phi\rangle$, which gives (eb) a quantified measure of the amount of entanglement present in a given system

G_{32} : Category one of Given a large number of Bell states the amount of entanglement present in a pure state $|\psi\rangle$ can then be defined as the ratio of n/m, called the distillable entanglement of a particular state $|\phi\rangle$,

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of quantified measure of the amount of entanglement present in a given system

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

The process of entanglement distillation aims (eb) to saturate this limiting ratio

G_{36} : Category one of process of entanglement distillation

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of saturate this limiting ratio

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Number of copies of a pure state that may be converted into a maximally entangled state is equal to (=) the von Neumann entropy S(p) of the state, which is an extension of the concept of classical entropy for quantum systems.

G_{40} : Category one of Number of copies of a pure state that may be converted into a maximally entangled state

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of von Neumann entropy S (p) of the state, which is an extension of the concept of classical entropy for quantum systems.

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Entanglement can then be quantified as the entropy of entanglement, which is the von Neumann entropy of either p_A or p_B as: $E = -\mathrm{Tr}(p_A \ln p_A) = -\mathrm{Tr}(p_B \ln p_B)$, This ranges from 0 for a product state to 1 for a maximally entangled state

G_{44} : Category one of LHS of the equation $E = -\mathrm{Tr}(p_A \ln p_A) = -\mathrm{Tr}(p_B \ln p_B),$

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of RHS of the equation $E = -\mathrm{Tr}(p_A \ln p_A) = -\mathrm{Tr}(p_B \ln p_B),$

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{20})^{(3)}\,+(a''_{20})^{(3)}(T_{21},t)\;+(a''_{16})^{(2,2,2)}(T_{17},t)\;+(a''_{13})^{(1,1,1,)}(T_{14},t)\\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\;+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\;+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\;+(a''_{40})^{(8,8,8,8)}(T_{41},t)\;+(a''_{44})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a'_{21})^{(3)}\,+(a''_{21})^{(3)}(T_{21},t)\;+(a''_{17})^{(2,2,2)}(T_{17},t)\;+(a''_{14})^{(1,1,1,)}(T_{14},t)\\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\;+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\;+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\;+(a''_{41})^{(8,8,8,8)}(T_{41},t)\;+(a''_{45})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{22})^{(3)}\,+(a''_{22})^{(3)}(T_{21},t)\;+(a''_{18})^{(2,2,2)}(T_{17},t)\;+(a''_{15})^{(1,1,1,)}(T_{14},t)\\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\;+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\;+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\;+(a''_{42})^{(8,8,8,8)}(T_{41},t)\;+(a''_{46})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\,-(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\,-(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\,-(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ \boxed{-(b_{24}'')^{(4)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{28}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{33}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{34}'')^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a_{28}')^{(5)}\boxed{+(a_{28}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a_{29}')^{(5)}\boxed{+(a_{29}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a_{30}')^{(5)}\boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

$Where\ \boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$-\begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(YYYYYYY) $\quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(ZZZZZZZ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(AAAAAAAA) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(BBBBBBBB)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CCCCCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DDDDDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40,41,42$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ee) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \le \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \le \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \ge (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \ge \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ee) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO D

Entanglement Concentration

INTRODUCTION—VARIABLES USED


Entanglement distillation (Wikipedia)

Motivation

(1) Suppose that two parties, Alice and Bob, would like to communicate (e&eb) classical information over a noisy quantum channel.
(2) Either classical or quantum information can be transmitted over a quantum channel by encoding (e) the information in a quantum state.
(3) With this knowledge, Alice encodes (e) the classical information that she intends to send to (e&eb) Bob in a (quantum) product state, as a tensor product of reduced density matrices $p_1 \otimes p_2 \otimes ....$ Where each p is diagonal and can only be used as a onetime input for a particular channel ϵ.
(4) The fidelity of the noisy quantum channel is (=) a measure of how closely the output of a quantum channel resembles the input, and is therefore a measure (eb) of how well a quantum channel preserves information.
(5) If a pure state ψ is sent into a quantum channel emerges as (=) the state represented by density matrix p, the **fidelity of transmission** is defined as $F = \langle \psi | p | \psi \rangle$.
(6) The problem that Alice and Bob now face is that quantum communication over large distances depends upon (=) successful distribution of highly entangled quantum states, and due (e) to unavoidable noise in quantum communication channels, the quality of entangled states generally decreases (e) exponentially with channel length as a function of the fidelity of the channel.
(7) Entanglement distillation addresses this problem of maintaining a high degree of entanglement between distributed quantum states by transforming N copies of an arbitrary entangled state ρ into (eb) approximately $S(\rho)N$ Bell pairs, using only local operations and classical communication.
(8) The objective is to share strongly correlated qubits between distant parties (Alice and Bob) in order to allow (eb) reliable quantum teleportation or quantum cryptography.

Entanglement Concentration

Pure States



The new fidelity after one iteration of the distillation protocol for pure states.

(9) Given n particles in the singlet state shared between Alice and Bob, local actions and classical communication will suffice to prepare (eb) m arbitrarily good copies of ϕ with a yield $\frac{m}{n}$ Approaching $\frac{1}{E(\phi)}$ as $n \to \infty$.

NOTATION

Module One

Suppose that two parties, Alice and Bob, would like to communicate (e&eb) classical information over a noisy quantum channel.

G_{13} : Category one of Alice and Bob, would like to communicate; classical information over a noisy quantum channel.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of classical information over a noisy quantum channel.; Alice and Bob, would like to communicate

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Either classical or quantum information can be transmitted over a quantum channel by encoding (e) the information in a quantum state

G_{16} : Category one of information in a quantum state

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of classical or quantum information can be transmitted over a quantum channel

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

With this knowledge, Alice encodes the classical information that she intends to send to (e&eb) Bob in a (quantum) product state, as a tensor product of reduced density matrices $p_1 \otimes p_2 \otimes$.... Where each p is diagonal and can only be used as a onetime input for a particular channel ϵ.

G_{20} : Category one of classical information that she intends to send to (e&eb) Bob in a (quantum) product state, as a tensor product of reduced density matrices $p_1 \otimes p_2 \otimes$.... Where each p is diagonal and can only be used as a onetime input for a particular channel ϵ.

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Alice

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Alice encodes the classical information that she intends to send to (e&eb) Bob in a (quantum) product state, as a tensor product of reduced density matrices $p_1 \otimes p_2 \otimes$.... Where each p is diagonal and can only be used as a onetime input for a particular channel ϵ.

G_{24} : Category one of Alice encodes the classical information that she intends to send; Bob in a (quantum) product state, as a tensor product of reduced density matrices $p_1 \otimes p_2 \otimes$.... Where each p is diagonal and can only be used as a onetime input for a particular channel ϵ.

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Bob in a (quantum) product state, as a tensor product of reduced density matrices $p_1 \otimes p_2 \otimes$.... Where each p is diagonal and can only be used as a onetime input for a particular channel ϵ.; Alice encodes the classical information that she intends to send

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Fidelity of the noisy quantum channel is a measure of how closely the output of a quantum channel resembles the input, and is therefore a measure (eb) of how well a quantum channel preserves information.

G_{28} : Category one of Fidelity of the noisy quantum channel;

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of measure of how closely the output of a quantum channel resembles the input, and is therefore a measure (eb) of how well a quantum channel preserves information

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

G_{32} : Category one of Fidelity of the noisy quantum channel is a measure of how closely the output of a quantum channel resembles the input

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of preservation of information by quantum channel (the quality thereof)

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

If a pure state ψ is sent into a quantum channel emerges as (=) the state represented by density matrix p, the **fidelity of transmission** is defined as $F = \langle\psi|p|\psi\rangle$.

G_{36} : Category one of pure state ψ is sent into a quantum channel emerges

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of state represented by density matrix p, the **fidelity of transmission** is defined as $F = \langle\psi|p|\psi\rangle$.

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Multiple modules: care

problem that Alice and Bob now face is that quantum communication over large distances depends upon successful distribution of highly entangled quantum states, and due to unavoidable noise in quantum communication channels, the quality of entangled states generally decreases (e) exponentially with channel length as a function of the fidelity of the channel

G_{40} : Category one of problem that Alice and Bob now face is that quantum communication over large distances; unavoidable noise in quantum communication channels, the quality of entangled states generally decreases (e) exponentially with channel length as a function of the fidelity of the channel; exponentially with channel length as a function of the fidelity of the channel

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of depends upon successful distribution of highly entangled quantum states, and due (e) to unavoidable noise in quantum communication channels, the quality of entangled states generally decreases (e) exponentially with channel length as a function of the fidelity of the channel; problem that Alice and Bob now face is that quantum communication over large distances depends upon successful distribution of highly entangled quantum states; problem that Alice and Bob now face is that quantum communication over large distances depends upon successful distribution of highly entangled quantum states, and due to unavoidable noise in quantum communication channels, the quality of entangled states

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Care: Multiple modules

Entanglement distillation addresses this problem of maintaining a high degree of entanglement between distributed quantum states by transforming N copies of an arbitrary entangled state ρ into (eb) approximately $S(\rho)N$ Bell pairs, using only local operations and classical communication

Objective is to share strongly correlated qubits between distant parties (Alice and Bob) in order to allow (eb) reliable quantum teleportation or quantum cryptography.

Given n particles in the singlet state shared between Alice and Bob, local actions and classical communication will suffice to prepare (eb) m arbitrarily good copies of ϕ with a yield $\frac{m}{n}$

Approaching $\frac{1}{E(\phi)}$ as $n \to \infty$.

G_{44} : Category one of Entanglement distillation addresses this problem of maintaining a high degree of entanglement between distributed quantum states by transforming N copies of an arbitrary entangled state ρ ; Objective is to share strongly correlated qubits between distant parties (Alice and Bob); n particles in the singlet state shared between Alice and Bob, local actions and classical communication

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of approximately $S(\rho)N$ Bell pairs, using only local operations and classical communication; reliable quantum teleportation or quantum cryptography;) m arbitrarily good copies of ϕ with a yield $\frac{m}{n}$ Approaching $\frac{1}{E(\phi)}$ as $n \to \infty$.

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}$ $(a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$ $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}&+(a''_{16})^{(2)}(T_{17},t)&+(a''_{13})^{(1,1,)}(T_{14},t)&+(a''_{20})^{(3,3,3)}(T_{21},t)\\+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{36})^{(7,7,7)}(T_{37},t)&+(a''_{40})^{(8,8,8)}(T_{41},t)&+(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)}&+(a''_{17})^{(2)}(T_{17},t)&+(a''_{14})^{(1,1,)}(T_{14},t)&+(a''_{21})^{(3,3,3)}(T_{21},t)\\+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{37})^{(7,7,7)}(T_{37},t)&+(a''_{41})^{(8,8,8)}(T_{41},t)&+(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)}&+(a''_{18})^{(2)}(T_{17},t)&+(a''_{15})^{(1,1,)}(T_{14},t)&+(a''_{22})^{(3,3,3)}(T_{21},t)\\+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)&+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)&+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\+(a''_{38})^{(7,7,7)}(T_{37},t)&+(a''_{42})^{(8,8,8)}(T_{41},t)&+(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)}&-(b''_{16})^{(2)}(G_{19},t)&-(b''_{13})^{(1,1,)}(G,t)&-(b''_{20})^{(3,3,3,)}(G_{23},t)\\-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)&-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)&-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\-(b''_{36})^{(7,7,7)}(G_{39},t)&-(b''_{40})^{(8,8,8)}(G_{43},t)&-(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt}=(b_{17})^{(2)}T_{16}-\begin{bmatrix}(b'_{17})^{(2)}&-(b''_{17})^{(2)}(G_{19},t)&-(b''_{14})^{(1,1,)}(G,t)&-(b''_{21})^{(3,3,3,)}(G_{23},t)\\-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)&-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)&-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)\\-(b''_{37})^{(7,7,7)}(G_{39},t)&-(b''_{41})^{(8,8,8)}(G_{43},t)&-(b''_{45})^{(9,9)}(G_{47},t)\end{bmatrix}T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(EEEEEEEE) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(FFFFFFFF) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(GGGGGGGG) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(HHHHHHHH)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(IIIIIIII) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(JJJJJJJJ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ff) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ff) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO E

Schmidt Decomposition, Probability Distribution And Tensor Product Of States

INTRODUCTION—VARIABLES USED

Entanglement distillation (Wikipedia)

(1) Let an entangled state $|\psi\rangle$ have a Schmidt decomposition: $|\psi\rangle = \sum_x \sqrt{p(x)}|x_A\rangle|x_B\rangle$

Here coefficients p(x) form a probability distribution, and thus are positive valued and sum to unity.

(2) The tensor product of this state is then,

$$|\psi\rangle^{\otimes m} = \sum_{x_1,x_2,\ldots,x_m} \sqrt{p(x_1)p(x_2)\ldots p(x_m)}|x_{1A}x_{2A}\ldots x_{mA}\rangle|x_{1B}x_{2B}\ldots x_{mB}\rangle$$

(3) Now, omitting all terms $x_1, \ldots, x_m$ which are not part of any sequence which is likely to occur with high probability, known as the typical set : $A_\epsilon^{(n)}$ the new state is (=)

$$|\phi_m\rangle = \sum_{x\epsilon A_\epsilon^{(n)}} \sqrt{p(x_1)p(x_2)\ldots p(x_m)}|x_{1A}x_{2A}\ldots x_{mA}\rangle|x_{1B}x_{2B}\ldots x_{mB}\rangle$$

(4) And renormalizing, $$|\phi_m'\rangle = \frac{|\phi_m\rangle}{\sqrt{\langle\phi_m|\phi_m\rangle}}$$

(5) Then the fidelity $F(|\psi\rangle^{\otimes m}, |\phi_m'\rangle) \to 1$ as $m \to \infty$.

(6) Suppose that Alice and Bob are (=) in possession of m copies of $|\psi\rangle$.

(7) Alice can perform a measurement onto the typical set $A_\epsilon^{(n)}$ subset of p_ψ, converting (eb) the state $|\psi\rangle^{\otimes m} \to |\phi_m\rangle$ with high fidelity.

(8) The theorem of typical sequences then shows us that $1-\delta$ is the probability that the given sequence is (=) part of the typical set, and may be made arbitrarily close to 1 for sufficiently large m, and therefore the Schmidt coefficients of the renormalized Bell state $|\phi_m'\rangle$ will be at most a factor $\frac{1}{\sqrt{1-\delta}}$ larger.

(9) Alice and Bob can now obtain a smaller set of n Bell states by performing LOCC on (e&eb) the state $|\phi_m'\rangle$ with which they can overcome (e) the noise of a quantum channel to communicate successfully.

NOTATION

Module One

Let an entangled state $|\psi\rangle$ have a Schmidt decomposition: $|\psi\rangle = \sum_x \sqrt{p(x)}|x_A\rangle|x_B\rangle$

Here coefficients p(x) form a probability distribution, and thus are positive valued and sum to unity.

$$|\psi\rangle = \sum_x \sqrt{p(x)}|x_A\rangle|x_B\rangle$$

G_{13} : Category one of LHS of

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of RHS of $|\psi\rangle = \sum_x \sqrt{p(x)}|x_A\rangle|x_B\rangle$ (apply the model to each term)

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

tensor product of this state is then,

$$|\psi\rangle^{\otimes m} = \sum_{x_1,x_2,\ldots,x_m} \sqrt{p(x_1)p(x_2)\ldots p(x_m)}|x_{1A}x_{2A}\ldots x_{mA}\rangle|x_{1B}x_{2B}\ldots x_{mB}\rangle$$

G_{16} : Category one of LHS of the tensor product $|\psi\rangle^{\otimes m} = \sum_{x_1,x_2,\ldots,x_m} \sqrt{p(x_1)p(x_2)\ldots p(x_m)}|x_{1A}x_{2A}\ldots x_{mA}\rangle|x_{1B}x_{2B}\ldots x_{mB}\rangle$

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of RHS of the tensor product $|\psi\rangle^{\otimes m} = \sum_{x_1,x_2,\ldots,x_m} \sqrt{p(x_1)p(x_2)\ldots p(x_m)}|x_{1A}x_{2A}\ldots x_{mA}\rangle|x_{1B}x_{2B}\ldots x_{mB}\rangle$

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Care: dual module

Now, omitting all terms $x_1, \ldots, x_m$ which are not part of any sequence which is likely to occur with high probability, known as the typical set : $A_\epsilon^{(n)}$ the new state is (=)

$$|\phi_m\rangle = \sum_{x\epsilon A_\epsilon^{(n)}} \sqrt{p(x_1)p(x_2)\ldots p(x_m)}|x_{1A}x_{2A}\ldots x_{mA}\rangle|x_{1B}x_{2B}\ldots x_{mB}\rangle$$

G_{20} : Category one of omitting all terms $x_1, \ldots, x_m$ which are not part of any sequence which is likely to occur with high probability, known as the typical set : $A_\epsilon^{(n)}$ the new state; LHS of the new state

G_{21} : Category two of SAS

G_{22} : Category three of SAS

$$|\phi_m\rangle = \sum_{x\epsilon A_\varepsilon^{(n)}} \sqrt{p(x_1)p(x_2)...p(x_m)}|x_{1A}x_{2A}...x_{mA}\rangle|x_{1B}x_{2B}...x_{mB}\rangle$$

T_{20} : Category one of ;RHS of the new state

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

renormalizing, $$|\phi_m'\rangle = \frac{|\phi_m\rangle}{\sqrt{\langle\phi_m|\phi_m\rangle}}$$

$$|\phi_m'\rangle = \frac{|\phi_m\rangle}{\sqrt{\langle\phi_m|\phi_m\rangle}}$$

G_{24} : Category one of renormalisation ; LHS of the

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of $|\phi_m'\rangle = \frac{|\phi_m\rangle}{\sqrt{\langle\phi_m|\phi_m\rangle}}$; RHS of $|\phi_m'\rangle = \frac{|\phi_m\rangle}{\sqrt{\langle\phi_m|\phi_m\rangle}}$

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Fidelity $F(|\psi\rangle^{\otimes m}, |\phi_m'\rangle) \to 1$ as $m \to \infty$.

G_{28} : Category one of fidelity

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of $F(|\psi\rangle^{\otimes m}, |\phi_m'\rangle) \to 1$ $m \to \infty$

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Alice and Bob are in possession of m copies of $|\psi\rangle$.

G_{32} : Category one of m copies of $|\psi\rangle$.

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Alice and Bob are in possession (there are lot of them, Alice and Bobs)

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Alice can perform a measurement onto the typical set $A_\epsilon^{(n)}$ subset of p_ψ, converting (eb) the state $|\psi\rangle^{\otimes m} \rightarrow |\phi_m\rangle$ with high fidelity

G_{36} : Category one of Alice can perform a measurement onto the typical set $A_\epsilon^{(n)}$ subset of p_ψ,

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of state $|\psi\rangle^{\otimes m} \rightarrow |\phi_m\rangle$ with high fidelity

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

theorem of typical sequences then shows us that $1-\delta$ is the probability that the given sequence is (=) part of the typical set, and may be made arbitrarily close to 1 for sufficiently large m, and therefore the Schmidt coefficients of the renormalized Bell state $|\phi_m^{'}\rangle$ will be at most a factor $\frac{1}{\sqrt{1-\delta}}$ larger

G_{40} : Category one of theorem of typical sequences then shows us that $1-\delta$ is the probability that the given sequence

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of part of the typical set, and may be made arbitrarily close to 1 for sufficiently large m, and therefore the Schmidt coefficients of the renormalized Bell state $|\phi_m^{'}\rangle$ will be at most a factor $\frac{1}{\sqrt{1-\delta}}$ larger

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Alice and Bob can now obtain a smaller set of n Bell states by performing LOCC on (e&eb) the state $|\phi'_m\rangle$ with which they can overcome (e) the noise of a quantum channel to communicate successfully.

G_{44} : Category one of Alice and Bob can now obtain a smaller set of n Bell states by performing LOCC; state $|\phi'_m\rangle$ with which they can overcome (e) the noise of a quantum channel to communicate successfully.

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of state $|\phi'_m\rangle$ with which they can overcome (e) the noise of a quantum channel to communicate successfully; Alice and Bob can now obtain a smaller set of n Bell states by performing LOCC

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{20})^{(3)}\,+(a''_{20})^{(3)}(T_{21},t)\;+(a''_{16})^{(2,2,2)}(T_{17},t)\;+(a''_{13})^{(1,1,1,)}(T_{14},t)\\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\;+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\;+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\;+(a''_{40})^{(8,8,8,8)}(T_{41},t)\;+(a''_{44})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a'_{21})^{(3)}\,+(a''_{21})^{(3)}(T_{21},t)\;+(a''_{17})^{(2,2,2)}(T_{17},t)\;+(a''_{14})^{(1,1,1,)}(T_{14},t)\\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\;+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\;+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\;+(a''_{41})^{(8,8,8,8)}(T_{41},t)\;+(a''_{45})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{22})^{(3)}\,+(a''_{22})^{(3)}(T_{21},t)\;+(a''_{18})^{(2,2,2)}(T_{17},t)\;+(a''_{15})^{(1,1,1,)}(T_{14},t)\\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\;+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\;+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\;+(a''_{42})^{(8,8,8,8)}(T_{41},t)\;+(a''_{46})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right] & \left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right] & \left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right] & \left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad (75)$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

$Where\ \boxed{-(b_{24}'')^{(4)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{28}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{33}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{34}'')^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a_{28}')^{(5)}\boxed{+(a_{28}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a_{29}')^{(5)}\boxed{+(a_{29}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a_{30}')^{(5)}\boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

$Where\ \boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} & -(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} & -(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} & -(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\left[-(b''_{40})^{(8)}(G_{43},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\left[-(b''_{41})^{(8)}(G_{43},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\left[-(b''_{42})^{(8)}(G_{43},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(KKKKKKKK) $\quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(LLLLLLLL) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(MMMMMMMM) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(NNNNNNNN)

$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(OOOOOOOO) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(PPPPPPPP) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(gg) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},, (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(gg) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO

Bizarre Quantum Physics And Encryption

INTRODUCTION—VARIABLES USED

More secure communications thanks to quantum physics Mar 12, 2014

(1) One of the recent revelations by Edward Snowden is that the U.S. National Security Agency is currently developing a quantum computer. Physicists aren't surprised by this news; such a computer could crack (e) the encryption that is commonly used today in no time and would therefore be highly attractive for the NSA.
(2) Professor Thomas Walther of the Institute of Applied Physics at the Technical University of Darmstadt is convinced that "Sooner or later, the quantum computer will arrive". Yet the quantum physicist is not worried. After all, he knows of an antidote: so-called quantum cryptography. This also uses (e) the bizarre rules of quantum physics, but not (e) to decrypt messages at a record pace.
(3) Quite the opposite – to encrypt it in a way that cannot be cracked by a quantum computer. To do this, a "key" that depends on the laws of quantum mechanics has to be exchanged (e&eb) between the communication partners; this then serves to encrypt the message.
(4) Physicists throughout the world are perfecting quantum cryptography to make it suitable (e) for particularly security-sensitive applications, such as for banking transactions or tap-proof communications. Walther's Ph.D. student Sabine Euler is one of them.
(5) As early as the 1980s, physicists Charles Bennett and Gilles Brassard thought about how quantum physics could help (e) transfer keys while avoiding eavesdropping.
(6) Something similar to Morse code is used, consisting (e) of a sequence of light signals from individual light particles (photons).
(7) The information is (=) in the different polarizations of successive photons.
(8) Eavesdropping is impossible due (e) to the quantum nature of photons.
(9) Any eavesdropper will inevitably be discovered because (e) the eavesdropper needs to do measurements on the photons, and these measurements will always be noticed.

NOTATION

Module One

One of the recent revelations by Edward Snowden is that the U.S. National Security Agency is currently developing a quantum computer. Physicists aren't surprised by this news; such a computer could crack (e) the encryption that is commonly used today in no time and would therefore be highly attractive for the NSA

G_{13} : Category one of encryption that is commonly used today in no time and would therefore be highly attractive for the NSA

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of quantum computer

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Professor Thomas Walther of the Institute of Applied Physics at the Technical University of Darmstadt is convinced that "Sooner or later, the quantum computer will arrive". Yet the quantum physicist is not worried. After all, he knows of an antidote: so-called quantum cryptography. This also uses (e) the bizarre

rules of quantum physics, but not (e) to decrypt messages at a record pace

G_{16} : Category one of bizarre rules of quantum physics, but not (e) to decrypt messages at a record pace

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of quantum cryptography

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

quantum cryptography knows not (e) to decrypt messages at a record pace

G_{20} : Category one of decrypt messages at a record pace

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of quantum cryptography

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Quite the opposite – to encrypt it in a way that cannot be cracked by a quantum computer. To do this, a "key" that depends on the laws of quantum mechanics has to be exchanged (e&eb) between the communication partners; this then serves to encrypt the message

G_{24} : Category one of “key" that depends on the laws of quantum mechanics; communication partners; this then serves to encrypt the message

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of communication partners; this then serves to encrypt the message; “key" that depends on the laws of quantum mechanics

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Physicists throughout the world are perfecting quantum cryptography to make it suitable (e) for particularly security-sensitive applications, such as for banking transactions or tap-proof communications. Walther's Ph.D. student Sabine Euler is one of them.

G_{28} : Category one of security-sensitive applications, such as for banking transactions or tap-proof communications

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of quantum cryptography is being improvised

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

As early as the 1980s, physicists Charles Bennett and Gilles Brassard thought about how quantum physics could help (e) transfer keys while avoiding eavesdropping

G_{32} : Category one of transfer keys while avoiding eavesdropping

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of quantum physics

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Something similar to Morse code is used, consisting (e) of a sequence of light signals from individual light particles (photons).

G_{36} : Category one of sequence of light signals from individual light particles (photons)

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of similar to Morse code is used

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Eavesdropping is impossible due (e) to the quantum nature of photons

G_{40} : Category one of quantum nature of photons

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Eavesdropping is impossible

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

eavesdropper will inevitably be discovered because (e) the eavesdropper needs to do measurements on the photons, and these measurements will always be noticed

G_{44} : Category one of eavesdropper needs to do measurements on the photons, and these measurements will always be noticed

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of eavesdropper will inevitably be discovered

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & \boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & \boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & \boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & \boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(QQQQQQQQ) $\quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(RRRRRRRR) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(SSSSSSSS) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(TTTTTTTT)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(UUUUUUUU) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(VVVVVVVV) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\left((G_{43}), t\right) = (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\left((G_{43}), t\right)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(hh) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying 193
Equations into itself

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)}t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (228)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$
Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(hh) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO A

Protocol Of Bennett And Brassard Reality

INTRODUCTION—VARIABLES USED

More secure communications thanks to quantum physics Mar 12, 2014

(1) "That's the theory" says Walther. However, there are ways to listen without (e) being noticed in practice. This has been demonstrated by hackers who specialize in quantum cryptography based on systems already available on the market.
(2) "Commercial systems have always relinquished a little bit of security in the past" says Walther. In order to make the **protocol of Bennett and Brassard reality**, you need, for example, light sources that are can be controlled (e&eb) so finely that they emit single photons in succession.
(3) Usually, a laser that is weakened so much that it emits single photons serves as (=) the light source.
(4) "But sometimes two photons can come out simultaneously, which might help (e) a potential eavesdropper to remain unnoticed" says Walther.
(5) The eavesdropper could intercept (e) the second photon and transmit the first one.
(6) Therefore, the team led by Sabine Euler uses (e) a light source that transmits (e&eb) a signal when it sends a single photon; this signal can be used (e) to select only the individually transmitted photons for communication.
(7) Nevertheless, there are still **vulnerabilities**. If the system changes the polarization of the light particles during coding, for example, (eb) the power consumption varies or the time interval of the pulses changes slightly.
(8) "An eavesdropper could tap (e) this information and read the message without the sender and receiver noticing" explains Walther. Sabine Euler and her colleagues at the Institute of Applied Physics are trying to eliminate these vulnerabilities. "They are demonstrating a lot of creativity here" says Walther approvingly. Thanks to such research, it will be harder and harder for hackers to take advantage of vulnerabilities in quantum cryptography systems.
(9) The TU Darmstadt quantum physicists want to make quantum cryptography not only more secure, but more manageable at the same time. "In a network in which many users wish to communicate securely with each other, the technology must be affordable," he says. Therefore, his team develops its systems in such a manner that they are as (=) simple as possible and can be miniaturized.

Controlled order rearrangement encryption for quantum key distribution Phys. Rev. A 68, 042315 – Published 14 October 2003 Fu-Guo Deng and G. L. Long.

(1) At the receiver's end, the order of the quantum information carrier is (=) restored.
(2) The order rearrangement operation in both parties is controlled by (e&eb) a prior shared control key which is used (e) repeatedly in a quantum key distribution session. DOI: http://dx.doi.org/10.1103/PhysRevA.68.042315

NOTATION

Module One

There are ways to listen without (e) being noticed in practice. This has been demonstrated by hackers who specialize in quantum cryptography based on systems already available on the market.

G_{13} : Category one of being noticed in practice

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of ways to listen

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

"Commercial systems have always relinquished a little bit of security in the past" says Walther. In order to make the **protocol of Bennett and Brassard reality**, you need, for example, light sources that can be controlled (e&eb) so finely that they emit single photons in succession

G_{16} : Category one of **protocol of Bennett and Brassard reality**, you need, for example, light sources; finely that they emit single photons in succession

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of finely that they emit single photons in succession **;protocol of Bennett and Brassard reality**, you need, for example, light sources

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

laser that is weakened so much that it emits single photons serves as (=) the light source

G_{20} : Category one of laser that is weakened so much that it emits single photons serves

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of light source

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

"But sometimes two photons can come out simultaneously, which might help (e) a potential eavesdropper to remain unnoticed" says Walther

G_{24} : Category one of potential eavesdropper to remain unnoticed

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of two photons can come out simultaneously

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

eavesdropper could intercept (e) the second photon and transmit the first one

G_{28} : Category one of second photon and transmit the first one

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of eavesdropper

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Therefore, the team led by Sabine Euler uses a light source that transmits a signal when it sends a single photon; this signal can be used (e) to select only the individually transmitted photons for communication

G_{32} : Category one of select only the individually transmitted photons for communication

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of light source that transmits a signal when it sends a single photon

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Nevertheless, there are still **vulnerabilities**. If the system changes the polarization of the light particles during coding, for example, (eb) the power consumption varies or the time interval of the pulses changes slightly.

G_{36} : Category one of system changes the polarization of the light particles during coding

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of power consumption varies or the time interval of the pulses changes slightly

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Eavesdropper could tap (e) this information and read the message without the sender and receiver noticing"

explains Walther. Sabine Euler and her colleagues at the Institute of Applied Physics are trying to eliminate these vulnerabilities

G_{40} : Category one of information and read the message without the sender and receiver noticing"

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Eavesdropper

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

A technique is devised to perform orthogonal state quantum key distribution. In this scheme, entangled parts of a quantum information carrier are (=) sent from Alice to Bob through two quantum channels. However, before the transmission, the order of the quantum information carrier in one channel is (=) reordered so that Eve cannot steal useful information

G_{44} : Category one of before the transmission, the order of the quantum information carrier in one channel

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of reordered so that Eve cannot steal useful information

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,

2 and 3

$[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)]$, $[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)]$, $[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}[+(a''_{16})^{(2)}(T_{17},t)] & [+(a''_{13})^{(1,1,)}(T_{14},t)] & [+(a''_{20})^{(3,3,3)}(T_{21},t)] \\ [+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)] & [+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)] & [+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)] \\ [+(a''_{36})^{(7,7,7)}(T_{37},t)] & [+(a''_{40})^{(8,8,8)}(T_{41},t)] & [+(a''_{44})^{(9,9)}(T_{45},t)] \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}[+(a''_{17})^{(2)}(T_{17},t)] & [+(a''_{14})^{(1,1,)}(T_{14},t)] & [+(a''_{21})^{(3,3,3)}(T_{21},t)] \\ [+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)] & [+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)] & [+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)] \\ [+(a''_{37})^{(7,7,7)}(T_{37},t)] & [+(a''_{41})^{(8,8,8)}(T_{41},t)] & [+(a''_{45})^{(9,9)}(T_{45},t)] \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}[+(a''_{18})^{(2)}(T_{17},t)] & [+(a''_{15})^{(1,1,)}(T_{14},t)] & [+(a''_{22})^{(3,3,3)}(T_{21},t)] \\ [+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)] & [+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)] & [+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)] \\ [+(a''_{38})^{(7,7,7)}(T_{37},t)] & [+(a''_{42})^{(8,8,8)}(T_{41},t)] & [+(a''_{46})^{(9,9)}(T_{45},t)] \end{bmatrix} G_{18} \quad (63)$$

Where $[+(a''_{16})^{(2)}(T_{17},t)]$, $[+(a''_{17})^{(2)}(T_{17},t)]$, $[+(a''_{18})^{(2)}(T_{17},t)]$ are first augmentation coefficients for category 1, 2 and 3

$[+(a''_{13})^{(1,1,)}(T_{14},t)]$, $[+(a''_{14})^{(1,1,)}(T_{14},t)]$, $[+(a''_{15})^{(1,1,)}(T_{14},t)]$ are second augmentation coefficient for category 1, 2 and 3

$[+(a''_{20})^{(3,3,3)}(T_{21},t)]$, $[+(a''_{21})^{(3,3,3)}(T_{21},t)]$, $[+(a''_{22})^{(3,3,3)}(T_{21},t)]$ are third augmentation coefficient for category 1, 2 and 3

$[+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)]$, $[+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)]$, $[+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)]$ are fourth augmentation coefficient for category 1, 2 and 3

$[+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)]$, $[+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)]$, $[+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)]$ are fifth augmentation coefficient for category 1, 2 and 3

$[+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)]$, $[+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)]$, $[+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)]$ are sixth augmentation coefficient for category 1, 2 and 3

$[+(a''_{36})^{(7,7,7)}(T_{37},t)]$, $[+(a''_{37})^{(7,7,7)}(T_{37},t)]$, $[+(a''_{38})^{(7,7,7)}(T_{37},t)]$ are seventh augmentation coefficient for category 1, 2 and 3

$[+(a''_{40})^{(8,8,8)}(T_{41},t)]$, $[+(a''_{41})^{(8,8,8)}(T_{41},t)]$, $[+(a''_{42})^{(8,8,8)}(T_{41},t)]$ are eight augmentation coefficient for category 1, 2 and 3

$[+(a''_{44})^{(9,9)}(T_{45},t)]$, $[+(a''_{45})^{(9,9)}(T_{45},t)]$, $[+(a''_{46})^{(9,9)}(T_{45},t)]$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}[-(b''_{16})^{(2)}(G_{19},t)] & [-(b''_{13})^{(1,1,)}(G,t)] & [-(b''_{20})^{(3,3,3,)}(G_{23},t)] \\ [-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)] & [-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)] & [-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)] \\ [-(b''_{36})^{(7,7,7)}(G_{39},t)] & [-(b''_{40})^{(8,8,8)}(G_{43},t)] & [-(b''_{44})^{(9,9)}(G_{47},t)] \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}[-(b''_{17})^{(2)}(G_{19},t)] & [-(b''_{14})^{(1,1,)}(G,t)] & [-(b''_{21})^{(3,3,3,)}(G_{23},t)] \\ [-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)] & [-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)] & [-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)] \\ [-(b''_{37})^{(7,7,7)}(G_{39},t)] & [-(b''_{41})^{(8,8,8)}(G_{43},t)] & [-(b''_{45})^{(9,9)}(G_{47},t)] \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$ 103

$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)}$ 104

$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)}$ 105

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\widehat{M}_{16})^{(2)}t}$ 107

$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\widehat{M}_{16})^{(2)}t}$ 108

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$\frac{1}{(\widehat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1$ 110

$\frac{1}{(\widehat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1$ 111

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(WWWWWWWW) $\quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(XXXXXXXX) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(YYYYYYYY) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(ZZZZZZZZ)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(AAAAAAAAA) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(BBBBBBBBB) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ii) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying 193
Equations into itself

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (228)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)}$$ 267

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)}$$ 268

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ii) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY TWO B

Quantum Key Distribution Protocol In The "Device-Independent" Security Scenario

INTRODUCTION—VARIABLES USED

More secure communications thanks to quantum physics Mar 12, 2014

The research team is part of the Center for Advanced Security Research Darmstadt (CASED), in which the TU Darmstadt, the Fraunhofer Institute for Secure Information Technology and the University of Darmstadt combine their expertise in current and future IT security issues. Over 200 scientists conduct research in CASED, funded by the State Initiative for Economic and Academic Excellence (LOEWE) of the Hessian Ministry for Science and the Arts. "We also exchange information with computer scientists, which is very exciting," says Walther. After all, the computer science experts deal with many of the same issues as Walther's quantum physicists. For example, Johannes Buchmann of the department of Computer Science at the TU Darmstadt is also working on encryption methods that theoretically cannot be cracked by a quantum computer. However, these are not based on quantum physics phenomena, but rather on an unsolvable math problem. Therefore, it may well be that the answer to the first code-cracking quantum computer comes from Darmstadt.

Bizarre quantum physics and encryption

(1) A quantum computer could quickly crack current encryptions because (e) it can test very many possibilities simultaneously, in the same way as (=) if you could try all possible variations for a password at once.
(2) After all, according to the quantum physics principle of superposition, atoms, electrons or photons can have (e) several states simultaneously; for example, they can rotate (e&eb) clockwise and counterclockwise at the same time.
(3) However, if you were to measure a property of a particle, such as the direction of rotation, (eb) the superposition is lost.
(4) This phenomenon is useful (e) for quantum cryptography.
(5) Eavesdroppers inevitably betray themselves because (e) their measurements of the photon change the photon's characteristics.
(6) Moreover, quantum physics forbids (e) them to copy the photon with all its properties.
(7) Therefore, they cannot siphon off (e) any information to retransmit the uninfluenced photons on to (e&eb) the sender of the message. Provided by Technische Universitat Darmstadt

Device-Independent Security of Quantum Cryptography against Collective Attacks Phys. Rev. Lett 98, 230501 – Published 4 June 2007 Antonio Acín

(8) Authors present the optimal collective attack on (e&eb) a **quantum key distribution protocol in the "device-independent" security scenario,** where no (e) assumptions are made about the way the quantum key distribution devices work or on (eb) what quantum system they operate.
(9) Our main result is (=) a tight bound on the Holevo information between one of the authorized parties and the eavesdropper, as (=) a function of the amount of violation of a Bell-type inequality. DOI: http://dx.doi.org/10.1103/PhysRevLett.98.230501

Controlled order rearrangement encryption for quantum key distribution Phys. Rev. A 68, 042315 – Published 14 October 2003 Fu-Guo Deng and G. L. Long.

(1) At the receiver's end, the order of the quantum information carrier is (=) restored.
(2) The order rearrangement operation in both parties is controlled by (e&eb) a prior shared control key which is used (e) repeatedly in a quantum key distribution session. DOI: http://dx.doi.org/10.1103/PhysRevA.68.042315

NOTATION

Module One

Quantum computer could quickly crack current encryptions because (e) it can test very many possibilities simultaneously, in the same way as (=) if you could try all possible variations for a password at once.

G_{13} : Category one of quantum computer can test very many possibilities simultaneously, in the same way as (=) if you could try all possible variations for a password at once.

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Quantum computer could quickly crack current encryptions

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

After all, according to the quantum physics principle of superposition, atoms, electrons or photons can have several states simultaneously; for example, they can rotate (e&eb) clockwise and counterclockwise at the same time

G_{16} : Category one of atoms, electrons or photons can have several states simultaneously; clockwise and counterclockwise at the same time

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of clockwise and counterclockwise at the same time; atoms, electrons or photons can have several states simultaneously

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

if you were to measure a property of a particle, such as the direction of rotation, (eb) the superposition is lost.

G_{20} : Category one of measurement of a property of a particle, such as the direction of rotation

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of superposition is lost.

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Eavesdroppers inevitably betray themselves because (e) their measurements of the photon change the photon's characteristics

G_{24} : Category one of measurements of the photon change the photon's characteristics

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Eavesdroppers inevitably betray themselves

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

quantum physics forbids (e) them to copy the photon with all its properties

G_{28} : Category one of copy the photon with all its properties

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of quantum physics

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Eavesdroppers cannot siphon off any information to retransmit the uninfluenced photons on to (e&eb) the sender of the message

G_{32} : Category one of information to retransmit the uninfluenced photons on to (e&eb) the sender of the message

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Eavesdroppers cannot siphon off

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Eavesdroppers cannot siphon off any information to retransmit the uninfluenced photons on to (e&eb) the sender of the message

G_{36} : Category one of Eavesdroppers cannot siphon off any information to retransmit the uninfluenced photons; sender of the message

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of sender of the message; Eavesdroppers cannot siphon off any information to retransmit the uninfluenced photons

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Authors present the optimal collective attack on a **quantum key distribution protocol in the "device-independent" security scenario,** where no (e) assumptions are made about the way the quantum key distribution devices work or on (eb) what quantum system they operate

G_{40} : Category one of optimal collective attack; **quantum key distribution protocol in the "device-independent" security scenario,** where no (e) assumptions are made about the way the quantum key distribution devices work or on (eb) what quantum system they operate

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **quantum key distribution protocol in the "device-independent" security scenario,** where no (e) assumptions are made about the way the quantum key distribution devices work or on (eb) what quantum system they operate; optimal collective attack

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

optimal collective attack on a **quantum key distribution protocol in the "device-independent" security scenario,** where no (e) assumptions are made about the way the quantum key distribution devices work or on what quantum system they operate

G_{44} : Category one of assumptions are made about the way the quantum key distribution devices work or on what quantum system they operate

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of optimal collective attack on a **quantum key distribution protocol in the "device-independent" security scenario**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad (57)$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad (60)$$

Where $-(b_{13}'')^{(1)}(G,t)$, $-(b_{14}'')^{(1)}(G,t)$, $-(b_{15}'')^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,)}(G_{19},t)$, $-(b_{17}'')^{(2,2,)}(G_{19},t)$, $-(b_{18}'')^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{16}')^{(2)} & +(a_{16}'')^{(2)}(T_{17},t) & +(a_{13}'')^{(1,1,)}(T_{14},t) \quad +(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a_{17}')^{(2)} & +(a_{17}'')^{(2)}(T_{17},t) & +(a_{14}'')^{(1,1,)}(T_{14},t) \quad +(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a_{18}')^{(2)} & +(a_{18}'')^{(2)}(T_{17},t) & +(a_{15}'')^{(1,1,)}(T_{14},t) \quad +(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad (63)$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\,-(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\,-(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\,-(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b_{20}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{16}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{13}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{15}'')^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{36}'')^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b_{38}'')^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{24}')^{(4)} & \boxed{+(a_{24}'')^{(4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a_{25}')^{(4)} & \boxed{+(a_{25}'')^{(4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{33}'')^{(6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{26}')^{(4)} & \boxed{+(a_{26}'')^{(4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{26} \quad 75$$

$\boxed{(a_{24}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{25}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{26}'')^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)}$

are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$

are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}-(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t)\\ -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t)\\ -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t)\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}-(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t)\\ -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t)\\ -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t)\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}-(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t)\\ -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t)\\ -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t)\end{bmatrix}T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\begin{bmatrix}(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t)\\ +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t)\\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients*

$for\ category$ 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ $are\ \ third\ \ detrition\ coefficients$ $for\ category$ 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt}=(a_{32})^{(6)}G_{33} \qquad 85$$

$$-\begin{bmatrix}(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t)\\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{32}$$

$$\frac{dG_{33}}{dt}=(a_{33})^{(6)}G_{32}-\begin{bmatrix}(a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t)\\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{33} \qquad 86$$

$$\frac{dG_{34}}{dt}=(a_{34})^{(6)}G_{33}-\begin{bmatrix}(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t)\\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)\\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)\end{bmatrix}G_{34} \qquad 87$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ $are\ \ first\ \ augmentation\ coefficients$ $for\ category$ 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ $are\ second\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ $are\ third\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$
seventh augmentation coefficients
$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$
Eighth augmentation coefficients
$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt}=(b_{32})^{(6)}T_{33}-\begin{bmatrix}(b'_{32})^{(6)}\,-(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t)\\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt}=(b_{33})^{(6)}T_{32}-\begin{bmatrix}(b'_{33})^{(6)}\,-(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t)\\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt}=(b_{34})^{(6)}T_{33}-\begin{bmatrix}(b'_{34})^{(6)}\,-(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t)\\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3
$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3
$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\boxed{-(b_{44}'')^{(9)}(G_{47},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\boxed{-(b_{45}'')^{(9)}(G_{47},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\boxed{-(b_{46}'')^{(9)}(G_{47},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} \ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$ 123

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(CCCCCCCCC) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(DDDDDDDDD) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(EEEEEEEEE) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(FFFFFFFFF)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(GGGGGGGGG) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(HHHHHHHHH) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{32})^{(6)}, T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(jj) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25}\ \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t\ > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} is bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\leq G_{15}(t)\leq\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\leq T_{13}(t)\leq\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\leq T_{15}(t)\leq \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\leq-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\leq-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\leq-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\leq-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^0 e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(jj) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523
A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^{*}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^{*}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b''_i)^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left(\ (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)\ (\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY THREE

Antibunching Dip In The Resonance Fluorescence

INTRODUCTION—VARIABLES USED

Highly stable quantum light source for applications in quantum information developed Nov 04, 2013

(1) Physicists at the University of Basel have been successful in generating photons - the quantum particles of light – with (e) only one color.
(2) This is useful for quantum information. The scientists have actively stabilized the wavelength of the photons emitted by a semiconductor thereby neutralizing (e) the charge noise in the semiconductor. The results were developed in close collaboration with the Universities of Bochum, Paderborn and Lyon and have been published in the magazine Physical Review X.
(3) Light consists of quantum particles, so-called photons. With a single photon it is possible to transfer (e&eb) quantum information.
(4) The information can be encoded in the polarization or in the phase of the photons' wave packets and can be used (e) in quantum communication and computation.
(5) In such applications, a single-photon source, a device that emits photons one by one, is a prerequisite. One of the most promising platforms for single-photon sources is based (e) on semiconductor quantum dots.
(6) One major unsolved problem is, however, that the "color" (or wavelength) of the photons emitted by a quantum dot is (=) not locked to a precise value; rather, it wanders around randomly.
(7) The fluctuations in the wavelength of the photons originate from (e) imperfections in the vicinity of the quantum dot.
(8) These imperfections can trap (e) electric charge in the semiconductor resulting (eb) in noise.
(9) To remove this "charge noise", Prof. Warburton of the Department of Physics at the University of Basel and his team has developed a quantum-classical hybrid system that connects a single quantum dot to (e&eb) a constant-wavelength laser.

NOTATION

Module One

Physicists at the University of Basel have been successful in generating photons - the quantum particles of light – with (e) only one color

G_{13} : Category one of photons - the quantum particles of light – with only one color

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of produced (generated)

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

scientists have actively stabilized the wavelength of the photons emitted by a semiconductor thereby neutralizing (e) the charge noise in the semiconductor

G_{16} : Category one of charge noise in the semiconductor

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of scientists have actively stabilized the wavelength of the photons emitted by a semiconductor thereby neutralizing

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Light consists of quantum particles, so-called photons. With a single photon it is possible to transfer (e&eb) quantum information

G_{20} : Category one of single photon it is possible to transfer; quantum information

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of quantum information ;single photon it is possible to transfer

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

information can be encoded in the polarization or in the phase of the photons' wave packets and can be used (e) in quantum communication and computation

G_{24} : Category one of quantum communication and computation

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of information can be encoded in the polarization or in the phase of the photons' wave packets

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

In such applications, a single-photon source, a device that emits photons one by one, is a prerequisite. One of the most promising platforms for single-photon sources is based (e) on semiconductor quantum dots.

G_{28} : Category one of semiconductor quantum dots

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of single-photon source, a device that emits photons one by one, is a prerequisite. One of the most promising platforms for single-photon sources

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

One major unsolved problem is, however, that the "color" (or wavelength) of the photons emitted by a quantum dot is (=) not locked to a precise value; rather, it wanders around randomly

G_{32} : Category one of major unsolved problem is, however, that the "color" (or wavelength) of the photons emitted by a quantum dot

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of not locked to a precise value; rather, it wanders around randomly

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Fluctuations in the wavelength of the photons originate from (e) imperfections in the vicinity of the quantum

dot

G_{36} : Category one of imperfections in the vicinity of the quantum dot

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Fluctuations in the wavelength of the photons originate

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Imperfections can trap electric charge in the semiconductor resulting (eb) in noise.

G_{40} : Category one of electric charge in the semiconductor resulting (eb) in noise

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Imperfections

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Imperfections can trap electric charge in the semiconductor resulting (eb) in noise.

G_{44} : Category one of Imperfections can trap electric charge in the semiconductor

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of noise.

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION TWENTY THREE A

Stream Of Single-Color Photons

INTRODUCTION—VARIABLES USED

Highly stable quantum light source for applications in quantum information developed Nov 04, 2013

(1) This stabilizing mechanism monitors (e&eb) continuously the fluctuations via the highly sensitive optical absorption of the quantum dot.
(2) By applying the exact opposite effect, the electrical field experienced by the quantum dot can be actively regulated.

Stream of single-color photons

(3) With this system, the scientists succeeded in generating (eb) a nearly perfect stream of single-color photons.
(4) A notable point is that a quantum system could be made technically useful by using (e) a classical feedback scheme, a general feature which has not been demonstrated up until now.
(5) This new scheme - through its highly effective removal (e) of the charge noise - potentially enables (eb) a stable single-photon source and may lead (eb) for example, to improvement in semiconductor-based spin qubits. The study was supported by the National Center of Competence in Research "QSIT – Quantum Science and Technology", for which the University of Basel acts as Co-Leading-House.

More information: Jonathan H. Prechtel, Andreas V. Kuhlmann, Julien Houel, Lukas Greuter, Arne Ludwig, Dirk Reuter, Andreas D. Wieck, and Richard J. Warburton, Frequency-Stabilized Source of Single Photons from a Solid-State Qubit, Phys. Rev. X 3, 041006 (2013) | DOI: 10.1103/PhysRevX.3.041006 Journal reference: Physical Review X Provided by University of Basel

(6) Single quantum dots are (=) solid-state emitters that mimic (e&eb) two-level atoms but with a highly enhanced spontaneous emission rate.
(7) A single quantum dot is (=) the basis for a potentially excellent single-photon source.
(8) One outstanding problem is that there is considerable noise in the emission frequency, making it very difficult to couple the quantum dot to (e&eb) another quantum system.
(9) Authors solve this problem here with a dynamic feedback technique that locks (e) the quantum-dot emission frequency to a reference.

NOTATION

Module One

This stabilizing mechanism monitors (e&eb) continuously the fluctuations via the highly sensitive optical absorption of the quantum dot

G_{13} : Category one of stabilizing mechanism **;** continuously the fluctuations via the highly sensitive optical absorption of the quantum dot

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of continuously the fluctuations via the highly sensitive optical absorption of the quantum dot ;stabilizing mechanism

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

By applying the exact opposite effect, the electrical field experienced by the quantum dot can be actively

regulated.

G_{16} : Category one of exact opposite effect; electrical field experienced by the quantum dot can be actively regulated

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of electrical field experienced by the quantum dot can be actively regulated; exact opposite effect

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

With this system, the scientists succeeded in generating (eb) a nearly perfect stream of single-color photons

G_{20} : Category one of electrical field experienced by the quantum dot can be actively regulated

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of perfect stream of single-color photons

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

A notable point is that a quantum system could be made technically useful by using (e) a classical feedback scheme, a general feature which has not been demonstrated up until now

G_{24} : Category one of

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of quantum system could be made technically useful

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

This new scheme - through its highly effective removal of the charge noise - potentially enables (eb) a stable single-photon source and may lead (eb) for example, to improvement in semiconductor-based spin qubits

G_{28} : Category one of charge noise - potentially enables (eb) a stable single-photon source and may lead (eb) for example, to improvement in semiconductor-based spin qubits

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of new scheme - through its highly effective

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

new scheme - through its highly effective removal of the charge noise - potentially enables a stable single-photon source and may lead (eb) for example, to improvement in semiconductor-based spin qubits

G_{32} : Category one of new scheme - through its highly effective removal of the charge noise - potentially enables

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of stable single-photon source and may lead (eb) for example, to improvement in semiconductor-based spin qubits

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

new scheme - through its highly effective removal of the charge noise - potentially enables a stable single-photon source and may lead (eb) for example, to improvement in semiconductor-based spin qubits

G_{36} : Category one of new scheme - through its highly effective removal of the charge noise - potentially enables a stable single-photon source

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of improvement in semiconductor-based spin qubits

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Single quantum dots are solid-state emitters that mimic (e&eb) two-level atoms but with a highly enhanced spontaneous emission rate.

G_{40} : Category one of Single quantum dots; Single quantum dots are solid-state emitters that mimic; two-level atoms but with a highly enhanced spontaneous emission rate.

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of solid-state emitters that mimic (e&eb) two-level atoms but with a highly enhanced spontaneous emission rate; two-level atoms but with a highly enhanced spontaneous emission rate.; Single quantum dots are solid-state emitters that mimic;

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

single quantum dot is (=) the basis for a potentially excellent single-photon source

G_{44} : Category one of single quantum dot

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of basis for a potentially excellent single-photon source

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(IIIIIIIII) $\qquad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(JJJJJJJJJ)The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(KKKKKKKKK) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(LLLLLLLLL)

$\lim_{G\to\infty}(b_i'')^{(7)}((G_{39}),t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}((G_{39}),t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(MMMMMMMMM) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(NNNNNNNNN) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$ *and* $(\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(kk) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$

$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$

$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)},\ if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, if\ (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$

and $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)},\ if\ (\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)},\ \boldsymbol{if}\ (\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)}$, ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \qquad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \qquad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \qquad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(kk) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY THREE B

Coulomb Interaction Between The Coupled Dots Mutually Affects The Energy Levels Of The Other Coupled Dots.

INTRODUCTION—VARIABLES USED

More information: Jonathan H. Prechtel, Andreas V. Kuhlmann, Julien Houel, Lukas Greuter, Arne Ludwig, Dirk Reuter, Andreas D. Wieck, and Richard J. Warburton, Frequency-Stabilized Source of Single Photons from a Solid-State Qubit, Phys. Rev. X 3, 041006 (2013) | DOI: 10.1103/PhysRevX.3.041006 Journal reference: Physical Review X Provided by University of Basel

(1) The incoherent scattering (resonance fluorescence) represents (eb) the single-photon output, whereas the coherent scattering (Rayleigh scattering) is used (e) for the feedback control.
(2) The fluctuations in emission frequency are (=) reduced to 20 MHz, just approximately 5% of the quantum-dot optical linewidth, even over several hours.
(3) By eliminating the 1/f-like noise, the relative fluctuations in quantum-dot noise power are (=) reduced to approximately 10−5 at low frequency.
(4) Under these conditions, the **antibunching dip in the resonance fluorescence** is described extremely well by (e) the two-level atom result.
(5) The technique represents a way of removing (e) charge noise from a quantum device. **DOI: http://dx.doi.org/10.1103/PhysRevX.3.041006**

Quantum gates by coupled asymmetric quantum dots and controlled-NOT-gate operation Phys. Rev. A 61, 022305 – Published 12 January 2000 Tetsufumi Tanamoto

(6) A quantum computer based (e) on an asymmetric coupled-dot system has been proposed and shown to operate as (=) a controlled-NOT gate.
(7) The basic ideas are the following. The electron is (=) localized in one of the asymmetric coupled dots.
(8) The electron transfer takes place from one dot to (e&eb) the other when the energy levels of the coupled dots are (=) set to be close.
(9) The Coulomb interaction between the coupled dots mutually affects (e&eb) the energy levels of the other coupled dots.
(10) Decoherence time of the quantum computation and the measurement time are estimated. The proposed system can be realized by (e) developing the technology of the single-electron memory using Si nanocrystals, and a direct combination of the quantum circuit and the conventional circuit is possible. DOI: http://dx.doi.org/10.1103/PhysRevA.61.022305

NOTATION

Module One

incoherent scattering (resonance fluorescence) represents the single-photon output, whereas the coherent scattering (Rayleigh scattering) is used (e) for the feedback control

G_{13} : Category one of incoherent scattering (resonance fluorescence)

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of single-photon output, whereas the coherent scattering (Rayleigh scattering) is used (e) for the feedback control

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Fluctuations in emission frequency are (=) reduced to 20 MHz, just approximately 5% of the quantum-dot optical linewidth, even over several hours.

G_{16} : Category one of Fluctuations in emission frequency

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of reduced to 20 MHz, just approximately 5% of the quantum-dot optical linewidth, even over several hours (take in to consideration different such experiments; model is applicable to all such experiments)

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

By eliminating the 1/f-like noise, the relative fluctuations in quantum-dot noise power are (=) reduced to approximately 10−5 at low frequency

G_{20} : Category one of elimination of the 1/f-like noise

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of relative fluctuations in quantum-dot noise power are reduced to approximately 10−5 at low frequency

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Antibunching dip in the resonance fluorescence is described extremely well by (e) the two-level atom result.

G_{24} : Category one of two-level atom result.

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **Antibunching dip in the resonance fluorescence** is described extremely well

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

technique represents a way of removing (e) charge noise from a quantum device

G_{28} : Category one of charge noise from a quantum device

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of technique represents a way of removing

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Quantum computer based (e) on an asymmetric coupled-dot system has been proposed and shown to operate as (=) a controlled-NOT gate.

G_{32} : Category one of asymmetric coupled-dot system has been proposed and shown to operate as a controlled-NOT gate

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Quantum computer

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Electron is (=) localized in one of the asymmetric coupled dots.

G_{36} : Category one of Electron

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of localized in one of the asymmetric coupled dots

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

electron transfer takes place from one dot to the other when the energy levels of the coupled dots are (=) set to be close

G_{40} : Category one of electron transfer takes place from one dot to the other when the energy levels of the coupled dots

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of set to be close

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Coulomb interaction between the coupled dots mutually affects (e&eb) the energy levels of the other coupled dots.

G_{44} : Category one of Coulomb interaction between the coupled dots; energy levels of the other coupled dots.

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of energy levels of the other coupled dots.; Coulomb interaction between the coupled dots

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30}$ 27

$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28}$ 28

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29}$ 29

$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30}$ 30

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32}$ 31

$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33}$ 32

$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34}$ 33

$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32}$ 34

$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33}$ 35

$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34}$ 36

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36}$ 37

$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37}$ *38*

$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38}$ *39*

$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36}$ *40*

$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37}$ *41*

$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38}$ *42*

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40}$ *43*

$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41}$ *44*

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{c} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{array}\right] T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{c} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{array}\right] T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{c} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{array}\right] T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category

1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t)+(a_{16}'')^{(2,2,2)}(T_{17},t)+(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)+(a_{40}'')^{(8,8,8,8)}(T_{41},t)+(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t)+(a_{17}'')^{(2,2,2)}(T_{17},t)+(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)+(a_{41}'')^{(8,8,8,8)}(T_{41},t)+(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t)+(a_{18}'')^{(2,2,2)}(T_{17},t)+(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)+(a_{42}'')^{(8,8,8,8)}(T_{41},t)+(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{32}'')^{(6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b_{13}'')^{(1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a_{28}')^{(5)} & +(a_{28}'')^{(5)}(T_{29},t) & +(a_{24}'')^{(4,4,)}(T_{25},t) & +(a_{32}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a_{29}')^{(5)} & +(a_{29}'')^{(5)}(T_{29},t) & +(a_{25}'')^{(4,4,)}(T_{25},t) & +(a_{33}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a_{30}')^{(5)} & +(a_{30}'')^{(5)}(T_{29},t) & +(a_{26}'')^{(4,4,)}(T_{25},t) & +(a_{34}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad (85)$$

$$- \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b_{28}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b_{24}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}, \boxed{(a_{37}'')^{(7)}(T_{37},t)}, \boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(OOOOOOOOO) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(PPPPPPPPP) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

(QQQQQQQQQ) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(RRRRRRRRR)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

(SSSSSSSSS) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

(TTTTTTTTT) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(ll) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right| \; e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)}$, ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(ll) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY FOUR

Principle Of Locality

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

Principle of locality

(1) Principle of locality states that an object is influenced (e&eb) directly by its immediate surroundings.
(2) A physical theory is said to be a **local theory** if it is consistent with (e&eb) the principle of locality.
(3) To date, no test has simultaneously closed (e) all loopholes to the idea that entangled particles violate (e) the principle of locality or engage in superluminal communication.
(4) In the 17th Century Newton's law of universal gravitation was formulated in terms of "action at a distance", thereby violating (e) the principle of locality.
(5) It is inconceivable that inanimate Matter should, without the Mediation of something else, which is not material, operate upon, and affect (e&eb) other matter without mutual Contact…
(6) That Gravity should be innate, inherent and essential to (e&eb) Matter, so that one body may act upon another at a distance thro' (e&eb) a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws; but whether this Agent is material or immaterial, I have left to the Consideration of my readers.

—Isaac Newton, Letters to Bentley, 1692/3

EPR paradox

(7) Albert Einstein argued that quantum mechanics was an incomplete physical theory. Using the principle of locality, he, Podolsky, and Rosen articulated the Einstein-Podolsky-Rosen paradox which showed that **quantum mechanics predicts (eb) non-locality** unless position and momentum were (=) simultaneous "real" properties of a particle.
(8) The locality question remained unverifiable for several decades. Then in 1964, John Stewart Bell derived his eponymous theorem, which describes quantum mechanical predictions that no theory of local hidden variables, no local realism, could ever reproduce (eb)
(9) In his groundbreaking 1964 paper, "On the Einstein Podolsky Rosen paradox", physicist John Stewart Bell presented an analogy (based on spin measurements on pairs of entangled electrons) to EPR's hypothetical paradox. Using their reasoning, he said, a choice of measurement setting here should not (e) affect the outcome of a measurement there (and vice versa). After providing a mathematical formulation of locality and realism based on this, he showed specific cases where this would be inconsistent with the predictions of QM theory.

NOTATION

Module One

Principle of locality states that an object is influenced (e&eb) directly by its immediate surroundings

G_{13} : Category one of Principle of locality states that an object ; directly by its immediate surroundings

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of directly by its immediate surroundings; Principle of locality states that an object

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Physical theory is said to be a **local theory** if it is consistent with (e&eb) the principle of locality.

G_{16} : Category one of Physical theory is said to be a **local theory;** principle of locality

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of principle of locality ;Physical theory is said to be a **local theory**

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

no test has simultaneously closed (e) all loopholes to the idea that entangled particles violate (e) the principle of locality or engage in superluminal communication

entangled particles violate (e) the principle of locality or engage in superluminal communication

G_{20} : Category one of all loopholes to the idea that entangled particles violate (e) the principle of locality or engage in superluminal communication; principle of locality or engage in superluminal communication

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of no test has simultaneously; entangled particles

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Newton's law of universal gravitation was formulated in terms of "action at a distance", thereby violating (e) the principle of locality

G_{24} : Category one of principle of locality

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of law of universal gravitation was formulated in terms of "action at a distance",

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

It is inconceivable that inanimate Matter should, without the Mediation of something else, which is not material, operate upon, and affect (e&eb) other matter without mutual Contact

G_{28} : Category one of inanimate Matter should, without the Mediation of something else, which is not material, operate upon, and affect; matter without mutual Contact

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of matter without mutual Contact; inanimate Matter should, without the Mediation of something else, which is not material, operate upon, and affect

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance thro' (e&eb) a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws; but whether this Agent is material or immaterial, I have left to the Consideration of my readers.

G_{32} : Category one of Gravity should be innate, inherent and essential; Matter, so that one body may act upon another at a distance thro' (e&eb) a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws; but whether this Agent is material or immaterial, I have left to the Consideration of my readers

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Matter, so that one body may act upon another at a distance thro' (e&eb) a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws; but whether this Agent is material or immaterial, I have left to the Consideration of my readers; Gravity should be innate, inherent and essential

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a

distance thro' a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws

G_{36} : Category one of Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance; Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to (e&eb) another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws; Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance thro' a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws

G_{40} : Category one of Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance thro' a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one; another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws; Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance thro' a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Gravity should be innate, inherent and essential to Matter, so that one body may act upon another at a distance thro' a Vacuum, without the Mediation of any thing else, by and through which their Action and Force may be conveyed from one to another, is to me so great an Absurdity that I believe no Man who has in philosophical Matters a competent Faculty of thinking can ever fall into it. Gravity must be caused by (e) an Agent acting constantly according to certain laws

G_{44} : Category one of Agent acting constantly according to certain laws

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Gravity

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{c} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{array}\right] T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{c} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{array}\right] T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{c} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{array}\right] T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category

1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t)+(a_{16}'')^{(2,2,2)}(T_{17},t)+(a_{13}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t)+(a_{40}'')^{(8,8,8,8)}(T_{41},t)+(a_{44}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t)+(a_{17}'')^{(2,2,2)}(T_{17},t)+(a_{14}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t)+(a_{41}'')^{(8,8,8,8)}(T_{41},t)+(a_{45}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t)+(a_{18}'')^{(2,2,2)}(T_{17},t)+(a_{15}'')^{(1,1,1,)}(T_{14},t)\\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t)+(a_{42}'')^{(8,8,8,8)}(T_{41},t)+(a_{46}'')^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad 69$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation

coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad (72)$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{32}'')^{(6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b_{13}'')^{(1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a_{28}')^{(5)} & +(a_{28}'')^{(5)}(T_{29},t) & +(a_{24}'')^{(4,4,)}(T_{25},t) & +(a_{32}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a_{29}')^{(5)} & +(a_{29}'')^{(5)}(T_{29},t) & +(a_{25}'')^{(4,4,)}(T_{25},t) & +(a_{33}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a_{30}')^{(5)} & +(a_{30}'')^{(5)}(T_{29},t) & +(a_{26}'')^{(4,4,)}(T_{25},t) & +(a_{34}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad 81$$

Where $+(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad (85)$$

$$- \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad (90)$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b_{28}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,5)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b_{24}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4,4,4,)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}, \boxed{(a_{37}'')^{(7)}(T_{37},t)}, \boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\left[-(b''_{36})^{(7)}(G_{39},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\left[-(b''_{37})^{(7)}(G_{39},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\left[-(b''_{38})^{(7)}(G_{39},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(UUUUUUUUU) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(VVVVVVVVV) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$ 132

(WWWWWWWWW) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(XXXXXXXXX)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(YYYYYYYYY) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(ZZZZZZZZZ) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(mm) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$

$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$

$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$
Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq-(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq-(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)},\ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)},\ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(mm) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY FIVE

Local Realism

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) These three key concepts – locality, realism, freedom – are highly technical and much debated. In particular, the **concept of realism** is now somewhat different from what it was in discussions in the 1930s. It is (=) more precisely called counterfactual definiteness
(2) **Realism** means that we may think of outcomes of measurements that were (=) not actually performed as (=) being just as much part of reality as (=) those that were made.
(3) Locality is short for (e) local relativistic causality.
(4) Freedom refers to the physical possibility to determine (eb) settings on measurement devices independently of (e) the internal state of the physical system being measured.
(5) Einstein assumed that the principle of locality was necessary, and that there could be no violations of it. He said: "(...) the following idea characterises (eb) the relative independence of objects far apart in space, A and B
(6) external influence on A has no (e) direct influence on B
(7) This is known as (=) the Principle of Local Action, which is used (e) consistently only in field theory.
(8) If this axiom were to be completely abolished, the idea of the existence of quasienclosed systems, and thereby the postulation of laws which can be checked empirically in the accepted sense, would become (=) impossible. (...)"
(9) Local realism is (=) the combination of the principle of locality with (e&eb) the "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement before (e) the measurement is made.

NOTATION

Module One

Albert Einstein argued that quantum mechanics was an incomplete physical theory. Using the principle of locality, he, Podolsky, and Rosen articulated the Einstein-Podolsky-Rosen paradox which showed that **quantum mechanics predicts (eb) non-locality** unless position and momentum were (=) simultaneous "real" properties of a particle.

G_{13} : Category one of **quantum mechanics**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **non-locality** unless position and momentum were (=) simultaneous "real" properties of a particle.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Using their reasoning, he said, a choice of measurement setting here should not (e) affect the outcome of a measurement there (and vice versa). After providing a mathematical formulation of locality and realism based on this, he showed specific cases where this would be inconsistent with the predictions of QM theory

G_{16} : Category one of outcome of a measurement there (and vice versa)

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of choice of measurement setting

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Concept of realism is now somewhat different from what it was in discussions in the 1930s. It is (=) more precisely called counterfactual definiteness

G_{20} : Category one of **Concept of realism**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of counterfactual definiteness

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Realism means that we may think of outcomes of measurements that were not actually performed as (=) being just as much part of reality as (=) those that were made

G_{24} : Category one of **Realism** means that we may think of outcomes of measurements; **Realism** means that we may think of outcomes of measurements that were not actually performed

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of not actually performed as (=) being just as much part of reality as (=) those that were made; part of reality as (=) those that were made

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Locality is short for (e) local relativistic causality

G_{28} : Category one of local relativistic causality

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Locality

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Freedom refers to the physical possibility to determine settings on measurement devices independently of (e) the internal state of the physical system being measured

G_{32} : Category one of Freedom refers to the physical possibility

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of settings on measurement devices independently of (e) the internal state of the physical system being measured

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Freedom refers to the physical possibility to determine settings on measurement devices independently of (e) the internal state of the physical system being measured

G_{36} : Category one of internal state of the physical system being measured

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Freedom refers to the physical possibility to determine settings on measurement devices independently

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Principle of Local Action, which is used (e) consistently only in field theory.

G_{40} : Category one of field theory

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Principle of Local Action

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Local realism is (=) the combination of the principle of locality with (e&eb) the "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement before the measurement is made.

G_{44} : Category one of principle of locality; "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement before the measurement is made.

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement before the measurement is made.; principle of locality

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21},t)\ +(a''_{16})^{(2,2,2)}(T_{17},t)\ +(a''_{13})^{(1,1,1,)}(T_{14},t)\\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)\ +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)\ +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7,7)}(T_{37},t)\ +(a''_{40})^{(8,8,8,8)}(T_{41},t)\ +(a''_{44})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{c}(a'_{21})^{(3)}+(a''_{21})^{(3)}(T_{21},t)\ +(a''_{17})^{(2,2,2)}(T_{17},t)\ +(a''_{14})^{(1,1,1,)}(T_{14},t)\\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)\ +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)\ +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7,7)}(T_{37},t)\ +(a''_{41})^{(8,8,8,8)}(T_{41},t)\ +(a''_{45})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{c}(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21},t)\ +(a''_{18})^{(2,2,2)}(T_{17},t)\ +(a''_{15})^{(1,1,1,)}(T_{14},t)\\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)\ +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)\ +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7,7)}(T_{37},t)\ +(a''_{42})^{(8,8,8,8)}(T_{41},t)\ +(a''_{46})^{(9,9,9)}(T_{45},t)\end{array}\right]G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right] & \left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right] & \left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right] & \left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ \boxed{-(b_{24}'')^{(4)}(G_{27},t)}, \boxed{-(b_{25}'')^{(4)}(G_{27},t)}, \boxed{-(b_{26}'')^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{28}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{29}'')^{(5,5,)}(G_{31},t)}, \boxed{-(b_{30}'')^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{32}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{33}'')^{(6,6,)}(G_{35},t)}, \boxed{-(b_{34}'')^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{13}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{14}'')^{(1,1,1,1)}(G,t)}, \boxed{-(b_{15}'')^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{16}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{17}'')^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b_{18}'')^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{20}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{21}'')^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b_{22}'')^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b_{46}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{45}'')^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b_{44}'')^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & \boxed{+(a_{28}'')^{(5)}(T_{29},t)} & \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & \boxed{+(a_{29}'')^{(5)}(T_{29},t)} & \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & \boxed{+(a_{30}'')^{(5)}(T_{29},t)} & \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)} \quad \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

$Where\ \boxed{+(a_{28}'')^{(5)}(T_{29},t)}, \boxed{+(a_{29}'')^{(5)}(T_{29},t)}, \boxed{+(a_{30}'')^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a_{24}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{25}'')^{(4,4,)}(T_{25},t)}, \boxed{+(a_{26}'')^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{32}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{33}'')^{(6,6,6)}(T_{33},t)}, \boxed{+(a_{34}'')^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)} & \boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & \boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & \boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & \boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & \boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} & \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} & \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(AAAAAAAAAA) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(BBBBBBBBBB) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(CCCCCCCCCC) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(DDDDDDDDDD)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(EEEEEEEEEE) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(FFFFFFFFFF) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(nn) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0$ for $t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t} \; and \; (\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(nn) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY SIX

Mind-Independent Property Does Not Have To Be The Value Of Some Physical Variable Such As Position Or Momentum

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) Local realism is (=) the combination of the principle of locality with (e&eb) the "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement before (e) the measurement is made.
(2) Local realism is (=) a significant feature of classical mechanics, of general relativity, and of electrodynamics; but quantum mechanics largely rejects (e) this principle due to the theory of distant quantum entanglements, an interpretation rejected by Einstein in the EPR paradox but subsequently proven by (e) Bell's inequalities
(3) Any theory, such as quantum mechanics, that violates (e) Bell's inequalities must abandon (e) either locality or realism
(4) some physicists dispute that experiments have demonstrated (eb) Bell's violations, on the grounds (e) that the sub-class of inhomogeneous Bell inequalities has not been tested or due (e) to experimental limitations in the tests.
(5) Different interpretations of quantum mechanics violate (e) different parts of local realism and/or counterfactual definiteness.

Realism

(6) Realism in the sense used by (e) physicists does not equate to (=) realism in metaphysics.
(7) The physicist's Realism is the claim that the world is (=) in some sense mind-independent even if the results of a possible measurement do not (e) pre-exist the act of measurement, which does not (e) require that they are the creation of the observer (contrary to the "consciousness causes (eb) collapse" interpretation of quantum mechanics).
(8) Furthermore, a **mind-independent property does not (e) have to be the value of some physical variable such as position or momentum.**
(9) A property can be dispositional (or potential), i.e., it can be (=) a tendency: in the way that glass objects tend to (e) break, or are disposed to(e) break, even if they do not actually break

NOTATION

Module One

Local realism is the combination of the principle of locality with the "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement before (e) the measurement is made

G_{13} : Category one of Local realism is the combination of the principle of locality with the "realistic" assumption that all objects must objectively have a pre-existing value for any possible measurement

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of measurement is made

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Local realism is a significant feature of classical mechanics, of general relativity, and of electrodynamics; but quantum mechanics largely rejects (e) this principle due to the theory of distant quantum entanglements, an interpretation rejected by Einstein in the EPR paradox but subsequently proven by (e) Bell's inequalities

G_{16} : Category one of principle due to the theory of distant quantum entanglements, an interpretation rejected by Einstein in the EPR paradox but subsequently proven by (e) Bell's inequalities

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Local realism is a significant feature of classical mechanics, of general relativity, and of electrodynamics; but quantum mechanics

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Any theory, such as quantum mechanics, that violates Bell's inequalities must abandon (e) either locality or realism

G_{20} : Category one of **Bell's inequalities must abandon (e) either locality or realism**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **Any theory, such as quantum mechanics, that violates**

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Any theory, such as quantum mechanics, that violates Bell's inequalities must abandon (e) either locality or realism

G_{24} : Category one of **either locality or realism**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **Any theory, such as quantum mechanics, that violates Bell's inequalities**

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Different interpretations of quantum mechanics violate (e) different parts of local realism and/or counterfactual definiteness.

G_{28} : Category one of different parts of local realism and/or counterfactual definiteness.

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Different interpretations of quantum mechanics

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Realism in the sense used by (e) physicists does not equate to realism in metaphysics

G_{32} : Category one of physicists does not equate to realism in metaphysics

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Realism in the sense

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

physicist's Realism is the claim that the world is in some sense mind-independent even if the results of a possible measurement do not (e) pre-exist the act of measurement, which does not (e) require that they are the creation of the observer (contrary to the "consciousness causes (eb) collapse" interpretation of quantum mechanics).

G_{36} : Category one of physicist's Realism is the claim that the world

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of some sense mind-independent even if the results of a possible measurement do not (e) pre-exist the act of measurement, which does not (e) require that they are the creation of the observer (contrary to the "consciousness causes (eb) collapse" interpretation of quantum mechanics).

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

physicist's Realism is the claim that the world is in some sense mind-independent even if the results of a possible measurement do not pre-exist the act of measurement, which does not require that they are the creation of the observer (contrary to the "consciousness causes (eb) collapse" interpretation of quantum mechanics).

G_{40} : Category one of pre-exist the act of measurement, which does not (e) require that they are the creation of the observer (contrary to the "consciousness causes (eb) collapse" interpretation of quantum mechanics

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of physicist's Realism is the claim that the world is in some sense mind-independent even if the results of a possible measurement

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Physicist's Realism is the claim that the world is in some sense mind-independent even if the results of a possible measurement do not pre-exist the act of measurement, which does not require that they are the creation of the observer (contrary to the "consciousness causes (eb) collapse" interpretation of quantum mechanics).

G_{44} : Category one of Physicist's Realism is the claim that the world is in some sense mind-independent even if the results of a possible measurement do not pre-exist the act of measurement, which does not require that they are the **creation of the observer (contrary to the "consciousness**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **collapse" interpretation of quantum mechanics).**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$

$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$
$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,

2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & \boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} & \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & \boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} & \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & \boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & \boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} & \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} & \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$

$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$

$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$

$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)}$ 102

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(GGGGGGGGGG) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(HHHHHHHHHH) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(IIIIIIIII) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(JJJJJJJJJ)

$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(KKKKKKKKKK) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(LLLLLLLLLL) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$ are positive constants and $i = 40,41,42$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(oo) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)} t} \leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}} \left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)} (T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)} \left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$
$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G^0_{28}e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G^0_{28}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G^0_{30}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \qquad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \qquad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \qquad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(oo) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY SEVEN

"Detection Loophole" Or The "Communication Loophole."

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) Likewise, the mind-independent properties of quantum systems could consist (e) of a tendency to respond (e&eb) to particular measurements with particular values with ascertainable probability.
(2) Such ontology would be metaphysically realistic, without (e) being realistic in the physicist's sense of "local realism" (which would require that a single value be produced (eb) with certainty).
(3) A closely related term is counterfactual definiteness (CFD), used (e) to refer to the claim that one can meaningfully speak of the definiteness of results of measurements that have not been performed (i.e., the ability to assume (eb) the existence of objects, and properties of objects, even when they have (e) not been measured).

Bell's Theorem First published Wed Jul 21, 2004; substantive revision Thu Jun 11, 2009 (Stanford Encyclopedia)

(4) Bell's Theorem is the collective name for a family of results, all showing the (eb) impossibility of a Local Realistic interpretation of quantum mechanics.
(5) There are variants of the Theorem with different meanings of "Local Realistic." In John S. Bell's pioneering paper of 1964 the realism consisted (e) in postulating in addition to the quantum state a "complete state", which determines (eb) the results of measurements on the system, either by assigning (eb) a value to the measured quantity that is revealed by (e) the measurement regardless of the details of the measurement procedure, or by enabling (eb) the system to elicit (eb) a definite response whenever it is measured, but a response which may depend on (eb) the macroscopic features of the experimental arrangement or even on (eb) the complete state of the system together with that arrangement.
(6) Locality is (=) a condition on composite systems with (e&eb) spatially separated constituents, requiring (e) an operator which is (=) the product of operators associated with (e&eb) the individual constituents to be assigned (eb) a value which is (=) the product of the values assigned to the factors, and requiring (e) the value assigned to an operator associated with (e&eb) an individual constituent to be (=) independent of what is measured on (eb) any other constituent.
(7) From his assumptions Bell proved an inequality (the prototype of "Bell's Inequality") which is (=) violated (e) by the Quantum Mechanical predictions made from (e) an entangled state of the composite system.
(8) In other variants the complete state assigns (eb) probabilities to (e) the possible results of measurements of the operators rather than determining (eb) which result will be obtained, and nevertheless inequalities are(=) derivable; and still other variants dispense (e) with inequalities.
(9) The incompatibility of Local Realistic Theories with Quantum Mechanics permits (eb) adjudication by experiments, some of which are described here.
(10) Most of the dozens of experiments performed so far have favored Quantum Mechanics, but not decisively because (e) of the **"detection loophole" or the "communication loophole."**

NOTATION

Module One

Likewise, the mind-independent properties of quantum systems could consist of a tendency to respond (e&eb) to particular measurements with particular values with ascertainable probability

G_{13} : Category one of tendency to respond (e&eb) to particular measurements with particular values with ascertainable probability

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of mind-independent properties of quantum systems

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

mind-independent properties of quantum systems could consist of a tendency to respond (e&eb) to particular measurements with particular values with ascertainable probability

G_{16} : Category one of mind-independent properties of quantum systems could consist of a tendency; particular measurements with particular values with ascertainable probability

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of particular measurements with particular values with ascertainable probability; mind-independent properties of quantum systems could consist of a tendency

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Such ontology would be metaphysically realistic, without (e) being realistic in the physicist's sense of "local realism" (which would require that a single value be produced (eb) with certainty).

G_{20} : Category one of realistic in the physicist's sense of "local realism" (which would require that a single value be produced (eb) with certainty).

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of Such ontology would be metaphysically realistic

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Counterfactual definiteness (CFD), used to refer to the claim that one can meaningfully speak of the definiteness of results of measurements that have not been performed (i.e., the ability to assume (eb) the existence of objects, and properties of objects, even when they have (e) not been measured).

G_{24} : Category one of claim that one can meaningfully speak of the definiteness of results of measurements that have not been performed (i.e., the ability to assume (eb) the existence of objects, and properties of objects, even when they have (e) not been measured).

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of counterfactual definiteness (CFD)

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Ability to assume the existence of objects, and properties of objects, even when they have (e) not been measured).

G_{28} : Category one of even when they have not been measured).

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Ability to assume the existence of objects, and properties of objects

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Bell's Theorem is the collective name for a family of results, all showing the (eb) impossibility of a Local Realistic interpretation of quantum mechanics

G_{32} : Category one of Bell's Theorem is the collective name for a family of results

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of impossibility of a Local Realistic interpretation of quantum mechanics

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Bell proved an inequality (the prototype of "Bell's Inequality") which is violated (e) by the Quantum Mechanical predictions made from (e) an entangled state of the composite system.

G_{36} : Category one of Quantum Mechanical predictions made from (e) an entangled state of the composite system.. there are millions of systems satisfying the conditionalities so specified in the foregoing

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of inequality (the prototype of "Bell's Inequality") which is violated

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

The incompatibility of Local Realistic Theories with Quantum Mechanics permits (eb) adjudication by experiments, some of which are described here

G_{40} : Category one of incompatibility of Local Realistic Theories with Quantum Mechanics

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of adjudication by experiments, some of which are described here

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Dozens of experiments performed so far have favored Quantum Mechanics, but not decisively because (e) of the **"detection loophole" or the "communication loophole."**

G_{44} : Category one of **"detection loophole" or the "communication loophole."**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Dozens of experiments performed so far have favored Quantum Mechanics

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION TWENTY EIGHT

No Physical Theory Which Is Realistic And Also Local In A Specified Sense Can Agree With All Of The Statistical Implications Of Quantum Mechanics

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) The latter has been nearly decisively blocked (e) by a recent experiment and there is a good

prospect for blocking the former.

(2) The refutation of the family of Local Realistic Theories would imply (eb) that certain peculiarities of Quantum Mechanics will remain part (e&eb) of our physical worldview: notably, the objective indefiniteness of properties, (e&eb) the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory.

Introduction

(3) In 1964 John S. Bell, a native of Northern Ireland and a staff member of CERN (European Organisation for Nuclear Research) whose primary research concerned theoretical high energy physics, published a paper in the short-lived journal Physics which transformed the study of the foundation of Quantum Mechanics (Bell 1964). The paper showed (under conditions which were relaxed in later work by Bell (1971, 1985, 1987) himself and by his followers (Clauser et al. 1969, Clauser and Horne 1974, Mermin 1986, Aspect 1983)) that **no physical theory which is (=) realistic and also local in a specified sense can agree with (=) all of the statistical implications of Quantum Mechanics**.

(4) One line of investigation in the prehistory of Bell's Theorem concerned the conjecture that the Quantum Mechanical state of a system needs (e) to be supplemented by further "elements of reality" or "hidden variables" or "complete states" in order to provide (eb) a complete description, the incompleteness of the quantum state being the explanation for (e) the statistical character of Quantum Mechanical predictions concerning the system.

(5) There are actually two main classes of hidden-variables theories. In one, which is usually called "**non-contextual",** the complete state of the system determines (eb) the value of a quantity (equivalently, an eigenvalue of the operator representing that quantity) that will be obtained by (e) any standard measuring procedure of that quantity, regardless of what other quantities are (=) simultaneously measured (eb) or what the complete state of the system and the measuring apparatus may be.

(6) The hidden-variables theories of Kochen and Specker (1967) are explicitly of this type. In the other, which is usually called **"contextual",** the **value obtained depends (=) upon what quantities are simultaneously measured and/or on the details of the complete state of the measuring apparatus**.

(7) This distinction was first explicitly pointed out by Bell (1966) but without using the terms "contextual" and "non-contextual". There actually are two quite different versions of contextual hidden-variables theories, depending upon the character of the context: an **"algebraic context"** is (=) one which specifies the quantities (or the operators representing them) which are measured (eb) jointly with the quantity (or operator) of primary interest, whereas an **"environmental context"** is (=) a specification of the physical characteristics of the measuring apparatus whereby it simultaneously measures (eb) several distinct co-measurable quantities.

(8) In Bohm's hidden-variables theory (1952) the context is (=) environmental, whereas in those of Bell (1966) and Gudder (1970) the context is (=) algebraic.

(9) A pioneering version of a "hidden variables theory" was proposed by **Louis de Broglie** in 1926–7 (de Broglie 1927, 1928) and a more complete version by David Bohm in 1952 (Bohm 1952; see also the entry on Bohmian mechanics). In these theories the entity supplementing (eb) the quantum state (which is (=) a wave function in the position representation) is (=) typically a **classical entity, located in (eb) a classical phase space and therefore characterized by (e) both position and momentum variables**.

NOTATION

Module One

refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics will remain part (e&eb) of our physical worldview: notably, the objective indefiniteness of

properties, (e&eb) the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory

G_{13} : Category one of refutation of the family of Local Realistic Theories

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of certain peculiarities of Quantum Mechanics will remain part (e&eb) of our physical worldview: notably, the objective indefiniteness of properties, (e&eb) the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics will remain part of our physical worldview: notably, the objective indefiniteness of properties, the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory

G_{16} : Category one of refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics; physical worldview: notably, the objective indefiniteness of properties, (e&eb) the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of physical worldview: notably, the objective indefiniteness of properties, (e&eb) the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory; refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics will remain part of our physical worldview: notably, the objective indefiniteness of properties, the indeterminacy of measurement results, and the tension between quantum nonlocality and (e&eb) the locality of Relativity Theory

G_{20} : Category one of refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics will remain part of our physical worldview: notably, the objective indefiniteness of properties, the indeterminacy of measurement results, and the tension between quantum nonlocality; locality of Relativity Theory

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **locality of Relativity Theory** ;refutation of the family of Local Realistic Theories would imply that certain peculiarities of Quantum Mechanics will remain part of our physical worldview: notably, the objective indefiniteness of properties, the indeterminacy of measurement results, and the tension between quantum nonlocality

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

No physical theory which is realistic and also local in a specified sense can agree with (=) all of the statistical implications of Quantum Mechanics.

G_{24} : Category one of **No physical theory which is realistic and also local in a specified sense can agree with**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **all of the statistical implications of Quantum Mechanics**.

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

One line of investigation in the prehistory of Bell's Theorem concerned the conjecture that the Quantum Mechanical state of a system needs (e) to be supplemented by further "elements of reality" or "hidden variables" or "complete states" in order to provide (eb) a complete description, the incompleteness of the quantum state being the explanation for (e) the statistical character of Quantum Mechanical predictions concerning the system

G_{28} : Category one of supplemented by further "elements of reality" or "hidden variables" or "complete states" in order to provide (eb) a complete description, the incompleteness of the quantum state being the explanation for (e) the statistical character of Quantum Mechanical predictions concerning the system

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of conjecture that the Quantum Mechanical state of a system

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

There are actually two main classes of hidden-variables theories. In one, which is usually called "**non-contextual",** the complete state of the system determines the value of a quantity (equivalently, an eigenvalue of the operator representing that quantity) that will be obtained by (e) any standard measuring procedure of that quantity, regardless of what other quantities are (=) simultaneously measured (eb) or what the complete

state of the system and the measuring apparatus may be.

G_{32} : Category one of called "**non-contextual "hidden variables theories ,** the complete state of the system

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of value of a quantity (equivalently, an eigenvalue of the operator representing that quantity) that will be obtained by (e) any standard measuring procedure of that quantity, regardless of what other quantities are (=) simultaneously measured (eb) or what the complete state of the system and the measuring apparatus may be

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

There are actually two main classes of hidden-variables theories. In one, which is usually called "**non-contextual",** the complete state of the system determines the value of a quantity (equivalently, an eigenvalue of the operator representing that quantity) that will be obtained by (e) any standard measuring procedure of that quantity, regardless of what other quantities are (=) simultaneously measured (eb) or what the complete state of the system and the measuring apparatus may be.

G_{36} : Category one of'**Non-contextual",** the complete state of the system determines the value of a quantity (equivalently, an eigenvalue of the operator representing that quantity) that will be obtained by (e) any standard measuring procedure of that quantity, regardless of what other quantities

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of simultaneously measured (eb) or what the complete state of the system and the measuring apparatus may be.

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

In Bohm's hidden-variables theory (1952) the context is (=) environmental, whereas in those of Bell (1966) and Gudder (1970) the context is (=) algebraic

G_{40} : Category one of context hidden-variables theory (B theories)

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of environmental

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

A pioneering version of a "hidden variables theory" was proposed by **Louis de Broglie** in 1926–7 (de Broglie 1927, 1928) and a more complete version by David Bohm in 1952 (Bohm 1952; see also the entry on Bohmian mechanics). In these theories the entity supplementing the quantum state (which is (=) a wave function in the position representation) is (=) typically a **classical entity, located in (eb) a classical phase space and therefore characterized by (e) both position and momentum variables**

G_{44} : Category one of entity supplementing the quantum state (which is (=) a wave function in the position representation)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **classical entity, located in (eb) a classical phase space and therefore characterized by (e) both position and momentum variables**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$, $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(MMMMMMMMMM) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36{,}37{,}38$ 131

(NNNNNNNNNN) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$

132

(OOOOOOOOOO) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(PPPPPPPPPP)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(QQQQQQQQQQ) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(RRRRRRRRRR) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(pp) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)}\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} +$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{13})^{(1)}t} \leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right) d\left(\left(G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13{,}14{,}15$ depend only on T_{14} and respectively on G(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$
$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $\mathrm{G}_i\,(\mathrm{t}) = 0$ and $\mathrm{T}_i\,(\mathrm{t}) = 0$ 200

it results

$\mathrm{G}_i\,(\mathrm{t}) \geq \mathrm{G}_i^0\mathrm{e}^{\left[-\int_0^{\mathrm{t}}\{(a_i')^{(2)} - (a_i'')^{(2)}(\mathrm{T}_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$\mathrm{T}_i\,(\mathrm{t}) \geq \mathrm{T}_i^0\mathrm{e}^{(-(b_i')^{(2)}\mathrm{t})} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1$, $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2$ and $\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$\mathrm{G}_{16} < (\widehat{\mathrm{M}}_{16})^{(2)}$ it follows $\frac{\mathrm{dG}_{17}}{\mathrm{dt}} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}\mathrm{G}_{17}$ and by integrating

$\mathrm{G}_{17} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2 = \mathrm{G}_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$\mathrm{G}_{18} \leq \left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_3 = \mathrm{G}_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{\mathrm{M}}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{\mathrm{t}\to\infty}((b_i'')^{(2)}\,((G_{19})(\mathrm{t}), \mathrm{t})) = (b_{17}')^{(2)}$ then $\mathrm{T}_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}}) \right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},\ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)},\ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)},\ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \leq (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t)\right) - (a_{41}'')^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(pp) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)} G_{14}^{*}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)} G_{14}^{*}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)} T_{14}^{*}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)} T_{14}^{*}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$G_{32}^{*} = \frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}$, $G_{34}^{*} = \frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$

$T_{32}^{*} = \frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}$, $T_{34}^{*} = \frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]}$ 521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$G_{36}^{*} = \frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}$, $G_{38}^{*} = \frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$

$T_{36}^{*} = \frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}$, $T_{38}^{*} = \frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$G_{40}^{*} = \frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}$, $G_{42}^{*} = \frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$

$T_{40}^{*} = \frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}$, $T_{42}^{*} = \frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$G_{44}^{*} = \frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}$, $G_{46}^{*} = \frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$

$T_{44}^{*} = \frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}$, $T_{46}^{*} = \frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\ B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWENTY NINE

Guidance Equation Ensures Agreement With The Statistical Predictions Of Standard Quantum Mechanics

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) The classical dynamics of this entity is modified by a contribution from the wave function **ψ(x, t) = R(x, t)exp[iS(x, t)/ħ]**

(2) Temporal evolution of equation (1) is governed by (e&eb) the Schrödinger Equation. Both de Broglie and Bohm assert that the velocity of the particle satisfies the "guidance equation" **(2) v = grad(S/m),**

(3) Whereby the wave function ψ acts upon (e&eb) the particles as a "guiding wave".

(4) De Broglie (1928) and the school of "Bohmian mechanics" (notably Dürr, Goldstein, and Zanghì (1992)) postulate the guidance equation without an attempt to derive it from (e) a more fundamental principle.

(5) Bohm (1952), however, proposes a deeper justification of the guidance equation. He postulates (eb) a modified version of Newton's second law of motion: **(3) m d^2x/dt^2 = −grad[V(x,t) + U(x,t)],** where V(x, t) is the standard classical potential and U(x, t) is a new entity, the "quantum potential",

(4) U(x, t) = − ($\hbar^2$/2m) $grad^2$R(x, t)/R(x, t),

and he proves that if the guidance equation holds at an initial time t0, then it follows from Eqs (2), (3), (4) and the time dependent Schrödinger Equation that it holds for all time

(6) Although Bohm deserves credit for attempting to justify the guidance equation, there is in fact tension between that equation and the modified Newtonian equation (3), which has been analyzed by Baublitz and Shimony (1996). Eq. (3) is a second order differential equation in time and does not (e) determine a definite solution for all t without two initial conditions — x and v at t0.

(7) Since v at t0 is a contingency, the validity of the guidance equation at t0 (and hence at all other times) is (=) contingent.

(8) Bohm recognizes this gap in his theory and discusses possible solutions (Bohm 1952, p. 179), without reaching a definite proposal. If, however, this difficulty is set aside, a solution is provided (eb) to the measurement problem of standard quantum mechanics, i.e., the problem of accounting for (e) the occurrence of (e) a definite outcome when the system of interest is (=) prepared in a superposition of eigenstates of the operator which is subjected (e&eb)to measurement.

(9) Furthermore, the **guidance equation ensures (eb) agreement with the statistical predictions of standard quantum mechanics.**

NOTATION

Module One

In these theories the entity supplementing the quantum state (which is (=) a wave function in the position representation) is typically a **classical entity, located in (eb) a classical phase space and therefore characterized by (e) both position and momentum variables**

G_{13} : Category one of **position and momentum variables**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of entity supplementing the quantum state (which is (=) a wave function in the position representation) is typically a **classical entity, located in (eb) a classical phase space and therefore characterized**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

classical dynamics of this entity is modified by a contribution from the wave function **ψ(x, t) = R(x, t)exp[iS(x, t)/ħ]**

G_{16} : Category one of classical dynamics of this entity is modified by a contribution from the wave function **ψ(x, t)**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of **R(x, t)exp[iS(x, t)/ħ]**

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Wave function ψ acts upon (e&eb) the particles as a "guiding wave".

G_{20} : Category one of Wave function ψ; particles as a "guiding wave".

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of particles as a "guiding wave".; Wave function ψ

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

De Broglie (1928) and the school of "Bohmian mechanics" (notably Dürr, Goldstein, and Zanghì (1992)) postulate the guidance equation without an attempt to derive it from (e) a more fundamental principle

G_{24} : Category one of more fundamental principle

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of school of "Bohmian mechanics" (notably Dürr, Goldstein, and Zanghì (1992)) postulate the guidance equation without an attempt to derive

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

$$U(x, t) = -(\hbar^2/2m)\, \mathrm{grad}^2 R(x, t)/R(x, t)$$

G_{28} : Category one of LHS of the above equation

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of RHS of the above equation

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Although Bohm deserves credit for attempting to justify the guidance equation, there is in fact tension between that equation and the modified Newtonian equation (3), which has been analyzed by Baublitz and Shimony (1996). Eq. (3) is a second order differential equation in time and does not (e) determine a definite solution for all t without two initial conditions — x and v at t0.

G_{32} : Category one of determine a definite solution for all t without two initial conditions — x and v at t0.

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of second order differential equation in time

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Since v at t0 is a contingency, the validity of the guidance equation at t0 (and hence at all other times) is (=) contingent.

G_{36} : Category one of v at t0 is a contingency, the validity of the guidance equation at t0 (and hence at all other times)

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of contingent.

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Bohm recognizes this gap in his theory and discusses possible solutions (Bohm 1952, p. 179), without reaching a definite proposal. If, however, this difficulty is set aside, a solution is provided (eb) to the measurement problem of standard quantum mechanics, i.e., the problem of accounting for (e) the occurrence of (e) a definite outcome when the system of interest is (=) prepared in a superposition of eigenstates of the

operator which is subjected (e&eb)to measurement

G_{40} : Category one of system of interest

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of prepared in a superposition of eigenstates of the operator which is subjected (e&eb)to measurement

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

system of interest is prepared in a superposition of eigenstates of the operator which is subjected (e&eb)to measurement

G_{44} : Category one of system of interest is prepared in a superposition of eigenstates of the operator; measurement

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of measurement ;system of interest is prepared in a superposition of eigenstates of the operator

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY

Existence Of Such Mappings For Which These Intuitive Interpretations Are Maintained And Conditions (1) – (4) (See Previous Module And Extant One) Are Satisfied.

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) The hidden variables model using (e) the guidance equation inspired Bell to take seriously the hidden variables interpretation of Quantum Mechanics, and the nonlocality of this model suggested (eb) his theorem.
(2) Another approach to the hidden variables conjecture has been to investigate (e&eb) the consistency of the algebraic structure of the physical quantities characterized by (e) Quantum Mechanics with a

hidden variables interpretation.

(3) Standard Quantum Mechanics assumes (eb) that the "propositions" concerning a physical system are (=) isomorphic to the lattice L(H) of closed linear subspaces of a Hilbert space H (equivalently, to the lattice of projection operators on H) with the following conditions: (1) the proposition whose truth value is (=) necessarily 'true' is matched with (=) the entire space H; (2) the proposition whose truth value is necessarily 'false' is (=) matched with the empty subspace 0; (3) if a subspace S is matched with a proposition q, then the orthogonal complement of S is matched with the negation of q; (4) the proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false', is matched with (=) the closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors which can be expressed as (eb) the sum of a vector in S1 and a vector in S2.

NOTATION

Module One

Hidden variables model using (e) the guidance equation inspired Bell to take seriously the hidden variables interpretation of Quantum Mechanics, and the nonlocality of this model suggested (eb) his theorem.

G_{13} : Category one of nonlocality of model

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of theorem.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Another approach to the hidden variables conjecture has been to investigate the consistency of the algebraic structure of the physical quantities characterized by (e) Quantum Mechanics with a hidden variables interpretation

G_{16} : Category one of Quantum Mechanics with a hidden variables interpretation

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of approach to the hidden variables conjecture has been to investigate the consistency of the algebraic structure of the physical quantities

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Standard Quantum Mechanics assumes that the "propositions" concerning a physical system are (=) isomorphic to the lattice L(H) of closed linear subspaces of a Hilbert space H (equivalently, to the lattice of projection operators on H) with the following conditions: (1) the proposition whose truth value is (=) necessarily 'true' is matched with (=) the entire space H; (2) the proposition whose truth value is necessarily

'false' is (=) matched with the empty subspace 0; (3) if a subspace S is matched with a proposition q, then the orthogonal complement of S is matched with the negation of q; (4) the proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false', is matched with (=) the closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors which can be expressed as (eb) the sum of a vector in S1 and a vector in S2.

G_{20} : Category one of Standard Quantum Mechanics

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of propositions" concerning a physical system are (=) isomorphic to the lattice L(H) of closed linear subspaces of a Hilbert space H

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Standard Quantum Mechanics assumes that the "propositions" concerning a physical system are (=) isomorphic to the lattice L(H) of closed linear subspaces of a Hilbert space H (equivalently, to the lattice of projection operators on H)

G_{24} : Category one of Standard Quantum Mechanics assumes that the "propositions" concerning a physical system

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of isomorphic to the lattice L(H) of closed linear subspaces of a Hilbert space H (equivalently, to the lattice of projection operators on H)

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

proposition whose truth value is (=) necessarily 'true' is matched with (=) the entire space H;

G_{28} : Category one of proposition whose truth value

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of necessarily 'true' is matched with the entire space H

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

proposition whose truth value is necessarily 'false' is (=) matched with the empty subspace 0;

G_{32} : Category one of proposition whose truth value is necessarily 'false'

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of matched with the empty subspace 0

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

subspace S is matched with a proposition q, then the orthogonal complement of S is matched with the negation of q

G_{36} : Category one of subspace S is matched with a proposition q, then the orthogonal complement of S ; negation of q

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of negation of q; subspace S is matched with a proposition q, then the orthogonal complement of S

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false', is matched with the closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors which can be expressed as (eb) the sum of a vector in S1 and a vector in S2.

G_{40} : Category one of proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false'; closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors which can be expressed as (eb) the sum of a vector in S1 and a vector in S2.

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors which can be expressed as (eb) the sum of a vector in S1 and a vector in S2.; proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false'

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false', is matched with the closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors which can be expressed as (eb) the sum of a vector in S1 and a vector in S2.

G_{44} : Category one of proposition q, whose truth value is 'true' if the truth-value of either q1 or q2 is 'true' and is 'false' if the answers to both q1 and q2 are 'false', is matched with the closure of the set theoretical union of the spaces S1 and S2 respectively matched to q1 and q2, the closure being the set of all vectors

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of sum of a vector in S1 and a vector in S2.

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY A

"Non-Contextual Hidden Variables Interpretation"

INTRODUCTION—VARIABLES USED

(1) It should be emphasized that this matching does not presuppose that a proposition is (=) necessarily either true or false and hence is compatible with (=) the quantum mechanical indefiniteness of a truth value, which in turn underlies (e) the feature of quantum mechanics that a physical quantity may be (=) indefinite in value.

(2) The type of hidden variables interpretation which has been most extensively treated in the literature (often called a "non-contextual hidden variables interpretation" for a reason which will soon be apparent) is a mapping m of the lattice L into (e&eb) the pair {1, 0}, where m(S) =1 intuitively means that the proposition matched with S is true and m(S) =0 means intuitively that the proposition matched with S is false.

(3) A mathematical question of importance is whether **there exist such mappings for (e) which these intuitive interpretations are maintained and conditions (1) – (4) are satisfied**.

(4) A negative answer to this question for all L (H) where the Hilbert space H has dimensionality

greater than 2 is implied by(e) a deep theorem of Gleason (1957) (which does more, by providing a complete catalogue of possible probability functions on L (H)).

(5) The same negative answer is provided much more simply by (e) John Bell (1966) (but without the complete catalogue of probability functions achieved by Gleason), who also provides (eb) a positive answer to the question in the case of dimensionality 2 ; and independently these results were also achieved by (e) Kochen and Specker (1967).

(6) It should be added that in the case of dimensionality 2 the statistical predictions of (e) any quantum state can be recovered by (eb) an appropriate mixture of the mappings m. (See also the entry on the Kochen-Specker theorem.)

NOTATION

Module One

It should be emphasized that this matching does not presuppose that a proposition is necessarily either true or false and hence is compatible with (=) the quantum mechanical indefiniteness of a truth value, which in turn underlies (e) the feature of quantum mechanics that a physical quantity may be (=) indefinite in value

G_{13} : Category one of matching does not presuppose that a proposition

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of necessarily either true or false and hence is compatible with (=) the quantum mechanical indefiniteness of a truth value, which in turn underlies (e) the feature of quantum mechanics that a physical quantity may be (=) indefinite in value

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

matching does not presuppose that a proposition is necessarily either true or false and hence is compatible with the quantum mechanical indefiniteness of a truth value, which in turn underlies (e) the feature of quantum mechanics that a physical quantity may be (=) indefinite in value

G_{16} : Category one of matching does not presuppose that a proposition is necessarily either true or false and hence is compatible

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of quantum mechanical indefiniteness of a truth value, which in turn underlies (e) the feature of quantum mechanics that a physical quantity may be (=) indefinite in value

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

matching does not presuppose that a proposition is necessarily either true or false and hence is compatible with the quantum mechanical indefiniteness of a truth value, which in turn underlies the feature of quantum mechanics that a physical quantity may be (=) indefinite in value

G_{20} : Category one of feature of quantum mechanics that a physical quantity may be (=) indefinite in value

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of matching does not presuppose that a proposition is necessarily either true or false and hence is compatible with the quantum mechanical indefiniteness of a truth value, which in turn underlies

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

matching does not presuppose that a proposition is necessarily either true or false and hence is compatible with the quantum mechanical indefiniteness of a truth value, which in turn underlies the feature of quantum mechanics that a physical quantity may be (=) indefinite in value

G_{24} : Category one of matching does not presuppose that a proposition is necessarily either true or false and hence is compatible with the quantum mechanical indefiniteness of a truth value, which in turn underlies the feature of quantum mechanics that a physical quantity

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of indefinite in value

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

type of hidden variables interpretation which has been most extensively treated in the literature (often called a “non-contextual hidden variables interpretation” for a reason which will soon be apparent) is a mapping m of the lattice L into (e&eb) the pair {1, 0}, where m(S) =1 intuitively means that the proposition matched with S is true and m(S) =0 means intuitively that the proposition matched with S is false

G_{28} : Category one of type of hidden variables interpretation which has been most extensively treated in the literature (often called a “non-contextual hidden variables interpretation” for a reason which will soon be apparent) is a mapping m of the lattice L ; pair {1, 0}, where m(S) =1 intuitively means that the proposition matched with S is true and m(S) =0 means intuitively that the proposition matched with S is false

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of pair {1, 0}, where m(S) =1 intuitively means that the proposition matched with S is true and m(S) =0 means intuitively that the proposition matched with S is false ;type of hidden variables interpretation which has been most extensively treated in the literature (often called a “non-contextual

hidden variables interpretation" for a reason which will soon be apparent) is a mapping m of the lattice L

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

mathematical question of importance is whether **there exist such mappings for (e) which these intuitive interpretations are maintained and conditions (1) – (4) are satisfied**

G_{32} : Category one of **which these intuitive interpretations are maintained and conditions (1) – (4) are satisfied**

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of **there exist such mappings**

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Negative answer to this question for all L (H) where the Hilbert space H has dimensionality greater than 2 is implied by (e) a deep theorem of Gleason (1957) (which does more, by providing a complete catalogue of possible probability functions on L (H))

G_{36} : Category one of theorem of Gleason (1957) (which does more, by providing a complete catalogue of possible probability functions on L (H)). **Note that we are here talking of the systems that satisfy the conditionalities of the Gleason Theorem. Characterstics of such systems are taken in to consideration in the classification scheme.**

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Negative answer to this question for all L (H) where the Hilbert space H has dimensionality greater than 2

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

same negative answer is provided much more simply by John Bell (1966) (but without the complete catalogue of probability functions achieved by Gleason), who also provides (eb) a positive answer to the question in the case of dimensionality 2 ; and independently these results were also achieved by (e) Kochen and Specker (1967)

G_{40} : Category one of John Bell (1966) (but without the complete catalogue of probability functions achieved by Gleason), who also provides (eb) a positive answer to the question in the case of dimensionality 2 ; and independently these results were also achieved by (e) Kochen and Specker (1967)

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of negative answer

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

same negative answer is provided much more simply by John Bell (1966) (but without the complete catalogue of probability functions achieved by Gleason), who also provides (eb) a positive answer to the question in the case of dimensionality 2

G_{44} : Category one of same negative answer is provided much more simply by John Bell (1966) (but without the complete catalogue of probability functions achieved by Gleason)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of positive answer to the question in the case of dimensionality 2

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY ONE

Truth Value Which M Assigns To A Subspace S Depends Upon The Context C Of Propositions Measured In Tandem With The One Associated With S

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) In Bell (1966), after presenting a strong case against the hidden variables program (except for the special case of dimensionality 2) Bell performs a dramatic reversal by introducing (eb) a new type of hidden variables interpretation – one in which the **truth value which m assigns to (e) a subspace S depends upon the context C of propositions measured in (e&eb) tandem with the one associated with S.**

(2) In the new type of hidden variables interpretation the truth value into which m maps (e&eb) S is (=) depends upon the context C.

(3) These interpretations are (=) commonly referred to as "contextual hidden variables interpretations", whereas those in which there is no dependence upon the context are (=) called "non-contextual."

(4) Bell proves the consistency of contextual hidden variables interpretations with (e&eb) the algebraic structure of the lattice L (H) for two examples of H with dimension greater than 2. (His proposal has been systematized (e&eb) by Gudder (1970), who takes a context C to be (=) a maximal Boolean subalgebra of the lattice L (H) of subspaces.)

(5) Another line leading to Bell's Theorem was the investigation of Quantum Mechanically entangled states, that is, quantum states of a composite system that cannot be (e) expressed as direct products of quantum states of the individual components.

(6) That **Quantum Mechanics admits (e) of such entangled states was discovered by Erwin Schrödinger (1926) in one of his pioneering papers, but the significance of this discovery was not emphasized until the paper of Einstein, Podolsky, and Rosen (1935).**

(7) They examined correlations between the positions and (e&eb) the linear momenta of two well separated spinless particles and concluded that (eb) in order to avoid (e) an appeal to nonlocality these correlations could only be explained by (e) "elements of physical reality" in each particle — specifically, both definite position and definite momentum — and since this description is (=) richer than permitted by the uncertainty principle of Quantum Mechanics their conclusion is (=) effectively an argument for a hidden variables interpretation.

(8) It should be emphasized that their argument does not (e) depend upon counter-factual reasoning, which is (=) reasoning about what would be observed if a quantity were measured (eb) other than the one that was in fact measured; instead their argument can be reformulated entirely in the ordinary inductive logic, as emphasized by d'Espagnat (1976) and Shimony (2001).

(9) This reformulation is important because (e) it diminishes the force of Bohr's (1935) rebuttal of Einstein, Podolsky, and Rosen on the grounds that one is (=) not entitled to draw conclusions about the existence of elements of physical reality from (e) considerations of what would be seen if a measurement (eb) other than the actual one had been performed.

NOTATION

Module One

In Bell (1966), after presenting a strong case against the hidden variables program (except for the special case of dimensionality 2) Bell performs a dramatic reversal by introducing a new type of hidden variables interpretation – one in which the **truth value which m assigns to (e) a subspace S depends upon the context C of propositions measured in (e&eb) tandem with the one associated with S.**

G_{13} : Category one of dramatic reversal is performed by introduction

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of new type of hidden variables interpretation – one in which the **truth value which m assigns to (e) a subspace S depends upon the context C of propositions measured in (e&eb) tandem with the one associated with S.**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

dramatic reversal is by introducing a new type of hidden variables interpretation – one in which the **truth value which m assigns to (e) a subspace S depends upon the context C of propositions measured in**

(e&eb) tandem with the one associated with S.

G_{16} : Category one of **subspace S depends upon the context C of propositions measured in tandem with the one associated with S.**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of dramatic reversal is by introducing a new type of hidden variables interpretation – one in which the **truth value which m assigns**

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

New type of hidden variables interpretation has the truth value into which m maps (e&eb) S is dependent upon the context C.

G_{20} : Category one of New type of hidden variables interpretation has the truth value into which m maps; S is dependent upon the context C.

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of S is dependent upon the context C.; New type of hidden variables interpretation has the truth value into which m maps

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Bell proves the consistency of contextual hidden variables interpretations with (e&eb) the algebraic structure of the lattice L (H) for two examples of H with dimension greater than 2. (His proposal has been systematized (e&eb) by Gudder (1970), who takes a context C to be (=) a maximal Boolean subalgebra of the lattice L (H) of subspaces.)

G_{24} : Category one of consistency of contextual hidden variables interpretations; algebraic structure of the lattice L (H) for two examples of H with dimension greater than 2. (His proposal has been systematized (e&eb) by Gudder (1970), who takes a context C to be (=) a maximal Boolean subalgebra of the lattice L (H) of subspaces.)

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of algebraic structure of the lattice L (H) for two examples of H with dimension greater than 2. (His proposal has been systematized (e&eb) by Gudder (1970), who takes a context C to be (=) a maximal Boolean subalgebra of the lattice L (H) of subspaces.); consistency of contextual hidden variables interpretations

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

quantum states of a composite system that cannot be (e) expressed as direct products of quantum states of the individual components

G_{28} : Category one of expressed as direct products of quantum states of the individual components

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of quantum states of a composite system

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Quantum Mechanics admits (e) of such entangled states was discovered by Erwin Schrödinger (1926) in one of his pioneering papers, but the significance of this discovery was not emphasized until the paper of Einstein, Podolsky, and Rosen (1935)

G_{32} : Category one of **entangled states**

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of **Quantum Mechanics**

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

They examined correlations between the positions and the linear momenta of two well separated spinless particles and concluded that in order to avoid an appeal to nonlocality these correlations could only be explained by "elements of physical reality" in each particle — specifically, both definite position and definite momentum — and since this description is (=) richer than permitted by the uncertainty principle of Quantum Mechanics their conclusion is (=) effectively an argument for a hidden variables interpretation.

G_{36} : Category one of "elements of physical reality" in each particle — specifically, both definite position and definite momentum — and since this description is (=) richer than permitted by the uncertainty principle of Quantum Mechanics their conclusion is (=) effectively an argument for a hidden variables interpretation

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of correlations between the positions and the linear momenta of two well separated spinless particles and concluded that in order to avoid an appeal to nonlocality these correlations

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

argument does not depend upon counter-factual reasoning, which is reasoning about what would be observed if a quantity were measured (eb) other than the one that was in fact measured; instead their argument can be reformulated entirely in the ordinary inductive logic, as emphasized by d'Espagnat (1976) and Shimony (2001)

G_{40} : Category one of argument does not depend upon counter-factual reasoning

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of reasoning about what would be observed if a quantity were measured (eb) other than the one that was in fact measured; instead their argument can be reformulated entirely in the ordinary inductive logic, as emphasized by d'Espagnat (1976) and Shimony (2001)

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

argument does not depend upon counter-factual reasoning, which is reasoning about what would be observed if a quantity were measured (eb) other than the one that was in fact measured; instead their argument can be reformulated entirely in the ordinary inductive logic, as emphasized by d'Espagnat (1976) and Shimony (2001)

G_{44} : Category one of argument does not depend upon counter-factual reasoning, which is reasoning about what would be observed if a quantity

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of other than the one that was in fact measured; instead their argument can be reformulated entirely in the ordinary inductive logic, as emphasized by d'Espagnat (1976) and Shimony (2001)

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},\ \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ ,$
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor
$-(b''_{16})^{(2)}((G_{19}),t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$
$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}), t\right)\right]T_{45} \quad 53$$
$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}), t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}), t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) \quad +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) \quad +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) \quad +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{array}\right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)}\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20} \quad (70)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21} \quad (71)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22} \quad (72)$$

$\boxed{-(b_{20}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{16}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{13}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,)}(G,t)}$, $\boxed{-(b_{15}'')^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b_{36}'')^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b_{38}'')^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{24}')^{(4)} & \boxed{+(a_{24}'')^{(4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a_{25}')^{(4)} & \boxed{+(a_{25}'')^{(4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{33}'')^{(6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a_{26}')^{(4)} & \boxed{+(a_{26}'')^{(4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)} \quad \boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9)}(T_{45},t)} \end{array}\right] G_{26} \quad (75)$$

$\boxed{(a_{24}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{25}'')^{(4)}(T_{25},t)}$, $\boxed{(a_{26}'')^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a_{28}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a_{32}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a_{13}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{14}'')^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1)}(T_{14},t)}$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a_{16}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2)}(T_{17},t)}$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a_{20}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3)}(T_{21},t)}$

are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$

are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \tag{76}$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \tag{77}$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{26} \tag{78}$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt}=(a_{28})^{(5)}G_{29}-\begin{bmatrix}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{28} \tag{79}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad 81$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients*

$for\ category$ 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ $are\ \ third\ \ detrition\ coefficients$ $for\ category$ 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34} \quad (87)$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ $are\ \ first\ \ augmentation\ coefficients$ $for\ category$ 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ $are\ second\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ $are\ third\ augmentation$ $coefficients\ for\ category$ 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$
seventh augmentation coefficients
$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$
Eighth augmentation coefficients
$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt}=(b_{32})^{(6)}T_{33}-\begin{bmatrix}(b'_{32})^{(6)}\,-(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t)\\ -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{32} \quad (88)$$

$$\frac{dT_{33}}{dt}=(b_{33})^{(6)}T_{32}-\begin{bmatrix}(b'_{33})^{(6)}\,-(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t)\\ -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{33} \quad (89)$$

$$\frac{dT_{34}}{dt}=(b_{34})^{(6)}T_{33}-\begin{bmatrix}(b'_{34})^{(6)}\,-(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t)\\ -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)\\ -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{34} \quad (90)$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3
$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3
$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3
$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3
$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

96

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second

augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3
$+(a_{38}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{37}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a_{36}'')^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a_{40}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b_{44}')^{(9)}\left[-(b_{44}'')^{(9)}(G_{47},t)\right] & \left[-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b_{45}')^{(9)}\left[-(b_{45}'')^{(9)}(G_{47},t)\right] & \left[-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b_{46}')^{(9)}\left[-(b_{46}'')^{(9)}(G_{47},t)\right] & \left[-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)},\ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} +\ (\hat{A}_{20})^{(3)} +\ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} +\ (\hat{B}_{20})^{(3)} +\ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le\ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) =\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\left((G_{35}), t\right) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(SSSSSSSSSS) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

(TTTTTTTTTT) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$

132

(UUUUUUUUUU) $\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(VVVVVVVVV)

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(WWWWWWWWWW) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(XXXXXXXXXX) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be

noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+\ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+\ (\hat{Q}_{44})^{(9)}\ \ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t)\leq\ \left(\hat{P}_{13}\right)^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\leq\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\ ,T_i(0)$

$$G_i(t)\leq\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\leq\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\leq\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\ ,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{\left(\hat{M}_{40}\right)^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{13})^{(1)}, T_i^0 \le (\hat{Q}_{13})^{(1)}$, 155

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 156

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\tilde{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad 169$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(qq) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left(e^{(\tilde{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\tilde{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left(e^{(\tilde{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\tilde{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left(e^{(\tilde{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\tilde{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\tilde{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\tilde{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\tilde{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[(\widehat{P}_{13})^{(1)}+\left((\hat{P}_{13})^{(1)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on

$G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 19 to 24 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(1)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1 / (a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2 / (a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)}+\left((\hat{P}_{16})^{(2)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{16})^{(2)}\right]\leq(\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on

(G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)} t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18}. indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16}, G_{18} and G_{16}, G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18}. The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2:

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\,(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on

$(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on

$(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 228

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(4)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\,((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}\,(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\,((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})},\widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})},\widetilde{(T_{35})}\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose

$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)}+\left((\hat{P}_{36})^{(7)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\,(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it

suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1 / (a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2 / (a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 274

$(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t), t(t), t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t}\le G_{15}(t)\le\right. \quad (286)$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad (287)$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad (288)$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad (289)$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- (290)

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation (291)

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: (292)

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(T_{17},t)+(a''_{17})^{(2)}(T_{17},t)\le-(\sigma_1)^{(2)} \quad (293)$$

$$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)} \quad (294)$$

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: (295)

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots (296)

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (297)

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and (298)

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: (299)

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the (300)

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ (301)

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}$, ***if*** $(v_0)^{(4)} < (v_1)^{(4)}$

$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}$, ***if*** $(v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)}$,
and $\boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}$, ***if*** $(\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)}G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, $\boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}$, ***if*** $(v_0)^{(7)} < (v_1)^{(7)}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}$, ***if*** $(v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)}$,

and $\boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big) - (b''_{45})^{(9)}\big((G_{47}),t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t})$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^{0}}{G_{14}^{0}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)}+(C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)}-(v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)}-(v_0)^{(1)}}{(v_0)^{(1)}-(v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}$, $\boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}$, $\boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25},t)\right) - (a_{25}'')^{(4)}(T_{25},t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then
$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

411

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \le u^{(6)}(t) \le (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines** $(v_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)}=\frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)\leq\frac{dv^{(7)}}{dt}\leq-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0<\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}<(\nu_1)^{(7)}<(\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t)\geq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_0)^{(7)}\right)t\right]}}\quad,\quad\boxed{(C)^{(7)}=\frac{(\nu_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(7)}=\frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)}\leq\nu^{(7)}(t)\leq(\bar{\nu}_1)^{(7)}$

If $0<(\nu_1)^{(7)}<(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}<(\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)}\leq\frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}\leq\nu^{(7)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\bar{\nu}_1)^{(7)}$$

If $0<(\nu_1)^{(7)}\leq(\bar{\nu}_1)^{(7)}\leq\boxed{(\nu_0)^{(7)}=\frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)}\leq\nu^{(7)}(t)\leq\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}\leq(\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)}\leq\nu^{(7)}(t)\leq(m_1)^{(7)},\quad\boxed{\nu^{(7)}(t)=\frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for**

the special case .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(v_1)^{(9)}$ and $(\bar{v}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(qq) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that

there exists a unique G_{37}^{*} such that $\varphi(G_{39}{}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^{*} such that $\varphi(G_{43}{}^{*}) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^{*} such that $\varphi((G_{47})^{*}) = 0$

Finally we obtain the unique solution 511

G_{14}^{*} given by $\varphi(G^{*}) = 0$, T_{14}^{*} given by $f(T_{14}^{*}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \quad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i\qquad,T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^{*}\,)=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt}=-\big((a'_{13})^{(1)}+(p_{13})^{(1)}\big)\mathbb{G}_{13}+(a_{13})^{(1)}\mathbb{G}_{14}-(q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt}=-\big((a'_{14})^{(1)}+(p_{14})^{(1)}\big)\mathbb{G}_{14}+(a_{14})^{(1)}\mathbb{G}_{13}-(q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt}=-\big((a'_{15})^{(1)}+(p_{15})^{(1)}\big)\mathbb{G}_{15}+(a_{15})^{(1)}\mathbb{G}_{14}-(q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt}=-\big((b'_{13})^{(1)}-(r_{13})^{(1)}\big)\mathbb{T}_{13}+(b_{13})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt}=-\big((b'_{14})^{(1)}-(r_{14})^{(1)}\big)\mathbb{T}_{14}+(b_{14})^{(1)}\mathbb{T}_{13}+\textstyle\sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt}=-\big((b'_{15})^{(1)}-(r_{15})^{(1)}\big)\mathbb{T}_{15}+(b_{15})^{(1)}\mathbb{T}_{14}+\textstyle\sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^{*}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^{*}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^{*}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^{*}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^{*}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^{*}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^{*}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^{*}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^{*}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}\ and\ (b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \ , \ \frac{\partial(b_i'')^{(9)}}{\partial G_j}(\ (G_{47})^*\ ) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\ C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\ D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\ E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\ F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\ G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY TWO

Outcome Of A Measurement Made On One Constituent Depends Upon The Action Of The "Guiding Wave" Upon Constituents That Are Far Off, Which In General Will Depend On The Measurement Arrangement On That Side

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) **Bell was skeptical of Bohr's rebuttal for other reasons, essentially that he regarded it to be (=) anthropocentric**.

(2) At the conclusion of Bell (1966), in which Bell gives a new lease on life to the hidden variables program by introducing (eb) the notion of contextual hidden variables, he performs another dramatic reversal by raising (eb) a question about a composite system consisting of two well separated particles 1 and 2:

(3) Suppose a proposition associated with a (e&eb) subspace S1 of the Hilbert space of particle 1 is subjected to (e&eb) measurement, and a contextual hidden variables theory assigns (eb) the truth valuem (S1/C) to this proposition.

(4) What physically reasonable conditions can be imposed upon (e&eb) the context C?

(5) Bell suggests that C should consist (e) only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will (=) depend upon what operations are performed upon (e&eb) a remote particle 2, and that would be (=) non-locality.

(6) This condition raises the question: can the statistical predictions of Quantum Mechanics concerning the entangled state be duplicated by (e) a contextual hidden variables theory in which the context C is (=) localized?

(7) It is interesting to note that this question also arises from (e) a consideration of the de Broglie-Bohm model

(8) when Bohm derives the statistical predictions of a Quantum Mechanically entangled system whose constituents are well separated, the **outcome of a measurement made on one constituent depends upon (=) the action of the "guiding wave" upon constituents that are far off, which in general will depend on (=)the measurement arrangement on that side.**

(9) Bell was thus heuristically led to ask whether a lapse of locality is (=) necessary for the recovery of

Quantum Mechanical statistics.

NOTATION

Module One

Bell was skeptical of Bohr's rebuttal for other reasons, essentially that he regarded it to be (=) anthropocentric.

G_{13} : Category one of **rebuttal was for other reasons, essentially it was regarded theorem to be**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **anthropocentric**.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Dual module

Bell gives a new lease on life to the hidden variables program by introducing the notion of contextual hidden variables, he performs another dramatic reversal by raising (eb) a question about a composite system consisting of two well separated particles 1 and 2

G_{16} : Category one of hidden variables program; hidden variables program by introducing the notion of contextual hidden variables, he performs another dramatic reversal by raising

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of notion of contextual hidden variables, he performs another dramatic reversal by raising (eb) a question about a composite system consisting of two well separated particles 1 and 2; question about a composite system consisting of two well separated particles 1 and 2

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Suppose a proposition associated with a (e&eb) subspace S1 of the Hilbert space of particle 1 is subjected to measurement, and a contextual hidden variables theory assigns (eb) the truth valuem (S1/C) to this proposition

G_{20} : Category one of proposition; subspace S1 of the Hilbert space of particle 1

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of subspace S1 of the Hilbert space of particle 1 ;proposition

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

subspace S1 of the Hilbert space of particle 1 is subjected to measurement, and a contextual hidden variables theory assigns (eb) the truth valuem (S1/C) to this proposition

G_{24} : Category one of subspace S1 of the Hilbert space of particle 1; measurement, and a contextual hidden variables theory assigns (eb) the truth valuem (S1/C) to this proposition

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of measurement, and a contextual hidden variables theory assigns (eb) the truth valuem (S1/C) to this proposition ;subspace S1 of the Hilbert space of particle 1

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

subspace S1 of the Hilbert space of particle 1 is subjected to measurement, and a contextual hidden variables theory assigns (eb) the truth valuem (S1/C) to this proposition

G_{28} : Category one of subspace S1 of the Hilbert space of particle 1 is subjected to measurement, and a contextual hidden variables theory

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of truth valuem (S1/C) to this proposition

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

he performs another dramatic reversal by raising (eb) a question about a composite system consisting of two well separated particles 1 and 2

G_{32} : Category one of two well separated particles 1 and 2

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of question about a composite system

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

What physically reasonable conditions can be imposed upon (e&eb) the context C?

G_{36} : Category one of physically reasonable conditions; context C

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of context C ;physically reasonable conditions

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Bell suggests that C should consist only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will (=) depend upon what operations are performed upon (e&eb) a remote particle 2, and that would be (=) non-locality

G_{40} : Category one of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will (=) depend upon what operations are performed upon (e&eb) a remote particle 2, and that would be (=) non-locality

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of C

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

C should consist only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will depend upon what operations are performed upon a remote particle 2, and that would be (=) non-locality

G_{44} : Category one of C should consist only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1; C should consist only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will depend upon what operations; remote particle 2, and that would be (=) non-locality; C should consist only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will depend upon what operations are performed upon a remote particle 2

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of depend upon what operations are performed upon (e&eb) a remote particle 2, and that would be (=) non-locality; remote particle 2, and that would be (=) non-locality; C should consist only of propositions concerning particle 1, for otherwise the outcome of the measurement upon 1 will depend upon what operations; non-locality

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY THREE

Violation Constitutes A Special Case Of Bell's Theorem For It Shows That No Local Realistic Theory Subsumed Under The Framework Of Bell's 1964 Paper Can Agree With All Of The Statistical Predictions Of Quantum Theory.

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

Proof of a Theorem of Bell's Type

(1) Bell (1964) gives a pioneering proof of the theorem that bears his name, by first making explicit a conceptual framework within which expectation values can be calculated for (e) the spin components of a pair of spin-half particles, then showing that (eb) regardless of the choices that are (=) made for certain unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality."

(2) That term is now commonly used (e) to denote collectively a family of Inequalities derived in (eb) conceptual frameworks similar to but more general than the original one of Bell.

(3) Sometimes these Inequalities are referred to as "Inequalities of Bell's type." Each of these conceptual frameworks incorporates (e) some type of hidden variables theory and obeys (eb) a locality assumption.

NOTATION

Module One

Bell (1964) gives a pioneering proof of the theorem that bears his name, by first making explicit a conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that (eb) regardless of the choices that are (=) made for certain unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality."

G_{13} : Category one of spin components of a pair of spin-half particles, then showing that (eb) regardless of the choices that are (=) made for certain unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality."

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of conceptual framework within which expectation values can be calculated

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices that are (=) made for certain unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality."

G_{16} : Category one of conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of regardless of the choices that are (=) made for certain unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices that are made for certain unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality."

G_{20} : Category one of unspecified functions that occur in (eb) the framework, the expectation values obey (eb) a certain inequality which has come to be called "Bell's Inequality."

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices that are made for certain unspecified functions that occur in the framework, the expectation values obey (eb) a certain inequality which has come

to be called “Bell's Inequality.”

G_{24} : Category one of conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices that are made for certain unspecified functions that occur

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of framework, the expectation values obey (eb) a certain inequality which has come to be called “Bell's Inequality.”

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices that are made for certain unspecified functions that occur in the framework, the expectation values obey a certain inequality which has come to be called “Bell's Inequality.”

G_{28} : Category one of conceptual framework within which expectation values can be calculated for the spin components of a pair of spin-half particles, then showing that regardless of the choices that are made for certain unspecified functions that occur in the framework, the expectation values

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of certain inequality which has come to be called “Bell's Inequality.”

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Term is now commonly used to denote collectively a family of Inequalities derived in (eb) conceptual frameworks similar to but more general than the original one of Bell.

G_{32} : Category one of denote collectively a family of Inequalities derived in (eb) conceptual frameworks similar to but more general than the original one of Bell.

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Term

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Term is now commonly used to denote collectively a family of Inequalities derived in conceptual frameworks similar to but more general than the original one of Bell.

G_{36} : Category one of Term is now commonly used to denote collectively a family of Inequalities

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of conceptual frameworks similar to but more general than the original one of Bell.

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Sometimes these Inequalities are referred to as "Inequalities of Bell's type." Each of these conceptual frameworks incorporates some type of hidden variables theory and obeys (eb) a locality assumption

G_{40} : Category one of some type of hidden variables theory and obeys (eb) a locality assumption

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of Each of these conceptual frameworks

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Sometimes these Inequalities are referred to as "Inequalities of Bell's type." Each of these conceptual frameworks incorporates some type of hidden variables theory and obeys (eb) a locality assumption

G_{44} : Category one of Each of these conceptual frameworks incorporates some type of hidden variables theory and obeys

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of locality assumption

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY THREE A

"Local Realistic Theory"

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

Proof of a Theorem of Bell's Type

(1) The name "Local Realistic Theory" is also appropriate and will be used throughout this article because of its generality. Bell calculates the expectation values for (e) certain products of the form (σ1·â) (σ2·ê), where σ1 is (=) the vectorial Pauli spin operator for (e) particle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality.

NOTATION

Module One

"Local Realistic Theory" is also appropriate and will be used throughout this article because of its generality. Bell calculates the expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is (=) the vectorial Pauli spin operator for (e) particle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality.

G_{13} : Category one of certain products of the form (σ1·â) (σ2·ê), where σ1 is (=) the vectorial Pauli spin operator for (e) particle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of expectation values

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for (e) particle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{16} : Category one of expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of vectorial Pauli spin operator for (e) particle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{20} : Category one ofparticle 1 and σ2 is the vectorial Pauli spin operator for (e) particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{24} : Category one of particle 2 (both particles having spin ½), and â and ê are (=) unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin

½), and â and ê are unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{28} : Category one of expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin ½), and â and ê

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of unit vectors in three-space, and then shows that (eb) these Quantum Mechanical expectation values violate (e) Bell's Inequality

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin ½), and â and ê are unit vectors in three-space, and then shows that these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{32} : Category one of expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin ½), and â and ê are unit vectors in three-space

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of Quantum Mechanical expectation values violate (e) Bell's Inequality

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin ½), and â and ê are unit vectors in three-space, and then shows that these Quantum Mechanical expectation values violate (e) Bell's Inequality

G_{36} : Category one of Bell's Inequality (There is only one Bell's inequality. We are talking of systems which satisfy Bell's inequality and the characterstics thereof, for the classification scheme. On the other hand, it is possible to take all the categories as equivalent. Both the methods are feasible)

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of expectation values for certain products of the form (σ1·â) (σ2·ê), where σ1 is the vectorial Pauli spin operator for particle 1 and σ2 is the vectorial Pauli spin operator for particle 2 (both particles having spin ½), and â and ê are unit vectors in three-space, and then shows that these Quantum Mechanical expectation values

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

This **violation constitutes a special case of Bell's Theorem for it shows (eb) that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory**

G_{40} : Category one of **special case of Bell's Theorem for it shows (eb) that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory**

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of **violation**

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

violation constitutes a special case of Bell's Theorem for it shows (eb) that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory

G_{44} : Category one of **violation constitutes a special case of Bell's Theorem**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY THREE B

Each Pair Of Systems Is Characterized By A "Complete State" M Which Contains The Entirety Of The Properties Of The Pair At The Moment Of Generation

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

Proof of a Theorem of Bell's Type

(1) This **violation constitutes (e) a special case of Bell's Theorem for it shows (eb) that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory.**

(2) In the present section the pattern of Bell's 1964 paper will be followed: **formulation of (e) a framework, derivation of (e) an Inequality, demonstration of (e) a discrepancy between certain quantum mechanical expectation values and (e&eb) this Inequality.**

(3) However, a more general conceptual framework than his will be assumed (eb) and a somewhat more general Inequality will be derived, thus yielding (eb) a more general theorem than the one derived by Bell in 1964, but with the same strategy and in the same spirit. Papers which took the steps from Bell's 1964 demonstration to the one given here are Clauser et al. (1969), Bell (1971), Clauser and Horne (1974), Aspect (1983) and Mermin (1986).

(4) The conceptual framework in which a Bell-type Inequality will be demonstrated first of all postulates (eb) an ensemble of pairs of systems, the individual systems in each pair being labeled as 1 and 2.

(5) Each pair of systems is (=) characterized by a "complete state" m which contains (e) the entirety of the properties of the pair at the moment of generation.

NOTATION

Module One

violation constitutes a special case of Bell's Theorem for it shows (eb) that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory

G_{13} : Category one of **special case of Bell's Theorem for it shows (eb) that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with (=) all of the statistical predictions of quantum theory**

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of **violation**

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

violation constitutes a special case of Bell's Theorem for it shows that no Local Realistic Theory subsumed under the framework of Bell's 1964 paper can agree with all of the statistical predictions of

quantum theory

G_{16} : Category one of **violation constitutes a special case of Bell's Theorem for it shows that no Local Realistic Theory**

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of **all of the statistical predictions of quantum theory**

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Formulation of a framework, derivation of an Inequality, demonstration of (e) a discrepancy between certain quantum mechanical expectation values and (e&eb) this Inequality.

G_{20} : Category one of formulation of a framework; derivation of an Inequality, demonstration of (e) a discrepancy between certain quantum mechanical expectation values and (e&eb) this Inequality**.**

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of formulation of a framework; certain quantum mechanical expectation values and (e&eb) this Inequality

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Derivation of an Inequality, demonstration of a discrepancy between certain quantum mechanical expectation values and (e&eb) this Inequality.

G_{24} : Category one of **Derivation of an Inequality; certain quantum mechanical expectation values and (e&eb) this Inequality.**

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of **certain quantum mechanical expectation values and (e&eb) this Inequality.; Derivation of an Inequality**

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

demonstration of a discrepancy between certain quantum mechanical expectation values and (e&eb) this Inequality

G_{28} : Category one of Bell's inequality; demonstration of a discrepancy between certain quantum mechanical expectation values

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of demonstration of a discrepancy between certain quantum mechanical expectation values; Bell's inequality

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

general conceptual framework than his will be assumed and a somewhat more general Inequality will be derived, thus yielding (eb) a more general theorem than the one derived by Bell in 1964, but with the same strategy and in the same spirit. Papers which took the steps from Bell's 1964 demonstration to the one given here are Clauser et al. (1969), Bell (1971), Clauser and Horne (1974), Aspect (1983) and Mermin (1986).

G_{32} : Category one of general conceptual framework

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of somewhat more general Inequality will be derived, thus yielding (eb) a more general theorem than the one derived by Bell in 1964,

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

general conceptual framework than his will be assumed and a somewhat more general Inequality will be derived, thus yielding (eb) a more general theorem than the one derived by Bell in 1964,

G_{36} : Category one of general conceptual framework than his will be assumed and a somewhat more general Inequality will be derived

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of more general theorem

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Conceptual framework in which a Bell-type Inequality will be demonstrated first of all postulates (eb) an ensemble of pairs of systems, the individual systems in each pair being labeled as 1 and 2.

G_{40} : Category one of Conceptual framework in which a Bell-type Inequality

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of ensemble of pairs of systems, the individual systems in each pair being labeled as 1 and 2.

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Each pair of systems is characterized by a "complete state" m which contains (e) the entirety of the properties of the pair at the moment of generation

G_{44} : Category one of **entirety of the properties of the pair at the moment of generation**

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of **Each pair of systems is characterized by a "complete state" m**

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY FOUR

Bell's Own Version Of His Theorem Assumed That S And T Are Both Bivalent

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) Different experiments can be performed on (eb) each system, those on (eb) 1 designated by a, a′, etc. and those on 2 by (e) b, b′, etc.

(2) One can in principle let a, a′, etc. also include (e) characteristics of the apparatus used for (e) the measurement, but since the dependence of the result upon (eb) the microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc.

(3) The remarkably good agreement between the experimental measurements of correlations in Bell-type experiments and (=) the quantum mechanical predictions of these correlations justifies (eb) restricting attention in practice to (e) macroscopic features of the apparatus.

NOTATION

Module One

Different experiments can be performed on each system, those on (eb) 1 designated by a, a′, etc. and those on 2 by (e) b, b′, etc.

G_{13} : Category one of Different experiments can be performed

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of each system, those on (eb) 1 designated by a, a′, etc. and those on 2 by (e) b, b′, etc.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Different experiments can be performed on each system, those on (eb) 1 designated by a, a′, etc. and those on 2 by (e) b, b′, etc.

G_{16} : Category one of Different experiments can be performed on each system, those

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of 1 designated by a, a′, etc. and those on 2 by (e) b, b′, etc.

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

One can in principle let a, a′, etc. also include characteristics of the apparatus used for (e) the measurement, but since the dependence of the result upon (eb) the microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc.

G_{20} : Category one of characteristics of the apparatus used for (e) the measurement, but since the dependence of the result upon (eb) the microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc.

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of let a, a′, etc. also include

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence of the result upon (eb) the microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

G_{24} : Category one of measurement, but since the dependence of the result upon (eb) the microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of let a, a′, etc. also include characteristics of the apparatus used

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence of the result upon the microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

G_{28} : Category one of let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence of the result

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of microscopic features of the apparatus is (=) not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence

of the result upon the microscopic features of the apparatus is not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

G_{32} : Category one of let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence of the result upon the microscopic features of the apparatus

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of indeterminable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence of the result upon the microscopic features of the apparatus is not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted (e) in practice in the descriptions a, a′, etc., and likewise for b, b′, etc

G_{36} : Category one of descriptions a, a′, etc., and likewise for b, b′, etc

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of let a, a′, etc. also include characteristics of the apparatus used for the measurement, but since the dependence of the result upon the microscopic features of the apparatus is not determinable experimentally, only the macroscopic features of the apparatus (such as orientations of the polarization analyzers) in their incompletely controllable environment need be admitted

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

remarkably good agreement between the experimental measurements of correlations in Bell-type experiments and the quantum mechanical predictions of these correlations justifies restricting attention in practice to (e) macroscopic features of the apparatus

G_{40} : Category one of remarkably good agreement between the experimental measurements of correlations in Bell-type experiments and the quantum mechanical predictions of these correlations

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of restricting attention in practice to (e) macroscopic features of the apparatus

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

remarkably good agreement between the experimental measurements of correlations in Bell-type experiments and the quantum mechanical predictions of these correlations justifies restricting attention in practice to (e) macroscopic features of the apparatus

G_{44} : Category one of macroscopic features of the apparatus

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of remarkably good agreement between the experimental measurements of correlations in Bell-type experiments and the quantum mechanical predictions of these correlations justifies restricting attention in practice

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY FOUR A

Bell's Own Version Of His Theorem Assumed That S And T Are Both Bivalent, Either −1 Or 1, But Other Ranges Are Assumed In Other Variants Of The Theorem

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) The result of an experiment on 1 is labeled by (e) s, which can take on (e) any of a discrete set of real numbers in the interval [−1, 1].
(2) Likewise the result of an experiment on 2 is labeled by (e) t, which can take on (e) any of a discrete set of real numbers in [−1, 1].
(3) Bell's own version of his theorem assumed (eb) that s and t are both bivalent, either −1 or 1, but other ranges are assumed (eb) in other variants of the theorem.
(4) The following probabilities are assumed to be well defined: (5) p1m (s |a, b, t) = the probability

that the outcome of the measurement performed on 1 is s when m is (=) the complete state, the measurements performed on 1 and 2 respectively are (=) a and b, and the result of the experiment on 2 is (=) t; (6) p2m (t |a, b, s) = the probability that the outcome of the measurement performed on 2 is (=) t when the complete state is m, the measurements performed on 1 and 2 are respectively a and b, and (eb) the result of the measurement a is s. (7) pm (s, t |a, b) = the probability that the results of the joint measurements a and b, when the complete state is m, are (=) respectively s and t.

(5) The probability function p will be assumed (eb) to be non-negative and to sum to unity when the summation is taken over all allowed values of s and t. (Note that the hidden variables theories considered in Section 1 can be subsumed under this conceptual framework by restricting (e) the values of the probability function p to 1 and 0, the former being identified with (=) the truth value "true" and the latter to the truth value "false".)

NOTATION

Module One

Result of an experiment on 1 is labeled by s, which can take on (e) any of a discrete set of real numbers in the interval [−1, 1]

G_{13} : Category one of s, which can take on (e) any of a discrete set of real numbers in the interval [−1, 1]

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of Result of an experiment on 1 is labeled

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Result of an experiment on 1 is labeled by s, which can take on (e) any of a discrete set of real numbers in the interval [−1, 1]

G_{16} : Category one of any of a discrete set of real numbers in the interval [−1, 1]

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Result of an experiment on 1 is labeled by s, which can take

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Likewise the result of an experiment on 2 is labeled by t, which can take on (e) any of a discrete set of real numbers in [−1, 1]

G_{20} : Category one of t, which can take on (e) any of a discrete set of real numbers in [−1, 1]

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of result of an experiment on 2 is labeled

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

result of an experiment on 2 is labeled by t, which can take on (e) any of a discrete set of real numbers in $[-1, 1]$

G_{24} : Category one of any of a discrete set of real numbers in $[-1, 1]$

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of result of an experiment on 2 is labeled by t, which can take on

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

The following probabilities are assumed to be well defined: (5) p1m (s |a, b, t) = the probability that the outcome of the measurement performed on 1 is s when m is (=) the complete state, the measurements performed on 1 and 2 respectively are (=) a and b, and the result of the experiment on 2 is (=) t; (6) p2m (t |a, b, s) = the probability that the outcome of the measurement performed on 2 is (=) t when the complete state is m, the measurements performed on 1 and 2 are respectively a and b, and (eb) the result of the measurement a is s. (7) pm (s, t |a, b) = the probability that the results of the joint measurements a and b, when the complete state is m, are (=) respectively s and t.

G_{28} : Category one of p1m (s |a, b, t)

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of probability that the outcome of the measurement performed on 1 is s when m is (=) the complete state, the measurements performed on 1 and 2 respectively are (=) a and b, and the result of the experiment on 2 is (=) t

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

p2m (t |a, b, s) = the probability that the outcome of the measurement performed on 2 is (=) t when the complete state is m, the measurements performed on 1 and 2 are respectively a and b, and the result of the

measurement a is s.

G_{32} : Category one of p2m (t |a, b, s) = the probability that the outcome of the measurement performed on 2

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of t when the complete state is m, the measurements performed on 1 and 2 are respectively a and b, and the result of the measurement a is s.

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

pm (s, t |a, b) = the probability that the results of the joint measurements a and b, when the complete state is m, are (=) respectively s and t.

G_{36} : Category one of complete state is m, are (=) respectively s and t.

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of pm (s, t |a, b) = the probability that the results of the joint measurements a and b

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

probability function p will be assumed to be non-negative and to sum to unity when the summation is taken over all allowed values of s and t. (Note that the hidden variables theories considered in Section 1 can be subsumed under this conceptual framework by restricting (e) the values of the probability function p to 1 and 0, the former being identified with (=) the truth value “true” and the latter to the truth value “false”.)

G_{40} : Category one of probability function p

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of non-negative and to sum to unity when the summation is taken over all allowed values of s and t. (Note that the hidden variables theories considered in Section 1 can be subsumed under this conceptual framework by restricting (e) the values of the probability function p to 1 and 0, the former being identified with (=) the truth value “true” and the latter to the truth value “false”.)

T_{41} : Category two of SAS

T_{42} : Category three of SAS

probability function p will be assumed to be non-negative and to sum to unity when the summation is taken over all allowed values of s and t. (Note that the hidden variables theories considered in Section 1 can be subsumed under this conceptual framework by restricting (e) the values of the probability function p to 1 and 0, the former being identified with (=) the truth value "true" and the latter to the truth value "false".)

G_{44} : Category one of values of the probability function p to 1 and 0, the former being identified with (=) the truth value "true" and the latter to the truth value "false".)

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of probability function p will be assumed to be non-negative and to sum to unity when the summation is taken over all allowed values of s and t. (Note that the hidden variables theories considered in Section 1 can be subsumed under this conceptual framework

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY FIVE

Charginos, Neutralinos ,Sleptons And Supersymmetric Standard Model

INTRODUCTION—VARIABLES USED

The Supersymmetric Standard Model, with a Brout-Englert-Higgs boson as spin-0 partner of the Z P. Fayet

(1) Supersymmetric extensions of the standard model lead (eb) to expect superpartners for all particles, spin-0 squarks and sleptons and spin-1/2 gluinos, charginos and neutralinos, with an odd R-parity making the lightest one stable.
(2) The electroweak breaking is induced by (e) a pair of spin-0 doublets, leading to several charged and neutral BE-Higgs bosons.
(3) These theories also lead to (eb) gauge/Higgs unification by providing (e) spin-0 bosons as extra states for spin-1 gauge bosons within (eb) massive gauge multiplets.
(4) In particular, the 125 GeV boson recently observed at CERN, most likely a BE-Higgs boson associated with the electroweak breaking, may also be interpreted, up to a mixing angle induced by supersymmetry breaking, as (=) the spin-0 partner of the Z under two supersymmetry transformations.
(5) Authors also discuss how the compactification of extra dimensions, relying on R-parity and other discrete symmetries, may determine both the grand-unification and supersymmetry-breaking scales. High Energy Physics - Phenomenology (hep-ph) arXiv: 1403.5951 [hep-ph]

Search for direct production of charginos, neutralinos and sleptons in final states with two leptons and missing transverse momentum in pp collisions at sqrt(s) = 8 TeV with the ATLAS detector

(6) Searches for the electroweak production of charginos, neutralinos and sleptons in final states characterized by (e) the presence of two leptons (electrons and muons) and missing transverse momentum are performed using (e) 20.3 fb-1 of proton-proton collision data at sqrt(s) = 8 TeV recorded with the ATLAS experiment at the Large Hadron Collider. **Here take that many such experiments with variables are performed or equate all the three categories**

(7) No significant excess beyond Standard Model expectations is observed. Limits are set on (eb) the masses of the lightest chargino, next-to-lightest neutralino and sleptons for different lightest-neutralino mass hypotheses in simplified models.

(8) Results are also interpreted in various scenarios of the phenomenological Minimal Supersymmetric Standard Model. High Energy Physics - Experiment (hep-ex) Report number: CERN-PH-EP-2014-037 arXiv: 1403.5294 [hep-ex]

Search for direct top squark pair production in events with a Z boson, b-jets and missing transverse momentum in sqrt(s)=8 TeV pp collisions with the ATLAS detector ATLAS Collaboration

(9) A search is presented for direct top squark pair production using (e) events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent with (e&eb) the Z boson mass, jets tagged as (=) originating from b-quarks and missing transverse momentum.

(10) The analysis is performed with proton-proton collision data at \sqrt{s}=8 TeV collected with the ATLAS detector at the LHC in 2012 corresponding (e&eb) to an integrated luminosity of 20.3 fb^{-1}.

(11) No excess beyond the Standard Model expectation is observed. Interpretations of the results are provided in models based (e) on the direct pair production of the heavier top squark state (\tilde{t}_2) followed by the decay to (eb) the lighter top squark state (\tilde{t}_1) via \tilde{t}_2 \to Z\tilde{t}_1, and for \tilde{t}_1 pair production in natural gauge-mediated supersymmetry breaking scenarios where the neutralino (\tilde{\chi}^0_1) is (=) the next-to-lightest supersymmetric particle and decays producing (eb) a Z boson and a gravitino (\tilde{G}) via the \tilde{\chi}^0_1 \to Z\tilde{G} process. Report number: CERN-PH-EP-2014-035 arXiv: 1403.5222 [hep-ex]

NOTATION

Module One

Supersymmetric extensions of the standard model lead (eb) to expect superpartners for all particles, spin-0 squarks and sleptons and spin-1/2 gluinos, charginos and neutralinos, with an odd R-parity making the lightest one stable.

G_{13} : Category one of Supersymmetric extensions of the standard model

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of expect superpartners for all particles, spin-0 squarks and sleptons and spin-1/2 gluinos, charginos and neutralinos, with an odd R-parity making the lightest one stable.

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

Electroweak breaking is induced by a pair of spin-0 doublets, leading to several charged and neutral BE-Higgs bosons.

G_{16} : Category one of pair of spin-0 doublets, leading to several charged and neutral BE-Higgs bosons.

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of Electroweak breaking

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Electroweak breaking is induced by a pair of spin-0 doublets, leading to several charged and neutral BE-Higgs bosons.

G_{20} : Category one of Electroweak breaking is induced by a pair of spin-0 doublets

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of several charged and neutral BE-Higgs bosons.

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

These theories also lead to (eb) gauge/Higgs unification by providing (e) spin-0 bosons as extra states for spin-1 gauge bosons within (eb) massive gauge multiplets.

G_{24} : Category one of gauge; Higgs

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of Higgs; gauge

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

gauge/Higgs unification is possible by providing (e) spin-0 bosons as extra states for spin-1 gauge bosons

within (eb) massive gauge multiplets

G_{28} : Category one of spin-0 bosons as extra states for spin-1 gauge bosons within (eb) massive gauge multiplets

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of gauge/Higgs unification is possible

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

125 GeV boson recently observed at CERN, most likely a BE-Higgs boson associated with the electroweak breaking, may also be interpreted, up to a mixing angle induced by supersymmetry breaking, as (=) the spin-0 partner of the Z under two supersymmetry transformations.

G_{32} : Category one of125 GeV boson recently observed at CERN, most likely a BE-Higgs boson associated with the electroweak breaking, may also be interpreted, up to a mixing angle induced by supersymmetry breaking (Take many such experiments in to consideration or alternatively equate all the categories to each other)

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of spin-0 partner of the Z under two supersymmetry transformations.

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Searches for the electroweak production of charginos, neutralinos and sleptons in final states characterized by (e) the presence of two leptons (electrons and muons) and missing transverse momentum are performed using (e) 20.3 fb-1 of proton-proton collision data at sqrt(s) = 8 TeV recorded with the ATLAS experiment at the Large Hadron Collider. **Here take that many such experiments with variables are performed or equate all the three categories**

G_{36} : Category one ofpresence of two leptons (electrons and muons) and missing transverse momentum are performed using (e) 20.3 fb-1 of proton-proton collision data at sqrt(s) = 8 TeV recorded with the ATLAS experiment at the Large Hadron Collider. **Here take that many such experiments with variables are performed or equate all the three categories**

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Searches for the electroweak production of charginos, neutralinos and sleptons in final states

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

A search is presented for direct top squark pair production using events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent with (e&eb) the Z boson mass, jets tagged as (=) originating from b-quarks and missing transverse momentum

G_{40} : Category one of events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent with (e&eb) the Z boson mass, jets tagged as (=) originating from b-quarks and missing transverse momentum

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of search is presented for direct top squark pair production

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Dual Modules : care

search is presented for direct top squark pair production using events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent with the Z boson mass, jets tagged as (=) originating from b-quarks and missing transverse momentum

G_{44} : Category one of search is presented for direct top squark pair production using events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent; Z boson mass, jets tagged as (=) originating from b-quarks and missing transverse momentum; search is presented for direct top squark pair production using events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent with the Z boson mass, jets tagged

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Z boson mass, jets tagged as (=) originating from b-quarks and missing transverse momentum; search is presented for direct top squark pair production using events with at least two leptons including a same-flavour opposite-sign pair with invariant mass consistent; originating from b-quarks and missing transverse momentum

T_{45} : Category two of SAS

T_{46} : Category three of SAS

SECTION THIRTY SIX

Embodiment And Expression Of Human Values Is Will To Power.

INTRODUCTION—VARIABLES USED

Friedrich Nietzsche (1844—1900): **Wikipedia**

(1) Here, apparently, Nietzsche's doctrine of the feeling of power has become (=) more than an observation on the natural history and psychology of morals.
(2) The "teaching" reaches into (eb) the heart of life and it says something absolute about obeying and commanding.
(3) But what is being obeyed, on the cosmological level, and what is being commanded? At this point, Zarathustra passes on a secret told to him by life itself: "behold [life says], I am that which must overcome (e) itself again and again…
(4) And you too, enlightened man, are (=) only a path and a footstep of my will: truly, my will to power walks with (e&eb) the feet of your will to truth."
(5) We see here that a principle, will to power, is embodied by (e) the human being's will to truth, and we may imagine it taking other forms as well.
(6) Reflecting on this insight, for example, Zarathustra claims to have solved (e) "the riddle of the hearts" of the creator of values:
(7) "You exert power with your values and doctrines of good and evil, you assessors of values….but a mightier power and a new overcoming grow from out of (e) your values…"
(8) That mightier power growing in and through the embodiment and expression of human values is (=) will to power.
(9) It is important not to disassociate will to (e&eb) power, as a cosmology, from the human being's drive (eb) to create values.

NOTATION

Module One

Nietzsche's doctrine of the feeling of power has become (=) more than an observation on the natural history and psychology of morals

G_{13} : Category one of doctrine of the feeling of power

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of more than an observation on the natural history and psychology of morals

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

"Teaching" reaches into (eb) the heart of life and it says something absolute about obeying and commanding.

G_{16} : Category one of "teaching"

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of heart of life and it says something absolute about obeying and commanding

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

But what is being obeyed, on the cosmological level, and what is being commanded? At this point, Zarathustra passes on a secret told to him by life itself: "behold [life says], I am that which must overcome (e) itself again and again…

G_{20} : Category one of itself again and again

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **behold [life says], I am that which must overcome**

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

And you too, enlightened man, are only a path and a footstep of my will: truly, my will to power walks with (e&eb) the feet of your will to truth."

G_{24} : Category one of path and a footstep of my will: truly, my will to power walks; feet of your will to truth."

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of feet of your will to truth."; path and a footstep of my will: truly, my will to power walks

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Will to power, is embodied by (e) the human being's will to truth, and we may imagine it taking other forms as well.

G_{28} : Category one of human being's will to truth, and we may imagine it taking other forms as well.

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Will to power

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

You exert power with your values and doctrines of good and evil, you assessors of values….but a mightier power and a new overcoming grow from out of (e) your values…"

G_{32} : Category one of exert power with your values and doctrines of good and evil

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of mightier power and a new overcoming grow from out of (e) your values…"

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Exert power with your values and doctrines of good and evil, you assessors of values….but a mightier power and a new overcoming grow from out of (e) your values…" (emphasis on the relativisticness of good and bad)

G_{36} : Category one of your values

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of Exert power with your values and doctrines of good and evil, you assessors of values….but a mightier power and a new overcoming grow

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

Mightier power growing in and through the embodiment and expression of human values is (=) will to power.

G_{40} : Category one of Mightier power growing in and through the embodiment and expression of human values

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of will to power.

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

It is important not to disassociate will to power, as a cosmology, from the human being's drive (eb) to create values.

G_{44} : Category one of will; power

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of power; will

T_{45} : Category two of SAS

T_{46} : Category three of SAS

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}\ (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$:
$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$,
$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, \ (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$
$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}\ (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$,
$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation

coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) & \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad 73$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad 74$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad 76$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)} & \boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & \boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & \boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & \boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & \boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} \quad 85$$

$$- \begin{bmatrix} (a'_{32})^{(6)} \boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \begin{bmatrix} (a'_{33})^{(6)} \boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \begin{bmatrix} (a'_{34})^{(6)} \boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34} \quad 87$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \begin{bmatrix} (b'_{32})^{(6)} \boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \begin{bmatrix} (b'_{33})^{(6)} \boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & \boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \begin{bmatrix} (b'_{40})^{(8)}\,-(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \begin{bmatrix} (b'_{41})^{(8)}\,-(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \begin{bmatrix} (b'_{42})^{(8)}\,-(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} - \begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} - \begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq \ (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = \ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \qquad 102$$

$$(b_i'')^{(2)}(G_{19},t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t) - (a_i'')^{(2)}(T_{17},t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - \ T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t) - (b_i'')^{(2)}\big((G_{19}),t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\big[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\big]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t) \le (p_i)^{(3)} \le (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t) \le (r_i)^{(3)} \le (b_i')^{(3)} \le (\hat{B}_{20})^{(3)}$$

$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t) = (p_i)^{(3)}$ 113

$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t) - (a_i'')^{(3)}(T_{21},t)| \le (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}',t) - (b_i'')^{(3)}(G_{23},t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big) \leq \ (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}\big((G_{27}),t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \ , \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and \ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big) \leq \ (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = \ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - \ T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}\big((G_{31}),t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\big((G_{35}),t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}\big((G_{35}),t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and \ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(YYYYYYYYYY) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

(ZZZZZZZZZZ) The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

(AAAAAAAAAAA) $\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$

(BBBBBBBBBBB)

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$

$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

(CCCCCCCCCCC) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

(DDDDDDDDDDD) There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42$,
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}\left[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}\right] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \ , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\overline{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^{0} + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy A

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\tilde{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\tilde{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left(e^{(\tilde{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\tilde{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\tilde{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \quad (169)$$

$$\left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(rr) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^{0}+\int_0^t\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)} = \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for G_{41} , $G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for G_{45} , $G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}$, $\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded.
The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1$, $\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. 205
The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$: $\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)}-G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+\ (a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded.
The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)}-G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24{,}25{,}26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded.
The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\tilde{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)} - G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded.
The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \qquad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded.
The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t),t(t),t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})} \right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{36})^{(7)} \left|G_{37}^{(1)} - G_{37}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}}\, ds_{(36)} +$$
$$\int_0^t \{(a'_{36})^{(7)} \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) \left|G_{36}^{(1)} - G_{36}^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} +$$
$$G_{36}^{(2)} \left|(a''_{36})^{(7)} \left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)} \left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)} s_{(36)}} e^{(\widehat{M}_{36})^{(7)} s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right| e^{-(\widehat{M}_{36})^{(7)} t} \leq \qquad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}\right) d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ *and* $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)} t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded.
The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\ ((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded.
The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded.
The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad (324)$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)},\ \boldsymbol{if}\ (\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},\ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G^0_{28} e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G^0_{28} e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G^0_{28}}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G^0_{30} e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+\ G_{30}^{0}e^{-(a'_{30})^{(5)}t}\Big)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t}\le T_{28}(t)\le\frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t}\le T_{30}(t)\le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)},$ ***if*** $(v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, $\boldsymbol{if}$ $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}$ $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}$ $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}$ $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right) - (b''_{41})^{(8)}\left((G_{43}), t\right) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}$,, $(\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}$ $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}$ $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t})$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\leq T_{40}(t)\leq \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\leq T_{42}(t)\leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)}\leq -(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\leq -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)}\leq -(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\leq -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)},(v_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},v^{(9)},u^{(9)}$:

By $(v_1)^{(9)}>0$, $(v_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)},(m_2)^{(9)},(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)},\ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t} \right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})} [e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}] + G_{46}^0 e^{-(a'_{46})^{(9)} t} \right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t} \right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

For $0 < \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \ge \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}}$$

it follows $(v_0)^{(1)} \le v^{(1)}(t) \le (v_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}}$$

From which we deduce $(v_0)^{(1)} \le v^{(1)}(t) \le (\bar{v}_1)^{(1)}$

If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 385

$$(v_1)^{(1)} \le \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((v_1)^{(1)} - (v_2)^{(1)}\right)t\right]}} \le v^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \le (\bar{v}_1)^{(1)} \le \boxed{(v_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$, we obtain 386

$$(v_1)^{(1)} \le v^{(1)}(t) \le \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}\right)t\right]}} \le (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}\nu^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

406

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case,

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

407

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \le \nu^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(7)} and\ (b''_i)^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(8)} and\ (b''_i)^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(9)} and\ (b''_i)^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\ ]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\ ]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\ ]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(rr) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)} G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)} G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)} T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)} T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^{*}=\frac{(b_{28})^{(5)}T_{29}^{*}}{[(b_{28}')^{(5)}-(b_{28}'')^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*}=\frac{(b_{30})^{(5)}T_{29}^{*}}{[(b_{30}')^{(5)}-(b_{30}'')^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*})=0$, T_{33}^{*} given by $f(T_{33}^{*})=0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a_{32}')^{(6)}+(a_{32}'')^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a_{34}')^{(6)}+(a_{34}'')^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b_{32}')^{(6)}-(b_{32}'')^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b_{34}')^{(6)}-(b_{34}'')^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a_{36}')^{(7)}+(a_{36}'')^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a_{38}')^{(7)}+(a_{38}'')^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b_{36}')^{(7)}-(b_{36}'')^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b_{38}')^{(7)}-(b_{38}'')^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a_{40}')^{(8)}+(a_{40}'')^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a_{42}')^{(8)}+(a_{42}'')^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b_{40}')^{(8)}-(b_{40}'')^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b_{42}')^{(8)}-(b_{42}'')^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a_{44}')^{(9)}+(a_{44}'')^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a_{46}')^{(9)}+(a_{46}'')^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b_{44}')^{(9)}-(b_{44}'')^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b_{46}')^{(9)}-(b_{46}'')^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a_{13}')^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \qquad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a_{14}')^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \qquad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a_{15}')^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \qquad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b_{13}')^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T_{13}^*\mathbb{G}_j\right) \qquad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b_{14}')^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T_{14}^*\mathbb{G}_j\right) \qquad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b_{15}')^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T_{15}^*\mathbb{G}_j\right) \qquad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \qquad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \qquad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \qquad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \qquad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \qquad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \qquad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^{*}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^*_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^*_{25}+(b_{25})^{(4)}s_{(24),(26)}T^*_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^*_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^*_{29}+(b_{29})^{(5)}s_{(28),(29)}T^*_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^*_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^*_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29}+(b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left(\left((\lambda)^{(5)}\right)^2+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29}+(b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\Big]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33}+(b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33}+(b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THIRTY SEVEN

− (¼) ≤ [D(π/8)/D0 − D(3π/8)/D0] ≤ ¼ :Consequence Of The BCHSH Inequality, And In Disagreement With The Quantum Mechanical Prediction

INTRODUCTION—VARIABLES USED

Bell's Inequalities: Stanford Encyclopedia and Wikipedia

(1) The first experimental test of Bell's Inequality, performed by Freedman and Clauser (1972), and proceeded by making two applications of Ineq. (31), one to the angles a = π/4, a′ = 0, b = π/8, b′ = 3π/8, yielding (eb) (32a) $-1 \leq [3D(\pi/8)/D0 - D(3\pi/8)/D0\ ] - D1/D0 - D2/D0 \leq 0$,

(2) Another to the angles a = 3π/4, a′ = 0, b = 3π/8, b′ = 9π/8, yielding (eb) (32b) $-1 \leq [3D(3\pi/8)/D0 - D(\pi/8)/D0\ ] - D1/D0 - D2/D0 \leq 0$.

(3) Combining Ineq. (32a) and Ineq. (32b) yields (eb) (33) **$-(¼) \leq [D(\pi/8)/D0 - D(3\pi/8)/D0\ ] \leq ¼$.**

(4) The Quantum Mechanical prediction for this arrangement, taking into account the uncertainties about the polarization analyzers and the angle from the source subtended by the analyzers, is (=) (34) $[D(\pi/8)/D0 - D(3\pi/8)/D0\ ]QM = (0.401+/-0.005) - (0.100+/-0.005) = 0.301+/-0.007$,

(5) The experimental result obtained by Freedman and Clauser was (35) $[D(\pi/8)/D0 - D(3\pi/8)/D0\ ]expt = 0.300 +/- 0.008$, which is 6.5 sd from the limit allowed by Ineq. (33) But in good agreement with the quantum mechanical prediction Eq. (34).

(6) This was a difficult experiment, requiring 200 hours of running time, much longer than in laser tests of Bell's Inequality, which were able to use (e) lasers for exciting the sources of photon pairs.

(7) Several dozen experiments have been performed to test Bell's Inequalities. References will now be given to some of the most noteworthy of these, along with references to survey articles which provide information about others. A discussion of those actual or proposed experiments which were designed to close two serious loopholes in the early Bell experiments, the "detection loophole" and the "communication loophole". Holt and Pipkin completed in 1973 (Holt 1973) an experiment very much like that of Freedman and Clauser, but examining photon pairs produced in (eb) the 91P1 →73S1→63P 0 cascade in the zero nuclear-spin isotope of mercury-198 after using (e) electron bombardment to pump the atoms to (eb) the first state in this cascade.

(8) The result of Holt and Pipkin was in fairly good agreement with Ineq. (33), which is (=) a **consequence of the BCHSH Inequality, and in disagreement with (e&eb) the quantum mechanical prediction** by nearly 4 sd — contrary to the results of Freedman and Clauser.

(9) Because of the discrepancy between these two early experiments Clauser (1976) repeated the Holt-Pipkin experiment, using (e) the same cascade and excitation method but a different spin-0 isotope of mercury, and his results agreed well with the quantum mechanical predictions but violated (e) the consequence of Bell's Inequality.

NOTATION

Module One

First experimental test of Bell's Inequality, performed by Freedman and Clauser (1972), and proceeded by making two applications of Ineq. (31), one to the angles a = π/4, a′ = 0, b = π/8, b′ = 3π/8, yielding (eb) (32a) $-1 \leq [3D(\pi/8)/D0 - D(3\pi/8)/D0\ ] - D1/D0 - D2/D0 \leq 0$

G_{13} : Category one of **two** applications of Ineq. (31), one to the angles a = π/4, a′ = 0, b = π/8, b′ = 3π/8

G_{14} : Category two of SAS(same as superior/above)

G_{15} : Category three of SAS

T_{13} : Category one of(32a) $-1 \leq [3D(\pi/8)/D0 - D(3\pi/8)/D0\ ] - D1/D0 - D2/D0 \leq 0$

T_{14} : Category two of SAS

T_{15} : Category three of SAS

Module Two

(32a) $-1 \leq [3D(\pi/8)/D0 - D(3\pi/8)/D0\ ] - D1/D0 - D2/D0 \leq 0$

G_{16} : Category one of LHS of the above equation

G_{17} : Category two of SAS

G_{18} : Category three of SAS

T_{16} : Category one of RHS of the above equation

T_{17} : Category two of SAS

T_{18} : Category three of SAS

Module three

Combining Ineq. (32a) and Ineq. (32b) yields (eb) (33) **$-(¼) \leq [D(\pi/8)/D0 - D(3\pi/8)/D0\ ] \leq ¼$.**

G_{20} : Category one of Combining Ineq. (32a) and Ineq. (32b)

G_{21} : Category two of SAS

G_{22} : Category three of SAS

T_{20} : Category one of **$-(¼) \leq [D(\pi/8)/D0 - D(3\pi/8)/D0\ ] \leq ¼$. Note the negative sign**

T_{21} : Category two of SAS

T_{22} : Category three of SAS

Module four

Quantum Mechanical prediction for this arrangement, taking into account the uncertainties about the polarization analyzers and the angle from the source subtended by the analyzers, is (=) (34)
$[D(\pi/8)/D0 - D(3\pi/8)/D0\ ]QM = (0.401+/-0.005) - (0.100+/-0.005) = 0.301+/-0.007$

G_{24} : Category one of Quantum Mechanical prediction for this arrangement, taking into account the uncertainties about the polarization analyzers and the angle from the source subtended by the analyzers

G_{25} : Category two of SAS

G_{26} : Category three of SAS

T_{24} : Category one of (34) $[D(\pi/8)/D0 - D(3\pi/8)/D0\ ]QM = (0.401+/-0.005) - (0.100+/-0.005) = 0.301+/-0.007$

T_{25} : Category two of SAS

T_{26} : Category three of SAS

Module five

Experimental result obtained by Freedman and Clauser was (35) $[D(\pi/8)/D0 - D(3\pi/8)/D0\ ]expt = 0.300 +/-0.008$, which is 6.5 sd from the limit allowed by Ineq. (33) But in good agreement with the quantum mechanical prediction Eq. (34).

G_{28} : Category one of limit allowed by Ineq. (33) But in good agreement with the quantum mechanical prediction Eq. (34).

G_{29} : Category two of SAS

G_{30} : Category three of SAS

T_{28} : Category one of Experimental result obtained by Freedman and Clauser was (35) [D(π/8)/D0 − D(3π/8)/D0]expt = 0.300 +/- 0.008, which is 6.5 sd

T_{29} : Category two of SAS

T_{30} : Category three of SAS

Module six

Result of Holt and Pipkin was in fairly good agreement with Ineq. (33), which is a **consequence of the BCHSH Inequality, and in disagreement with (e&eb) the quantum mechanical prediction** by nearly 4 sd — contrary to the results of Freedman and Clauser.

G_{32} : Category one of Result of Holt and Pipkin was in fairly good agreement with Ineq. (33) Take in to consideration various similar experiments or equate all the three classification categories which are based upon the characterstics of the system based on the satisfaction of the predications of the given system.

G_{33} : Category two of SAS

G_{34} : Category three of SAS

T_{32} : Category one of **consequence of the BCHSH Inequality, and in disagreement with (e&eb) the quantum mechanical prediction** by nearly 4 sd — contrary to the results of Freedman and Clauser.

T_{33} : Category two of SAS

T_{34} : Category three of SAS

Module seven

Result of Holt and Pipkin was in fairly good agreement with Ineq. (33), which is a **consequence of the BCHSH Inequality, and in disagreement with (e&eb) the quantum mechanical prediction** by nearly 4 sd — contrary to the results of Freedman and Clauser.

G_{36} : Category one of **BCHSH Inequality; quantum mechanical prediction** by nearly 4 sd — contrary to the results of Freedman and Clauser.

G_{37} : Category two of SAS

G_{38} : Category three of SAS

T_{36} : Category one of **quantum mechanical prediction** by nearly 4 sd — contrary to the results of Freedman and Clauser; **BCHSH Inequality**

T_{37} : Category two of SAS

T_{38} : Category three of SAS

Module eight

91P1 →73S1→63P 0 cascade in the zero nuclear-spin isotope of mercury-198 after using (e) electron bombardment to pump the atoms to (eb) the first state in this cascade

G_{40} : Category one of electron bombardment to pump the atoms to (eb) the first state in this cascade

G_{41} : Category two of SAS

G_{42} : Category three of SAS

T_{40} : Category one of91P1 →73S1→63P 0 cascade in the zero nuclear-spin isotope of mercury-198

T_{41} : Category two of SAS

T_{42} : Category three of SAS

Module Nine

Because of the discrepancy between these two early experiments Clauser (1976) repeated the Holt-Pipkin experiment, using the same cascade and excitation method but a different spin-0 isotope of mercury, and his results agreed well with the quantum mechanical predictions but violated (e) the consequence of Bell's Inequality.

G_{44} : Category one of same cascade and excitation method but a different spin-0 isotope of mercury, and his results agreed well with the quantum mechanical predictions; consequence of Bell's Inequality

G_{45} : Category two of SAS

G_{46} : Category three of SAS

T_{44} : Category one of Holt-Pipkin experiment was repeated ; Holt-Pipkin experiment was repeated

T_{45} : Category two of SAS

T_{46} : Category three of SAS

References:

(1) Jump up ^ Shultis, J.K. and Faw, R.E. (2002). Fundamentals of nuclear science and engineering. CRC Press. p. 151. ISBN 0-8247-0834-2.
(2) Jump up ^ Physics Flexbook. Ck12.org. Retrieved on 2012-12-19.
(3) Jump up ^ Bethe, Hans A. "The Hydrogen Bomb", Bulletin of the Atomic Scientists, April 1950, p. 99.

(4) Jump up ^ "Progress in Fusion". ITER. Retrieved 2010-02-15.
(5) Jump up ^ "ITER - the way to new energy". ITER. 2014.
(6) Jump up ^ "The National Ignition Facility: Ushering in a New Age for Science". National Ignition Facility. Retrieved 2009-09-13.
(7) Jump up ^ "DOE looks again at inertial fusion as potential clean-energy source", David Kramer, Physics Today, March 2011, p 26
(8) Jump up ^ The Most Tightly Bound Nuclei. Hyperphysics.phy-astr.gsu.edu. Retrieved on 2011-08-17.
(9) Jump up ^ http://www.scienceworldreport.com/articles/5763/20130323/lawson-criteria-make-fusion-power-viable-iter.htm
(10) Jump up ^ Jones, S.E. (1986). "Muon-Catalysed Fusion Revisited". Nature 321 (6066): 127–133. Bibcode:1986Natur.321..127J. doi:10.1038/321127a0.
(11) Jump up ^ Access: Desktop fusion is back on the table: Nature News. Nature.com. Retrieved on 2011-08-17.
(12) Jump up ^ Supplementary methods for "Observation of nuclear fusion driven by a pyroelectric crystal". Main article Naranjo, B.; Gimzewski, J.K.; Putterman, S. (2005). "Observation of nuclear fusion driven by a pyroelectric crystal". Nature 434 (7037): 1115–1117. Bibcode:2005Natur.434.1115N. doi:10.1038/nature03575. PMID 15858570.
(13) Jump up ^ UCLA Crystal Fusion. Rodan.physics.ucla.edu. Retrieved on 2011-08-17.
(14) Jump up ^ Schewe, Phil and Stein, Ben (2005). "Pyrofusion: A Room-Temperature, Palm-Sized Nuclear Fusion Device". Physics News Update 729 (1).
(15) Jump up ^ Coming in out of the cold: nuclear fusion, for real. Christiansciencemonitor.com (2005-06-06). Retrieved on 2011-08-17.
(16) Jump up ^ fusion on the desktop ... really!. MSNBC (2005-04-27). Retrieved on 2011-08-17.
(17) Jump up ^ Gerstner, E. (2009). "Nuclear energy: The hybrid returns". Nature 460 (7251): 25–8. doi:10.1038/460025a. PMID 19571861.
(18) Jump up ^ FusEdWeb | Fusion Education. Fusedweb.pppl.gov (1998-11-09). Retrieved on 2011-08-17.
(19) ^ Jump up to: a b Archived January 3, 2006 at the Wayback Machine. Retrieved on 2012-12-19.
(20) Jump up ^ Subsection 4.7.4c. Kayelaby.npl.co.uk. Retrieved on 2012-12-19.
(21) Jump up ^ Michael A. Nielsen and Isaac L. Chuang (2000). "Quantum Computation and Quantum Information". Cambridge University Press.
(22) Jump up ^ W.Shor, Peter (1995). "Scheme for reducing decoherence in quantum computer memory". AT&T Bell Laboratories.
(23) Jump up ^ D. G. Cory, M. D. Price, W. Maas, E. Knill, R. Laflamme, W. H. Zurek, T. F. Havel and S. S. Somaroo, "Experimental Quantum Error Correction," Phys. Rev. Lett. 81, 2152–2155 (1998), doi:10.1103/PhysRevLett.81.2152
(24) Jump up ^ T. B. Pittman, B. C. Jacobs and J. D. Franson, "Demonstration of quantum error correction using linear optics," Phys. Rev. A 71, 052332 (2005), doi:10.1103/PhysRevA.71.052332
(25) Jump up ^ J. Chiaverini, D. Leibfried, T. Schaetz, M. D. Barrett, R. B. Blakestad, J. Britton, W. M. Itano, J. D. Jost, E. Knill, C. Langer, R. Ozeri and D. J. Wineland, "Realization of quantum error correction," Nature 432, 602-605 (2004), doi:10.1038/nature03074
(26) Jump up ^ P. Schindler, J. T. Barreiro, T. Monz, V. Nebendahl, D. Nigg, M. Chwalla, M. Hennrich and R. Blatt, "Experimental Repetitive Quantum Error Correction," Science 332, 1059-1061 (2011), doi:10.1126/science.1203329
(27) Jump up ^ M. D. Reed, L. DiCarlo, S. E. Nigg, L. Sun, L. Frunzio, S. M. Girvin and R. J. Schoelkopf, "Realization of Three-Qubit Quantum Error Correction with Superconducting Circuits," Nature 482, 382-385 (2012), doi:10.1038/nature10786, arXiv:1109.4948

@@
@@
@@